

# EMBEDDING CYCLE SYSTEMS OF EVEN LENGTH

C. C. Lindner, C. A. Rodger  
Dept. of Algebra, Combinatorics and Analysis  
Auburn University  
Auburn, Alabama 36849-3501  
U.S.A.

and

D. R. Stinson  
Department of Computer Science  
University of Manitoba  
Winnipeg, Manitoba R3T 2N2  
Canada

**Abstract.** We prove that if  $m$  is even then a partial  $m$ -cycle system on  $n$  vertices can be embedded in an  $m$ -cycle system on  $2mn + 1$  vertices.

## 1. Introduction and notation.

Let  $V(G)$  and  $E(G)$  denote the vertex and edge sets of a graph  $G$ , respectively. Let  $Z_n = \{0, 1, \dots, n-1\}$ . Let  $K_n$  and  $K_{x,y}$  be the complete graph and the complete bipartite graph, respectively. An  $m$ -cycle is a simple graph with  $m$  vertices, say  $u_0, \dots, u_{m-1}$  in which the only edges are  $u_0 u_{m-1}$  and the edges joining  $u_i$  to  $u_{i+1}$  (for  $0 \leq i \leq m-2$ ). We represent this cycle by  $(u_0, \dots, u_{m-1})$  or  $(u_0, u_{m-1}, u_{m-2}, \dots, u_1)$  or any cyclic shift of these. A (partial)  $m$ -cycle system is an order pair  $(V, C(m))$  where  $C(m)$  is a set of edge-disjoint  $m$ -cycles which partition (a subset of) the edge set of the complete graph with vertex set  $V$ .

A partial  $m$ -cycle system  $(Z_n, C_1(m))$  is *embedded* in an  $m$ -cycle system  $(Z_v, C_2(m))$  if  $C_1(m) \subseteq C_2(m)$ . A natural problem then is to find as small a value of  $v$  as possible so that every partial  $m$ -cycle on  $n$  vertices can be embedded in an  $m$ -cycle system on  $v$  vertices. The best result to date is Wilson's theorem [8] which shows that all partial  $m$ -cycle systems can be finitely embedded, but the size  $v$  of the  $m$ -cycle system is an exponential function of  $n$ . (Of course, Wilson's result is proved for the embedding of partial graph decompositions in general, not just for  $m$ -cycle systems.) The only other results related to this problem deal with the particular case when  $m$  is odd. A 3-cycle system is more commonly known as a *Steiner triple system*. Originally, a finite embedding of a partial Steiner triple system on  $n$  vertices in a Steiner triple system on  $v$  vertices was found by Treash [7], but  $v$  is an exponential function of  $n$ . Gradually over the years, several results [1, 4] have culminated in reducing

---

C.C. Lindner's research supported by NSF grant DMS-8401263  
D.R. Stinson's research supported by NSERC grant A 9287.

$v$  to at most  $4n + 1$ . Most recently it has been shown [5] that if  $m$  is odd then a partial  $m$ -cycle system on  $n$  vertices can be embedded in an  $m$ -cycle system on at most  $m((m - 2)n(n - 1) + 2n + 1)$  vertices. There it is also shown that a partial weak Steiner  $m$ -cycle system on  $n$  vertices can be embedded in an  $m$ -cycle system on  $m(2n + 1)$  vertices when  $m$  is odd.

Here we consider embedding partial  $m$ -cycle systems when  $m$  is even. This seems to be an easier problem than when  $m$  is odd. We show that a partial  $m$ -cycle system on  $n$  vertices can be embedded in an  $m$ -cycle system on at most  $2mn + 1$  vertices.

Let  $\lambda K_n$  denote the graph on  $n$  vertices in which each pair of vertices is joined by exactly  $\lambda$  edges. Some results have been obtained on the generalized embedding problem when  $K_n$  is replaced by  $\lambda K_n$ . However, using the technique described in [3], it can be shown that if any partial  $m$ -cycle system of  $K_n$  can be embedded in an  $m$ -cycle system of  $K_{f(n)}$  then any partial  $m$ -cycle system of  $\lambda K_n$  can be embedded in an  $m$ -cycle system of  $\lambda K_v$ , where  $v \leq f(m(m - 2)\lambda n^2)$ . So, clearly, the case when  $\lambda = 1$  is of most interest.

All graphs in this paper are simple. See [2] for any graph theoretical terms that are not defined. Throughout the rest of this paper we shall assume that  $m$  is even and at least 4; we shall write  $m = 2k$ . Let  $[x]$  denote the greatest integer less than or equal to  $x$ .

## 2. Preliminary results.

In this paper, we make extensive use of the following result. We say that a graph  $G$  can be *decomposed* into  $m$ -cycles if there exists a set of  $m$ -cycles  $C(m)$ , the edges of which partition  $E(G)$ . Recall that we write  $m = 2k$ .

**Lemma 2.1.** [6]  $K_{x,y}$  can be decomposed into  $m$ -cycles if and only if  $x \geq k$ ,  $y \geq k$ ,  $m$  divides  $xy$  and  $x$  and  $y$  are even.

By Lemma 2.1,  $K_{m,m}$  can be decomposed into  $m$ -cycles; we denote such a set of  $m$ -cycles on the vertex set  $\{i, j\} \times \{0, 1, \dots, m - 1\}$  by  $C(i, j; m)$ .

Two partial  $m$ -cycle systems  $(Z_n, C_1)$  and  $(Z_n, C_2)$  are *mutually balanced* if for all  $ij \in E(K_n)$ ,  $ij$  is in a cycle in  $C_1$  if and only if  $ij$  is in a cycle in  $C_2$ . The following mutually balanced  $m$ -cycle systems, both defined on the vertex set  $Z_m \times Z_m$ , are of vital importance in proving our result. Let

$$A_1(m) = \bigcup_{i=0}^{m-1} C(i, i+1; m),$$

where all vertices are reduced modulo  $m$ , and

$$A_2(m) = \{ ((0, x), (1, x + y), (2, x + 2y), \dots, (m - 1, x + (m - 1)y)) \mid 0 \leq x \leq m - 1, 0 \leq y \leq m - 1 \}.$$

**Lemma 2.2.**  $(Z_m \times Z_m, A_1(m))$  and  $(Z_m \times Z_m, A_2(m))$  are mutually balanced partial  $m$ -cycle systems.

To construct  $m$ -cycle systems we need the following result. Again recall that  $m = 2k$ .

**Lemma 2.3.** *There exists an  $m$ -cycle system on  $2m + 1$  vertices.*

**Proof:** In each of 2 cases, we define an  $m$ -cycle  $a(m) = (a_1, a_2, \dots, a_m)$  as follows.

Let  $m = 4x$ . For  $1 \leq i \leq m/2$ , define

$$a_i = (-1)^i i, \text{ and}$$

$$a_{(m/2)+i} = \begin{cases} (m/2) - 2 + i & \text{if } i \text{ is odd,} \\ (3m/2) - 1 - i & \text{if } i \text{ is even,} \end{cases}$$

where everything is reduced modulo  $2m + 1$ .

Let  $m = 4x + 2$ . Define

$$a_i = (-1)^{i-1} (i - 1) \quad \text{for } 1 \leq i \leq m/2 - 2,$$

$$a_{k-1} = 1 - k, a_k = k - 2,$$

$$a_{k+1} = -k, a_{k+2} = k + 1,$$

$$a_{m-2i} = m - 1 + 2i \quad \text{for } 0 \leq i \leq (m - 6)/4, \text{ and}$$

$$a_{m-2i+1} = m - 1 - 2i \quad \text{for } 1 \leq i \leq (m - 6)/4,$$

where everything is reduced modulo  $2m + 1$ .

Now define  $a(m) + i$  to be the  $m$ -cycle formed by adding  $i$  modulo  $2m + 1$  to each vertex in  $a(m)$ . Then  $C(m) = \{a(m) + i \mid 0 \leq i \leq 2m\}$  is the required set of  $m$ -cycles.

**Example 2.4:**  $a(8) = (16, 2, 14, 4, 3, 9, 5, 7)$  and  $a(10) = (0, 20, 2, 17, 3, 16, 6, 11, 7, 9)$ .

Finally, we conclude this section with a construction of some  $m$ -cycle systems. Let  $(\{\infty\} \cup (\{i, j\} \times Z_m), B(i, j; m))$  be an  $m$ -cycle system (which exists by Lemma 2.3). Let  $E = \{(i, i + n) \mid 0 \leq i \leq n - 1\}$ .

**Theorem 2.5.** *For any  $n \geq 1$ ,  $(\{\infty\} \cup (Z_{2n} \times Z_m), D(n, m))$  is an  $m$ -cycle system, where we define*

$$D(n, m) = \left( \bigcup_{i=0}^{n-1} B(i, i + n; m) \right) \cup \left( \bigcup_{\substack{0 \leq i < j \leq 2n-1 \\ (i, j) \notin E}} C(i, j; m) \right).$$

### 3. Embedding partial $m$ -cycle systems.

**Theorem 3.1.** *A partial  $m$ -cycle system on  $n$  vertices can be embedded in an  $m$ -cycle system on  $2mn + 1$  vertices.*

Proof: Let  $(Z_n, C_1(m))$  be a partial  $m$ -cycle system. Let  $(\{\infty\} \cup (Z_{2n} \times Z_m), D(n, m))$  be the  $m$ -cycle system constructed in Theorem 2.5. For each  $m$ -cycle  $u = (u_0, u_1, \dots, u_{m-1}) \in C_1(m)$  let  $A_2(u; m)$  be the  $m$ -cycles formed from those in  $A_2(m)$  by replacing each vertex  $i$  with  $u_i$ . Then by Lemma 2.2,  $\bigcup_{i=0}^{m-1} C(u_i, u_{i+1}; m)$  (of course, reducing the subscripts of  $u_i$  modulo  $m$ ) and  $A_2(u; m)$  form mutually balanced partial  $m$ -cycle systems. For each  $u \in C_1(m)$ , remove the  $m$ -cycles in  $\bigcup_{u=0}^{m-1} C(u_i, u_{i+1}; m)$  from  $D(n, m)$  and replace them with the  $m$ -cycles in  $A_2(u; m)$ , thus forming another  $m$ -cycle system  $(\{\infty\} \cup (Z_{2n} \times Z_m), C_2(m))$ . Then since  $((u_0, 0), (u_1, 0), \dots, (u_{m-1}, 0)) \in A_2(u; m)$ ,  $(\{\infty\} \cup (Z_{2n} \times Z_m), C_2(m))$  is the required embedding of  $(Z_n, C_1(m))$ .

Remark: Clearly, Theorem 3.1 can be strengthened in several ways. For example, a partial  $m$ -cycle system on  $n$  vertices can be embedded in an  $m$ -cycle system on  $2mt + 1$  vertices for any  $t \geq n$ . Also, if  $ij$  is not an edge in any cycle in  $C_1(m)$  then the vertices in  $\{n + i, n + j\} \times Z_m$  need not be introduced in the embedding process, as  $B(i, j; m)$  can be used instead of the two  $m$ -cycle systems  $B(i, i + n; m)$  and  $B(j, j + n; m)$  when constructing  $D(n, m)$ . This observation shows that if  $G$  is the graph consisting of the cycles in  $C_1(m)$  and  $\alpha'$  is the maximum number of independent edges in the complement of  $G$  then  $(Z_n, C_1(m))$  can be embedded in an  $m$ -cycle system on  $1 + 2m(n - \alpha')$  vertices.

## References

1. L.D. Andersen, A.J.W. Hilton and E. Mendelsohn, *Embedding partial Steiner triple systems*, J. London Math. Soc. **41** (1980), 557-576.
2. J.A. Bondy and U.S.R. Murty, *Graph theory with applications*, North-Holland, New York. (1976).
3. C.J. Colbourn, R.C. Hamm, C.C. Lindner and C.A. Rodger, *Embedding partial graph designs, block designs and triple systems with  $\lambda > 1$* , Canad. Math. Bull. **29** (1986), 385-391.
4. C.C. Lindner, *A partial Steiner triple system of order  $n$  can be embedded in a Steiner triple system of order  $6n + 3$* , J. Combinatorial Theory (A), **18** (1975), 349-351.
5. C.C. Lindner, C.A. Rodger and D.R. Stinson, *Small embeddings for partial cycle systems of odd length*, Discrete Math. (to appear).
6. D. Sotheau, *Decomposition of  $K_{m,n}$  ( $K_{m,n}^*$ ) into cycles (circuits) of length  $2k$* , J. Combinatorial Theory (B)**30** (1981), 75-81.
7. C. Treash, *The completion of finite incomplete Steiner triple systems with applications to loop theory*, J. Combinatorial Theory (A)**10** (1971), 259-265.
8. R.M. Wilson, *Construction and uses of pairwise balanced designs*, Math. Centre Tracts **55** (1974), 18-41.