

8

Quantum channel capacities

This chapter is focused on *capacities* of quantum channels for transmitting information. The notion of a channel capacity has multiple, inequivalent formulations in the quantum setting. For example, one may consider the capacity with which classical or quantum information can be transmitted through a channel, and different resources may be available to assist with the information transmission, such as entanglement shared between a sender and receiver before the information transmission takes place.

Three fundamental theorems are presented, characterizing the capacities of quantum channels to transmit either classical or quantum information, both with and without the assistance of prior shared entanglement. When prior shared entanglement between the sender and receiver is not available, these characterizations have a somewhat undesirable property: they require a *regularization*—or an averaging over an increasingly large number of uses of a given channel—and fail to provide capacity formulas that are either explicit or efficiently computable for this reason. The apparent need for such regularizations is discussed in the last section of the chapter, along with the related phenomenon of *super-activation* of quantum capacity.

8.1 Classical information over quantum channels

The general scenario to be considered throughout this chapter involves two hypothetical individuals: a *sender* and a *receiver*. The sender attempts to transmit information, either classical or quantum, to the receiver through multiple, independent uses of a given channel Φ . Schemes are considered in which the sender prepares an input to these channel uses and the receiver processes the output in such a way that information is transmitted with a high degree of accuracy. As is standard in information theory, the chapter mainly deals with the asymptotic regime, making use of entropic notions

to analyze rates of information transmission in the limit of an increasingly large number of independent channel uses.

The subject of the present section is the capacity of quantum channels to transmit *classical* information, including both the case in which the sender and receiver share prior entanglement and in which they do not. The first subsection below introduces notions and terminology concerning channel capacities that will be needed throughout the section, as well as in later parts of the chapter. The second subsection is devoted to a proof of the *Holevo–Schumacher–Westmoreland theorem*, which characterizes the capacity of a channel to transmit classical information without the use of prior shared entanglement. The final subsection proves the *entanglement-assisted capacity theorem*, which characterizes the capacity of a channel to transmit classical information with the assistance of prior shared entanglement.

8.1.1 Classical capacities of quantum channels

Five quantities that relate to the information-transmitting capabilities of channels are defined below. The first two quantities—the *classical capacity* and the *entanglement-assisted classical capacity*—are fundamental within the subject of quantum channel capacities. The remaining three quantities are the *Holevo capacity*, the *entanglement-assisted Holevo capacity*, and the *coherent information*, all of which play important roles in the main results to be presented.

The classical capacity of a channel

Intuitively (and somewhat informally) speaking, the classical capacity of a channel describes the average number of classical bits of information that can be transmitted, with a high degree of accuracy, through each use of that channel. As is typical for information-theoretic notions, channel capacities are more formally defined in terms of asymptotic behaviors, where the limit of an increasing number of channel uses is considered.

When stating a precise mathematical definition of classical capacity, it is convenient to refer to the *emulation* of one channel by another.

Definition 8.1 Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ and $\Psi \in C(\mathcal{Z})$ be channels, for \mathcal{X} , \mathcal{Y} , and \mathcal{Z} being complex Euclidean spaces. It is said that the channel Φ *emulates* Ψ if there exist channels $\Xi_E \in C(\mathcal{Z}, \mathcal{X})$ and $\Xi_D \in C(\mathcal{Y}, \mathcal{Z})$ such that

$$\Psi = \Xi_D \Phi \Xi_E. \quad (8.1)$$

When this relationship holds, the channel Ξ_E is called an *encoding channel* and Ξ_D is called a *decoding channel*.

It is also convenient to refer to an *approximation* of a given channel by another. In this chapter, such an approximation is always assumed to be defined with respect to the completely bounded trace norm.

Definition 8.2 Let $\Psi_0, \Psi_1 \in C(\mathcal{Z})$ be channels, for \mathcal{Z} being a complex Euclidean space, and let $\varepsilon > 0$ be a positive real number. The channel Ψ_0 is an ε -approximation to Ψ_1 (equivalently, Ψ_1 is an ε -approximation to Ψ_0) if

$$\|\|\Psi_0 - \Psi_1\|\|_1 < \varepsilon. \quad (8.2)$$

The definition of the classical capacity of a channel, which makes use of the previous two definitions, is as follows.

Definition 8.3 (Classical capacity of a channel) Let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces and let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel. Let $\Gamma = \{0, 1\}$ denote the binary alphabet, let $\mathcal{Z} = \mathbb{C}^\Gamma$, and let $\Delta \in C(\mathcal{Z})$ denote the completely dephasing channel defined with respect to the space \mathcal{Z} .

1. A value $\alpha \geq 0$ is an *achievable rate* for classical information transmission through Φ if (i) $\alpha = 0$, or (ii) $\alpha > 0$ and the following holds for every positive real number $\varepsilon > 0$: for all but finitely many positive integers n , and for $m = \lfloor \alpha n \rfloor$, the channel $\Phi^{\otimes n}$ emulates an ε -approximation to the channel $\Delta^{\otimes m}$.
2. The *classical capacity* of Φ , denoted $C(\Phi)$, is the supremum value of all achievable rates for classical information transmission through Φ .

In the context of Definition 8.3, the completely dephasing channel Δ is to be viewed as an ideal channel for transmitting a single bit of classical information. When considering an emulation of the m -fold tensor product $\Delta^{\otimes m}$ of this ideal classical channel by the channel $\Phi^{\otimes n}$, no generality is lost in restricting one's attention to classical-to-quantum encoding channels Ξ_E and quantum-to-classical decoding channels Ξ_D . That is, one may assume

$$\Xi_E = \Xi_E \Delta^{\otimes m} \quad \text{and} \quad \Xi_D = \Delta^{\otimes m} \Xi_D. \quad (8.3)$$

This assumption causes no loss of generality because

$$\begin{aligned} & \|\|(\Delta^{\otimes m} \Xi_D) \Phi^{\otimes n} (\Xi_E \Delta^{\otimes m}) - \Delta^{\otimes m}\|\|_1 \\ &= \|\|\Delta^{\otimes m} (\Xi_D \Phi^{\otimes n} \Xi_E - \Delta^{\otimes m}) \Delta^{\otimes m}\|\|_1 \\ &\leq \|\|\Xi_D \Phi^{\otimes n} \Xi_E - \Delta^{\otimes m}\|\|_1; \end{aligned} \quad (8.4)$$

replacing a given choice of Ξ_E and Ξ_D by $\Xi_E \Delta^{\otimes m}$ and $\Delta^{\otimes m} \Xi_D$ will never decrease the quality of the emulation achieved.

In light of this observation, the implicit use of the completely bounded trace norm in Definition 8.3 may appear to be somewhat heavy-handed; an equivalent definition is obtained by requiring that $\Phi^{\otimes n}$ emulates some channel $\Psi \in C(\mathcal{Z}^{\otimes m})$ satisfying

$$\|\|(\Delta^{\otimes m} \Psi)(E_{a_1 \cdots a_m, a_1 \cdots a_m}) - E_{a_1 \cdots a_m, a_1 \cdots a_m}\|\|_1 < \varepsilon, \quad (8.5)$$

which is equivalent to

$$\langle E_{a_1 \cdots a_m, a_1 \cdots a_m}, \Psi(E_{a_1 \cdots a_m, a_1 \cdots a_m}) \rangle > 1 - \frac{\varepsilon}{2}, \quad (8.6)$$

for all $a_1 \cdots a_m \in \Gamma^m$. An interpretation of this requirement is that every string $a_1 \cdots a_m \in \Gamma^m$ is transmitted by Ψ with a probability of error smaller than $\varepsilon/2$.

There is, on the other hand, one benefit to using the stronger notion of channel approximation defined by the completely bounded trace norm in Definition 8.3, which is that it allows the quantum capacity (discussed later in Section 8.2) to be defined in an analogous manner to the classical capacity, simply replacing the dephasing channel Δ by the identity channel $\mathbb{1}_{L(\mathcal{Z})}$. (For the quantum capacity, the completely bounded trace norm provides the most natural notion of channel approximation.)

The following proposition is, perhaps, self-evident, but it is nevertheless worth stating explicitly. The same argument used to prove it may be applied to other notions of capacity as well; there is nothing specific to the classical capacity that is required by the proof.

Proposition 8.4 Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} , and let k be a positive integer. It holds that

$$C(\Phi^{\otimes k}) = k C(\Phi). \quad (8.7)$$

Proof If it is the case that α is an achievable rate for classical information transmission through Φ , then it follows trivially that αk is an achievable rate for classical information transmission through $\Phi^{\otimes k}$. It therefore holds that

$$C(\Phi^{\otimes k}) \geq k C(\Phi). \quad (8.8)$$

Now assume that $\alpha > 0$ is an achievable rate for classical information transmission through $\Phi^{\otimes k}$. For any $\varepsilon > 0$ and all but finitely many positive integers n , the channel $\Phi^{\otimes k \lfloor n/k \rfloor}$ therefore emulates an ε -approximation to $\Delta^{\otimes m}$ for $m = \lfloor \alpha \lfloor n/k \rfloor \rfloor$. It will be proved that $\alpha/k - \delta$ is an achievable rate for classical information transmission through Φ for all $\delta \in (0, \alpha/k)$. For any integer $n \geq k$, the channel $\Phi^{\otimes n}$ trivially emulates any channel emulated

by $\Phi^{\otimes k \lfloor n/k \rfloor}$, and for $\delta \in (0, \alpha/k)$, one has that $\alpha \lfloor n/k \rfloor \geq (\alpha/k - \delta)n$ for all but finitely many positive integers n . It therefore holds, for any $\varepsilon > 0$, and all but finitely many positive integers n , that the channel $\Phi^{\otimes n}$ emulates an ε -approximation to $\Delta^{\otimes m}$ for $m = \lfloor (\alpha/k - \delta)n \rfloor$, implying that $\alpha/k - \delta$ is an achievable rate for classical information transmission through Φ . In the case that $\alpha = 0$, one has that α/k is trivially an achievable rate for classical information transmission through Φ . Taking the supremum over all achievable rates, one finds that

$$C(\Phi) \geq \frac{1}{k} C(\Phi^{\otimes k}), \quad (8.9)$$

which completes the proof. \square

The entanglement-assisted classical capacity of a channel

The entanglement-assisted classical capacity of a channel is defined in a similar way to the classical capacity, except that one assumes the sender and receiver may share any state of their choosing prior to the transmission of information through the channel. (As separable states provide no advantage in this setting, the shared state is generally assumed to be entangled.) The ability of the sender and receiver to share entanglement, as compared with the situation in which they do not, can result in a significant increase in the classical capacity of a quantum channel. For instance, shared entanglement doubles the classical capacity of the identity channel through the use of dense coding (discussed in Section 6.3.1), and an arbitrary (constant-factor) increase is possible for other choices of channels.

A formal definition for the entanglement-assisted classical capacity of a channel requires only a minor change to the definition of the ordinary classical capacity: the definition of an emulation of one channel by another is modified to allow for the existence of a shared state as follows.

Definition 8.5 Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ and $\Psi \in C(\mathcal{Z})$ be channels, for \mathcal{X}, \mathcal{Y} , and \mathcal{Z} being complex Euclidean spaces. The channel Φ *emulates* Ψ *with the assistance of entanglement* if there exists a state $\xi \in D(\mathcal{V} \otimes \mathcal{W})$ and channels $\Xi_E \in C(\mathcal{Z} \otimes \mathcal{V}, \mathcal{X})$ and $\Xi_D \in C(\mathcal{Y} \otimes \mathcal{W}, \mathcal{Z})$, for complex Euclidean spaces \mathcal{V} and \mathcal{W} , such that

$$\Psi(Z) = (\Xi_D(\Phi \Xi_E \otimes \mathbf{1}_{L(\mathcal{W})}))(Z \otimes \xi) \quad (8.10)$$

for all $Z \in L(\mathcal{Z})$. (See Figure 8.1 for an illustration of the channel represented by the right-hand side of this equation.) When this relationship holds, the channel Ξ_E is called an *encoding channel*, Ξ_D is called a *decoding channel*, and ξ is referred to as the *shared state* that assists this emulation.




Figure 8.1 An illustration of the map $Z \mapsto (\Xi_D(\Phi \Xi_E \otimes \mathbf{1}_{L(\mathcal{W})}))(Z \otimes \xi)$ referred to in Definition 8.5.

Aside from the modification represented by the previous definition, the entanglement-assisted classical capacity is defined in an analogous way to the ordinary classical capacity.

Definition 8.6 (Entanglement-assisted classical capacity of a channel) Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} , let $\Gamma = \{0, 1\}$ denote the binary alphabet, let $\mathcal{Z} = \mathbb{C}^\Gamma$, and let $\Delta \in C(\mathcal{Z})$ denote the completely dephasing channel defined with respect to the space \mathcal{Z} .

1. A value $\alpha \geq 0$ is an *achievable rate* for entanglement-assisted classical information transmission through Φ if (i) $\alpha = 0$, or (ii) $\alpha > 0$ and the following holds for every positive real number $\varepsilon > 0$: for all but finitely many positive integers n , and for $m = \lfloor \alpha n \rfloor$, the channel $\Phi^{\otimes n}$ emulates an ε -approximation to $\Delta^{\otimes m}$ with the assistance of entanglement.
2. The *entanglement-assisted classical capacity* of Φ , denoted $C_E(\Phi)$, is the supremum over all achievable rates for entanglement-assisted classical information transmission through Φ .

Through the same argument used to prove Proposition 8.4, one has that the following simple proposition holds.

Proposition 8.7 Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} , and let k be a positive integer. It holds that

$$C_E(\Phi^{\otimes k}) = k C_E(\Phi). \quad (8.11)$$

The Holevo capacity of a channel

Suppose that \mathcal{X} is a complex Euclidean space, Σ is an alphabet, $p \in \mathcal{P}(\Sigma)$ is a probability vector, and $\{\rho_a : a \in \Sigma\} \subseteq D(\mathcal{X})$ is a collection of states. Letting $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$ be the ensemble defined as

$$\eta(a) = p(a)\rho_a \quad (8.12)$$

for each $a \in \Sigma$, one has that the Holevo information of η is given by

$$\chi(\eta) = \mathbb{H}\left(\sum_{a \in \Sigma} p(a)\rho_a\right) - \sum_{a \in \Sigma} p(a) \mathbb{H}(\rho_a). \quad (8.13)$$

Based on this quantity, one may define the *Holevo capacity* of a channel in the manner specified by Definition 8.8 below. This definition will make use of the following notation: for any ensemble $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$ and any channel $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$, one defines the ensemble $\Phi(\eta) : \Sigma \rightarrow \text{Pos}(\mathcal{Y})$ as

$$(\Phi(\eta))(a) = \Phi(\eta(a)) \quad (8.14)$$

for each $a \in \Sigma$. That is, $\Phi(\eta)$ is the ensemble obtained by evaluating Φ on the ensemble η in the most natural way.

Definition 8.8 Let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ be a channel, for \mathcal{X} and \mathcal{Y} being complex Euclidean spaces. The *Holevo capacity* of Φ is defined as

$$\chi(\Phi) = \sup_{\eta} \chi(\Phi(\eta)), \quad (8.15)$$

where the supremum is over all choices of an alphabet Σ and an ensemble of the form $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$.

Two restrictions may be placed on the supremum (8.15) in Definition 8.8 without decreasing the value that is defined for a given channel. The first restriction is that the supremum may be replaced by a maximum over all ensembles of the form $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$, for Σ being an alphabet of size

$$|\Sigma| = \dim(\mathcal{X})^2. \quad (8.16)$$

Second, the ensembles may be restricted to ones for which $\text{rank}(\eta(a)) \leq 1$ for each $a \in \Sigma$. The following proposition is useful for proving that this is so.

Proposition 8.9 Let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} , let Σ be an alphabet, and let $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$ be an ensemble. There exists an alphabet Γ and an ensemble $\theta : \Gamma \rightarrow \text{Pos}(\mathcal{X})$ such that

1. $\text{rank}(\theta(b)) \leq 1$ for each $b \in \Gamma$, and
2. $\chi(\Phi(\eta)) \leq \chi(\Phi(\theta))$.

Proof Assume that Λ is the alphabet for which $\mathcal{X} = \mathbb{C}^\Lambda$, and let

$$\eta(a) = \sum_{b \in \Lambda} \lambda_{a,b} x_{a,b} x_{a,b}^* \quad (8.17)$$

be a spectral decomposition of $\eta(a)$ for each $a \in \Sigma$. The requirements of the proposition hold for the ensemble $\theta : \Sigma \times \Lambda \rightarrow \text{Pos}(\mathcal{X})$ defined by

$$\theta(a, b) = \lambda_{a,b} x_{a,b} x_{a,b}^* \quad (8.18)$$

for each $(a, b) \in \Sigma \times \Lambda$. It is evident that the first property holds, so it remains to verify the second.

Define $\mathcal{Z} = \mathbb{C}^\Sigma$ and $\mathcal{W} = \mathbb{C}^\Lambda$, and consider three registers \mathcal{Y} , \mathcal{Z} , and \mathcal{W} corresponding to the spaces \mathcal{Y} , \mathcal{Z} , and \mathcal{W} , respectively. For the density operator $\rho \in \mathcal{D}(\mathcal{Y} \otimes \mathcal{Z} \otimes \mathcal{W})$ defined as

$$\rho = \sum_{(a,b) \in \Sigma \times \Lambda} \lambda_{a,b} \Phi(x_{a,b} x_{a,b}^*) \otimes E_{a,a} \otimes E_{b,b}, \quad (8.19)$$

one has that the following two equalities hold:

$$\begin{aligned} \chi(\Phi(\theta)) &= \mathcal{D}(\rho[\mathcal{Y}, \mathcal{Z}, \mathcal{W}] \parallel \rho[\mathcal{Y}] \otimes \rho[\mathcal{Z}, \mathcal{W}]), \\ \chi(\Phi(\eta)) &= \mathcal{D}(\rho[\mathcal{Y}, \mathcal{Z}] \parallel \rho[\mathcal{Y}] \otimes \rho[\mathcal{Z}]). \end{aligned} \quad (8.20)$$

The inequality $\chi(\Phi(\eta)) \leq \chi(\Phi(\theta))$ follows from the monotonicity of the quantum relative entropy function under partial tracing (which represents a special case of Theorem 5.35). \square

Theorem 8.10 Let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces, let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ be a channel, and let Σ be an alphabet having size $|\Sigma| = \dim(\mathcal{X})^2$. There exists an ensemble $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$ such that

$$\chi(\Phi(\eta)) = \chi(\Phi). \quad (8.21)$$

One may assume, in addition, that $\text{rank}(\eta(a)) \leq 1$ for each $a \in \Sigma$.

Proof Consider an arbitrary ensemble of the form $\theta : \Gamma \rightarrow \text{Pos}(\mathcal{X})$, for Γ being any alphabet, and let

$$\sigma = \sum_{a \in \Gamma} \theta(a) \quad (8.22)$$

denote the average state of the ensemble θ . Through Proposition 2.52, one finds that there must exist an alphabet Λ , a probability vector $p \in \mathcal{P}(\Lambda)$, and a collection of ensembles $\{\theta_b : b \in \Lambda\}$ taking the form $\theta_b : \Gamma \rightarrow \text{Pos}(\mathcal{X})$, each satisfying the constraint

$$\sum_{a \in \Gamma} \theta_b(a) = \sigma \quad (8.23)$$

and possessing the property

$$|\{a \in \Gamma : \theta_b(a) \neq 0\}| \leq \dim(\mathcal{X})^2, \quad (8.24)$$

so that θ is given by the convex combination

$$\theta = \sum_{b \in \Lambda} p(b) \theta_b. \quad (8.25)$$

By Proposition 5.48 it follows that

$$\chi(\Phi(\theta)) \leq \sum_{b \in \Lambda} p(b) \chi(\Phi(\theta_b)), \quad (8.26)$$

and so there must exist at least one choice of a symbol $b \in \Lambda$ for which $p(b) > 0$ and

$$\chi(\Phi(\theta)) \leq \chi(\Phi(\theta_b)). \quad (8.27)$$

Fix any such choice of $b \in \Lambda$, and let

$$\Gamma_0 = \{a \in \Gamma : \theta_b(a) \neq 0\}. \quad (8.28)$$

For an arbitrarily chosen injective mapping $f : \Gamma_0 \rightarrow \Sigma$, one obtains an ensemble $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$ such that

$$\chi(\Phi(\eta)) \geq \chi(\Phi(\theta)) \quad (8.29)$$

by setting $\eta(f(a)) = \theta_b(a)$ for every $a \in \Gamma_0$ and $\eta(c) = 0$ for $c \notin f(\Gamma_0)$.

Because the argument just presented holds for an arbitrary choice of an ensemble θ , it follows that

$$\chi(\Phi) = \sup_{\eta} \chi(\Phi(\eta)), \quad (8.30)$$

where the supremum is over all ensembles of the form $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$. As the set of all such ensembles is compact, there must exist an ensemble of the same form for which the equality (8.21) holds.

The additional restriction that $\text{rank}(\eta(a)) \leq 1$ for each $a \in \Sigma$ may be assumed by first using Proposition 8.9 to replace a given ensemble θ by one satisfying the restriction $\text{rank}(\theta(a)) \leq 1$ for each $a \in \Gamma$, and then proceeding with the argument above. This results in an ensemble $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$ with $\text{rank}(\eta(a)) \leq 1$ for each $a \in \Sigma$, and such that (8.21) holds, which completes the proof. \square

The entanglement-assisted Holevo capacity of a channel

Along similar lines to the entanglement-assisted classical capacity, which mirrors the definition of the classical capacity in a setting where the sender and receiver initially share a state of their choosing, one may define the entanglement-assisted Holevo capacity of a channel. The following definition is helpful when formalizing this notion.

Definition 8.11 Let Σ be an alphabet, let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces, let $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X} \otimes \mathcal{Y})$ be an ensemble, and let

$$\rho = \sum_{a \in \Sigma} \eta(a) \quad (8.31)$$

denote the average state of η . It is said that η is *homogeneous on \mathcal{Y}* if it holds that

$$\text{Tr}_{\mathcal{X}}(\eta(a)) = \text{Tr}(\eta(a)) \text{Tr}_{\mathcal{X}}(\rho) \quad (8.32)$$

for every $a \in \Sigma$.

A simple operational characterization of ensembles homogeneous on a given complex Euclidean space is provided by the following proposition. In essence, it states that this sort of ensemble is one obtained by applying a randomly selected channel to the opposite subsystem of a fixed bipartite state.

Proposition 8.12 Let Σ be an alphabet, let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces, and let $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X} \otimes \mathcal{Y})$ be an ensemble. The following three statements are equivalent:

1. The ensemble η is homogeneous on \mathcal{Y} .
2. There exists a complex Euclidean space \mathcal{Z} , a state $\sigma \in \text{D}(\mathcal{Z} \otimes \mathcal{Y})$, a collection of channels $\{\Phi_a : a \in \Sigma\} \subseteq \text{C}(\mathcal{Z}, \mathcal{X})$, and a probability vector $p \in \mathcal{P}(\Sigma)$, such that

$$\eta(a) = p(a) (\Phi_a \otimes \mathbb{1}_{\text{L}(\mathcal{Y})})(\sigma) \quad (8.33)$$

for every $a \in \Sigma$.

3. Statement 2 holds under the additional assumption that $\sigma = uu^*$ for some choice of a unit vector $u \in \mathcal{Z} \otimes \mathcal{Y}$.

Proof The fact that the second statement implies the first is immediate, and the third statement trivially implies the second. It therefore remains to prove that the first statement implies the third.

To this end, assume that η is homogeneous on \mathcal{Y} , let ρ denote the average state of the ensemble η , and let

$$\xi = \text{Tr}_{\mathcal{X}}(\rho). \quad (8.34)$$

Let \mathcal{Z} be a complex Euclidean space of dimension $\text{rank}(\xi)$, and let $u \in \mathcal{Z} \otimes \mathcal{Y}$ be a unit vector that purifies ξ :

$$\text{Tr}_{\mathcal{Z}}(uu^*) = \xi. \quad (8.35)$$

As η is homogeneous on \mathcal{Y} , it therefore holds that

$$\mathrm{Tr}(\eta(a)) \mathrm{Tr}_{\mathcal{Z}}(uu^*) = \mathrm{Tr}_{\mathcal{X}}(\eta(a)) \quad (8.36)$$

for every $a \in \Sigma$. By Proposition 2.29, one concludes that there must exist a channel $\Phi_a \in \mathcal{C}(\mathcal{Z}, \mathcal{X})$ such that

$$\eta(a) = \mathrm{Tr}(\eta(a))(\Phi_a \otimes \mathbf{1}_{\mathcal{L}(\mathcal{Y})})(uu^*) \quad (8.37)$$

for every $a \in \Sigma$. Setting $\sigma = uu^*$ and $p(a) = \mathrm{Tr}(\eta(a))$ for each $a \in \Sigma$ completes the proof. \square

Definition 8.13 Let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} . The *entanglement-assisted Holevo capacity* of Φ is the quantity $\chi_{\mathrm{E}}(\Phi)$ defined as

$$\chi_{\mathrm{E}}(\Phi) = \sup_{\eta} \chi((\Phi \otimes \mathbf{1}_{\mathcal{L}(\mathcal{W})})(\eta)), \quad (8.38)$$

where the supremum is over all choices of a complex Euclidean space \mathcal{W} , an alphabet Σ , and an ensemble $\eta : \Sigma \rightarrow \mathrm{Pos}(\mathcal{X} \otimes \mathcal{W})$ homogeneous on \mathcal{W} .

The relationship between the entanglement-assisted classical capacity and the entanglement-assisted Holevo capacity is discussed in Section 8.1.3. In this context, for a given ensemble that is homogeneous on \mathcal{W} , the bipartite state whose existence is implied by Proposition 8.12 may be seen as being representative of a state shared between a sender and receiver that facilitates information transmission.

The coherent information

The final quantity, associated with a given channel, that is to be defined in the present subsection is the *coherent information*.

Definition 8.14 Let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ be a channel and let $\sigma \in \mathcal{D}(\mathcal{X})$ be a state, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} . The *coherent information* of σ through Φ is the quantity $I_{\mathrm{C}}(\sigma; \Phi)$ defined as

$$I_{\mathrm{C}}(\sigma; \Phi) = \mathrm{H}(\Phi(\sigma)) - \mathrm{H}\left((\Phi \otimes \mathbf{1}_{\mathcal{L}(\mathcal{X})})(\mathrm{vec}(\sqrt{\sigma}) \mathrm{vec}(\sqrt{\sigma})^*)\right). \quad (8.39)$$

The *maximum coherent information* of Φ is the quantity

$$I_{\mathrm{C}}(\Phi) = \max_{\sigma \in \mathcal{D}(\mathcal{X})} I_{\mathrm{C}}(\sigma; \Phi). \quad (8.40)$$

In general terms, the coherent information of a state σ through a channel Φ quantifies the correlations that exist after Φ is applied to a purification of σ . The definition implicitly takes this purification to be $\mathrm{vec}(\sqrt{\sigma})$ for the

sake of simplicity and concreteness; any other purification would result in the same quantity.

Consider the state

$$\rho = (\Phi \otimes \mathbf{1}_{\mathcal{L}(\mathcal{X})})(\mathrm{vec}(\sqrt{\sigma}) \mathrm{vec}(\sqrt{\sigma})^*) \in \mathcal{D}(\mathcal{Y} \otimes \mathcal{X}) \quad (8.41)$$

of a pair of registers $(\mathcal{Y}, \mathcal{X})$, corresponding to the spaces \mathcal{Y} and \mathcal{X} , as suggested by the definition above. One has that the coherent information $I_{\mathrm{C}}(\sigma; \Phi)$ of σ through Φ is equal to $\mathrm{H}(\mathcal{Y}) - \mathrm{H}(\mathcal{Y}, \mathcal{X})$. The quantum mutual information between \mathcal{Y} and \mathcal{X} is therefore given by

$$I(\mathcal{Y} : \mathcal{X}) = I_{\mathrm{C}}(\sigma; \Phi) + \mathrm{H}(\sigma). \quad (8.42)$$

While it is not immediately clear that the coherent information is relevant to the notion of channel capacity, it will be proved later in the chapter that this quantity is fundamentally important with respect to the entanglement-assisted classical capacity and the quantum capacity (to be defined later in Section 8.2).

The following proposition establishes an intuitive fact: with respect to an arbitrary choice of an input state, feeding the output of one channel into a second channel cannot lead to an increase in coherent information.

Proposition 8.15 Let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ and $\Psi \in \mathcal{C}(\mathcal{Y}, \mathcal{Z})$ be channels and let $\sigma \in \mathcal{D}(\mathcal{X})$ be a state, for complex Euclidean spaces \mathcal{X} , \mathcal{Y} , and \mathcal{Z} . It holds that

$$I_{\mathrm{C}}(\sigma; \Psi\Phi) \leq I_{\mathrm{C}}(\sigma; \Phi). \quad (8.43)$$

Proof Choose complex Euclidean spaces \mathcal{W} and \mathcal{V} , along with isometries $A \in \mathcal{U}(\mathcal{X}, \mathcal{Y} \otimes \mathcal{W})$ and $B \in \mathcal{U}(\mathcal{Y}, \mathcal{Z} \otimes \mathcal{V})$, so that Stinespring representations of Φ and Ψ are obtained:

$$\Phi(X) = \mathrm{Tr}_{\mathcal{W}}(AXA^*) \quad \text{and} \quad \Psi(Y) = \mathrm{Tr}_{\mathcal{V}}(BYB^*) \quad (8.44)$$

for all $X \in \mathcal{L}(\mathcal{X})$ and $Y \in \mathcal{L}(\mathcal{Y})$. Define a unit vector $u \in \mathcal{Z} \otimes \mathcal{V} \otimes \mathcal{W} \otimes \mathcal{X}$ as

$$u = (B \otimes \mathbf{1}_{\mathcal{W}} \otimes \mathbf{1}_{\mathcal{X}})(A \otimes \mathbf{1}_{\mathcal{X}}) \mathrm{vec}(\sqrt{\sigma}). \quad (8.45)$$

Now, consider four registers \mathcal{Z} , \mathcal{V} , \mathcal{W} , and \mathcal{X} , corresponding to the spaces \mathcal{Z} , \mathcal{V} , \mathcal{W} , and \mathcal{X} , respectively. Assuming the compound register $(\mathcal{Z}, \mathcal{V}, \mathcal{W}, \mathcal{X})$ is in the pure state uu^* , one has the following expressions:

$$I_{\mathrm{C}}(\sigma; \Phi) = \mathrm{H}(\mathcal{Z}, \mathcal{V}) - \mathrm{H}(\mathcal{Z}, \mathcal{V}, \mathcal{X}), \quad (8.46)$$

$$I_{\mathrm{C}}(\sigma; \Psi\Phi) = \mathrm{H}(\mathcal{Z}) - \mathrm{H}(\mathcal{Z}, \mathcal{X}).$$

The proposition follows from the strong subadditivity of the von Neumann entropy (Theorem 5.36). \square

It is convenient to refer to the notion of *complementary channels* in some of the proofs to be found in the present chapter. This notion is defined as follows.

Definition 8.16 Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ and $\Psi \in C(\mathcal{X}, \mathcal{Z})$ be channels, for \mathcal{X} , \mathcal{Y} , and \mathcal{Z} being complex Euclidean spaces. It is said that Φ and Ψ are *complementary* if there exists an isometry $A \in U(\mathcal{X}, \mathcal{Y} \otimes \mathcal{Z})$ for which it holds that

$$\Phi(X) = \text{Tr}_{\mathcal{Z}}(AXA^*) \quad \text{and} \quad \Psi(X) = \text{Tr}_{\mathcal{Y}}(AXA^*) \quad (8.47)$$

for every $X \in L(\mathcal{X})$.

It is immediate from Corollary 2.27 that, for every channel $\Phi \in C(\mathcal{X}, \mathcal{Y})$, there must exist a complex Euclidean space \mathcal{Z} and a channel $\Psi \in C(\mathcal{X}, \mathcal{Z})$ that is complementary to Φ ; such a channel Ψ is obtained from any choice of a Stinespring representation of Φ .

Proposition 8.17 Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ and $\Psi \in C(\mathcal{X}, \mathcal{Z})$ be complementary channels and let $\sigma \in D(\mathcal{X})$ be a state, for complex Euclidean spaces \mathcal{X} , \mathcal{Y} , and \mathcal{Z} . It holds that

$$I_c(\sigma; \Phi) = H(\Phi(\sigma)) - H(\Psi(\sigma)). \quad (8.48)$$

Proof By the assumption that Φ and Ψ are complementary, there must exist an isometry $A \in U(\mathcal{X}, \mathcal{Y} \otimes \mathcal{Z})$ such that the equations (8.47) hold for every $X \in L(\mathcal{X})$. Let \mathbf{X} , \mathbf{Y} , and \mathbf{Z} be registers corresponding to the spaces \mathcal{X} , \mathcal{Y} , and \mathcal{Z} , define a unit vector $u \in \mathcal{Y} \otimes \mathcal{Z} \otimes \mathcal{X}$ as

$$u = (A \otimes \mathbb{1}_{\mathcal{X}}) \text{vec}(\sqrt{\sigma}). \quad (8.49)$$

With respect to the pure state uu^* of the compound register $(\mathbf{Y}, \mathbf{Z}, \mathbf{X})$, it holds that $H(\mathbf{Z}) = H(\mathbf{Y}, \mathbf{X})$, and therefore

$$H\left(\left(\Phi \otimes \mathbb{1}_{L(\mathcal{X})}\right)\left(\text{vec}(\sqrt{\sigma}) \text{vec}(\sqrt{\sigma})^*\right)\right) = H(\Psi(\sigma)), \quad (8.50)$$

from which the proposition follows. \square

8.1.2 The Holevo–Schumacher–Westmoreland theorem

The *Holevo–Schumacher–Westmoreland theorem*, which is stated and proved in the present section, establishes that the classical capacity of a quantum channel is lower-bounded by its Holevo capacity, and that by regularizing the Holevo capacity one obtains a characterization of the classical capacity. The notion of a *classical-to-quantum product state channel code*, along with

a few mathematical results that are useful for analyzing these codes, will be introduced prior to the statement and proof of the Holevo–Schumacher–Westmoreland theorem.

Classical-to-quantum product state channel codes

When studying the classical capacity of quantum channels, it is instructive to consider a related but somewhat more basic task of encoding classical information using fixed sets of quantum states. When this task is connected with the notion of the classical capacity of a channel, a link must be made between the particular set of states used to encode classical information and the given channel—but it is reasonable to begin by examining the task of encoding classical information into quantum states in isolation.

Throughout the discussion that follows, $\Gamma = \{0, 1\}$ will denote the binary alphabet and

$$\{\sigma_a : a \in \Sigma\} \subseteq D(\mathcal{X}) \quad (8.51)$$

will denote a fixed collection of states, for \mathcal{X} being a complex Euclidean space and Σ being an alphabet.¹ The situation to be considered is that binary strings, representing classical information, are to be encoded into tensor products of quantum states drawn from the collection (8.51) in such a way that each binary string can be recovered from its encoding with high probability.

In more precise terms, it is to be assumed that positive integers n and m have been selected, and that every binary string $b_1 \cdots b_m \in \Gamma^m$ of length m is to be *encoded* by a product state having the form

$$\sigma_{a_1} \otimes \cdots \otimes \sigma_{a_n} \in D(\mathcal{X}^{\otimes n}), \quad (8.52)$$

for some choice of a string $a_1 \cdots a_n \in \Sigma^n$. That is, a function $f : \Gamma^m \rightarrow \Sigma^n$ is to be selected, and each string $b_1 \cdots b_m \in \Gamma^m$ is to be encoded by the state (8.52) for $a_1 \cdots a_n = f(b_1 \cdots b_m)$. When discussing this sort of code, it is convenient to make use of the shorthand notation

$$\sigma_{a_1 \cdots a_n} = \sigma_{a_1} \otimes \cdots \otimes \sigma_{a_n} \quad (8.53)$$

for each string $a_1 \cdots a_n \in \Sigma^n$, and with respect to this notation one has that

$$\sigma_{f(b_1 \cdots b_m)} \in D(\mathcal{X}^{\otimes n}) \quad (8.54)$$

denotes the state that encodes the string $b_1 \cdots b_m \in \Gamma^m$.

¹ The entire discussion could be generalized to allow for arbitrary alphabets Γ in place of the binary alphabet. As there is little gain in doing this from the perspective of this book, the assumption that $\Gamma = \{0, 1\}$ is made in the interest of simplicity.

From the encoding of a given binary string, one may hope to *decode* this string by means of a measurement. Such a measurement takes the form $\mu : \Gamma^m \rightarrow \text{Pos}(\mathcal{X}^{\otimes n})$, and succeeds in successfully recovering a particular string $b_1 \cdots b_m$ from its encoding with probability

$$\langle \mu(b_1 \cdots b_m), \sigma_{f(b_1 \cdots b_m)} \rangle. \quad (8.55)$$

As a general guideline, one is typically interested in coding schemes for which the probability of a successful decoding is close to 1 and the ratio m/n , which represents the rate at which classical information is effectively transmitted, is as large as possible. The following definition summarizes these notions.

Definition 8.18 Let Σ be an alphabet, let \mathcal{X} be a complex Euclidean space, let

$$\{\sigma_a : a \in \Sigma\} \subseteq \mathcal{D}(\mathcal{X}) \quad (8.56)$$

be a collection of states, let $\Gamma = \{0, 1\}$ denote the binary alphabet, and let n and m be positive integers. A *classical-to-quantum product state channel code* for the collection of states (8.56) is a pair (f, μ) consisting of a function and a measurement of the forms

$$f : \Gamma^m \rightarrow \Sigma^n \quad \text{and} \quad \mu : \Gamma^m \rightarrow \text{Pos}(\mathcal{X}^{\otimes n}). \quad (8.57)$$

The *rate* of such a code is equal to the ratio m/n , and the code is said to have *error bounded by δ* if it holds that

$$\langle \mu(b_1 \cdots b_m), \sigma_{f(b_1 \cdots b_m)} \rangle > 1 - \delta \quad (8.58)$$

for every string $b_1 \cdots b_m \in \Gamma^m$.

Remark The term *channel code* is used in this definition to distinguish this type of code from a *source code*, as discussed in Chapter 5. The two notions are, in some sense, complementary. A channel code represents the situation in which information is encoded into a state that possesses some degree of randomness, while a source code represents the situation in which information produced by a random source is encoded into a chosen state.

It is evident that some choices of sets $\{\sigma_a : a \in \Sigma\}$ are better suited to the construction of classical-to-quantum product state channel codes than others, assuming one wishes to maximize the rate and minimize the error of such a code. For the most part, the analysis that follows will be focused on the situation in which a set of states has been fixed, and one is interested in understanding the capabilities of this particular set, with respect to classical-to-quantum product state channel codes.

Typicality for ensembles of states

The notion of *typicality* is central to the proofs of multiple theorems to be presented in the current chapter, including a fundamental theorem on the existence of classical-to-quantum product state channel codes possessing certain rates and error bounds.

A standard definition of typicality was introduced in Section 5.3.1—but it is an extension of this definition to ensembles of states that will be used in the context of channel coding. The following definition is a starting point for a discussion of this concept, providing a notion of typicality for joint probability distributions.

Definition 8.19 Let $p \in \mathcal{P}(\Sigma \times \Gamma)$ be a probability vector, for alphabets Σ and Γ , and let $q \in \mathcal{P}(\Sigma)$ be the marginal probability vector defined as

$$q(a) = \sum_{b \in \Gamma} p(a, b) \quad (8.59)$$

for each $a \in \Sigma$. For every choice of a positive real number $\varepsilon > 0$, a positive integer n , and a string $a_1 \cdots a_n \in \Sigma^n$ satisfying $q(a_1) \cdots q(a_n) > 0$, a string $b_1 \cdots b_n \in \Gamma^n$ is said to be *ε -typical conditioned on $a_1 \cdots a_n \in \Sigma^n$* if

$$2^{-n(\mathbb{H}(p) - \mathbb{H}(q) + \varepsilon)} < \frac{p(a_1, b_1) \cdots p(a_n, b_n)}{q(a_1) \cdots q(a_n)} < 2^{-n(\mathbb{H}(p) - \mathbb{H}(q) - \varepsilon)}. \quad (8.60)$$

One writes $K_{a_1 \cdots a_n, \varepsilon}(p)$ to denote the set of all such strings $b_1 \cdots b_n \in \Gamma^n$.

It is also convenient to define $K_{a_1 \cdots a_n, \varepsilon}(p) = \emptyset$ for any string $a_1 \cdots a_n \in \Sigma^n$ for which $q(a_1) \cdots q(a_n) = 0$. When a probability vector $p \in \mathcal{P}(\Sigma \times \Gamma)$ is fixed, or can safely be taken as being implicit, the notation $K_{a_1 \cdots a_n, \varepsilon}$ may be used in place of $K_{a_1 \cdots a_n, \varepsilon}(p)$.

Intuitively speaking, if one were to select strings $a_1 \cdots a_n \in \Sigma^n$ and $b_1 \cdots b_n \in \Gamma^n$ by independently choosing $(a_1, b_1), \dots, (a_n, b_n)$ at random, according to a given probability vector $p \in \mathcal{P}(\Sigma \times \Gamma)$, then it would be reasonable to expect $b_1 \cdots b_n$ to be contained in $K_{a_1 \cdots a_n, \varepsilon}(p)$, with this event becoming increasingly likely as n becomes large. This fact is established by the following proposition, which is based on the weak law of large numbers (Theorem 1.15)—the methodology is essentially the same as the analogous fact (Proposition 5.42) that was proved in regard to the standard definition of typicality discussed in Section 5.3.1.

Proposition 8.20 Let $p \in \mathcal{P}(\Sigma \times \Gamma)$ be a probability vector, for alphabets Σ and Γ . For every $\varepsilon > 0$ it holds that

$$\lim_{n \rightarrow \infty} \sum_{a_1 \cdots a_n \in \Sigma^n} \sum_{b_1 \cdots b_n \in K_{a_1 \cdots a_n, \varepsilon}} p(a_1, b_1) \cdots p(a_n, b_n) = 1. \quad (8.61)$$

Proof Let $q \in \mathcal{P}(\Sigma)$ be the marginal probability vector defined as

$$q(a) = \sum_{b \in \Gamma} p(a, b) \tag{8.62}$$

for each $a \in \Sigma$, and define a random variable $X : \Sigma \times \Gamma \rightarrow [0, \infty)$ as

$$X(a, b) = \begin{cases} -\log(p(a, b)) + \log(q(a)) & \text{if } p(a, b) > 0 \\ 0 & \text{if } p(a, b) = 0 \end{cases} \tag{8.63}$$

and distributed according to the probability vector p . The expected value of this random variable is given by

$$E(X) = H(p) - H(q). \tag{8.64}$$

Now, for any positive integer n , and for X_1, \dots, X_n being independent random variables, each identically distributed to X , one has

$$\begin{aligned} \Pr\left(\left|\frac{X_1 + \dots + X_n}{n} - (H(p) - H(q))\right| < \varepsilon\right) \\ = \sum_{a_1 \dots a_n \in \Sigma^n} \sum_{b_1 \dots b_n \in K_{a_1 \dots a_n, \varepsilon}} p(a_1, b_1) \dots p(a_n, b_n). \end{aligned} \tag{8.65}$$

The conclusion of the proposition therefore follows from the weak law of large numbers (Theorem 1.15). \square

The next proposition places an upper bound on the expected size of the set $K_{a_1 \dots a_n, \varepsilon}$. It is analogous to Proposition 5.43 for the standard definition of typicality.

Proposition 8.21 *Let $p \in \mathcal{P}(\Sigma \times \Gamma)$ be a probability vector, for alphabets Σ and Γ , and let $q \in \mathcal{P}(\Sigma)$ be the marginal probability vector defined as*

$$q(a) = \sum_{b \in \Gamma} p(a, b) \tag{8.66}$$

for each $a \in \Sigma$. For every positive integer n and every positive real number $\varepsilon > 0$, it holds that

$$\sum_{a_1 \dots a_n \in \Sigma^n} q(a_1) \dots q(a_n) |K_{a_1 \dots a_n, \varepsilon}(p)| < 2^{n(H(p) - H(q) + \varepsilon)}. \tag{8.67}$$

Proof For each string $a_1 \dots a_n \in \Sigma^n$ satisfying $q(a_1) \dots q(a_n) > 0$ and each string $b_1 \dots b_n \in K_{a_1 \dots a_n, \varepsilon}(p)$, one has

$$2^{-n(H(p) - H(q) + \varepsilon)} < \frac{p(a_1, b_1) \dots p(a_n, b_n)}{q(a_1) \dots q(a_n)}, \tag{8.68}$$

and therefore

$$\begin{aligned} 2^{-n(H(p) - H(q) + \varepsilon)} \sum_{a_1 \dots a_n \in \Sigma^n} q(a_1) \dots q(a_n) |K_{a_1 \dots a_n, \varepsilon}(p)| \\ = \sum_{a_1 \dots a_n \in \Sigma^n} \sum_{b_1 \dots b_n \in K_{a_1 \dots a_n, \varepsilon}(p)} q(a_1) \dots q(a_n) 2^{-n(H(p) - H(q) + \varepsilon)} \\ < \sum_{a_1 \dots a_n \in \Sigma^n} \sum_{b_1 \dots b_n \in K_{a_1 \dots a_n, \varepsilon}} p(a_1, b_1) \dots p(a_n, b_n) \leq 1, \end{aligned} \tag{8.69}$$

from which the proposition follows. \square

The notion of typicality for joint probability distributions established by Definition 8.19 may be extended to ensembles of quantum states in a fairly straightforward fashion, by referring to spectral decompositions of the states in an ensemble.

Definition 8.22 Let $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$ be an ensemble of states, for \mathcal{X} a complex Euclidean space and Σ an alphabet, and let Γ be an alphabet such that $|\Gamma| = \dim(\mathcal{X})$. By the spectral theorem (as stated by Corollary 1.4), it follows that one may write

$$\eta(a) = \sum_{b \in \Gamma} p(a, b) u_{a,b} u_{a,b}^* \tag{8.70}$$

for some choice of a probability vector $p \in \mathcal{P}(\Sigma \times \Gamma)$ and an orthonormal basis $\{u_{a,b} : b \in \Gamma\}$ of \mathcal{X} for each $a \in \Sigma$. With respect to the ensemble η , and for each positive real number $\varepsilon > 0$, each positive integer n , and each string $a_1 \dots a_n \in \Sigma^n$, the *projection onto the ε -typical subspace of $\mathcal{X}^{\otimes n}$ conditioned on $a_1 \dots a_n$* is defined as

$$\Lambda_{a_1 \dots a_n, \varepsilon} = \sum_{b_1 \dots b_n \in K_{a_1 \dots a_n, \varepsilon}(p)} u_{a_1, b_1} u_{a_1, b_1}^* \otimes \dots \otimes u_{a_n, b_n} u_{a_n, b_n}^*. \tag{8.71}$$

Remark For a fixed choice of a string $a_1 \dots a_n \in \Sigma^n$, one has that the inclusion of each string $b_1 \dots b_n$ in $K_{a_1 \dots a_n, \varepsilon}(p)$ is determined by the multiset of values $\{p(a_1, b_1), \dots, p(a_n, b_n)\}$ alone. Thus, the same is true regarding the inclusion of each rank-one projection in the summation (8.71). It follows that the projection $\Lambda_{a_1 \dots a_n, \varepsilon}$ specified by Definition 8.22 is uniquely defined by the ensemble η , and is independent of the particular choices of the spectral decompositions (8.70).

Facts analogous to the previous two propositions, holding for ensembles rather than joint probability distributions, follow directly.

Proposition 8.23 Let $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$ be an ensemble of states, for \mathcal{X} a complex Euclidean space and Σ an alphabet. For every $\varepsilon > 0$, it holds that

$$\lim_{n \rightarrow \infty} \sum_{a_1 \cdots a_n \in \Sigma^n} \langle \Lambda_{a_1 \cdots a_n, \varepsilon}, \eta(a_1) \otimes \cdots \otimes \eta(a_n) \rangle = 1, \quad (8.72)$$

where, for each positive integer n , and each string $a_1 \cdots a_n \in \Sigma^n$, $\Lambda_{a_1 \cdots a_n, \varepsilon}$ is the projection onto the ε -typical subspace of $\mathcal{X}^{\otimes n}$ conditioned on $a_1 \cdots a_n$, with respect to the ensemble η . Moreover, one has

$$\sum_{a_1 \cdots a_n \in \Sigma^n} \text{Tr}(\eta(a_1)) \cdots \text{Tr}(\eta(a_n)) \text{Tr}(\Lambda_{a_1 \cdots a_n, \varepsilon}) < 2^{n(\beta + \varepsilon)} \quad (8.73)$$

for

$$\beta = \sum_{\substack{a \in \Sigma \\ \eta(a) \neq 0}} \text{Tr}(\eta(a)) \text{H}\left(\frac{\eta(a)}{\text{Tr}(\eta(a))}\right). \quad (8.74)$$

Proof For each $a \in \Sigma$, let

$$\eta(a) = \sum_{b \in \Gamma} p(a, b) u_{a,b} u_{a,b}^* \quad (8.75)$$

be a spectral decomposition of $\eta(a)$, as described in Definition 8.22, and define $q \in \mathcal{P}(\Sigma)$ as

$$q(a) = \sum_{b \in \Gamma} p(a, b) \quad (8.76)$$

(which is equivalent to $q(a) = \text{Tr}(\eta(a))$). For each positive integer n , each positive real number $\varepsilon > 0$, and each string $a_1 \cdots a_n \in \Sigma^n$, one has

$$\begin{aligned} & \langle \Lambda_{a_1 \cdots a_n, \varepsilon}, \eta(a_1) \otimes \cdots \otimes \eta(a_n) \rangle \\ &= \sum_{b_1 \cdots b_n \in K_{a_1 \cdots a_n, \varepsilon}} p(a_1, b_1) \cdots p(a_n, b_n), \end{aligned} \quad (8.77)$$

and moreover

$$\beta = \text{H}(p) - \text{H}(q) \quad \text{and} \quad \text{Tr}(\Lambda_{a_1 \cdots a_n, \varepsilon}) = |K_{a_1 \cdots a_n, \varepsilon}|. \quad (8.78)$$

The proposition therefore follows from Propositions 8.20 and 8.21. \square

A useful operator inequality

It is helpful to make use of an operator inequality, stated as Lemma 8.25 below, when analyzing the performance of classical-to-quantum product state channel codes. The proof of this inequality makes use of the following fact regarding square roots of positive semidefinite operators.

Lemma 8.24 (Operator monotonicity of the square root function) Let \mathcal{X} be a complex Euclidean space and let $P, Q \in \text{Pos}(\mathcal{X})$ be positive semidefinite operators. It holds that

$$\sqrt{P} \leq \sqrt{P+Q}. \quad (8.79)$$

Proof The block operator

$$\begin{pmatrix} P & \sqrt{P} \\ \sqrt{P} & \mathbf{1} \end{pmatrix} + \begin{pmatrix} Q & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} P+Q & \sqrt{P} \\ \sqrt{P} & \mathbf{1} \end{pmatrix} \quad (8.80)$$

is positive semidefinite. As $[P+Q, \mathbf{1}] = 0$ and \sqrt{P} is Hermitian, it follows by Lemma 5.29 that

$$\sqrt{P} \leq \sqrt{P+Q} \sqrt{\mathbf{1}} = \sqrt{P+Q}, \quad (8.81)$$

as required. \square

Remark It is not difficult to prove Lemma 8.24 directly, without relying on Lemma 5.29, by using spectral properties of operators that were also employed in the proof of that lemma.

Lemma 8.25 (Hayashi–Nagaoka) Let \mathcal{X} be a complex Euclidean space, let $P, Q \in \text{Pos}(\mathcal{X})$ be positive semidefinite operators, and assume $P \leq \mathbf{1}$. It holds that

$$\mathbf{1} - \sqrt{(P+Q)^+} P \sqrt{(P+Q)^+} \leq 2(\mathbf{1} - P) + 4Q. \quad (8.82)$$

Proof For every choice of operators $A, B \in \text{L}(\mathcal{X})$, one has

$$0 \leq (A - B)(A - B)^* = AA^* + BB^* - (AB^* + BA^*), \quad (8.83)$$

and therefore $AB^* + BA^* \leq AA^* + BB^*$. Setting

$$A = X\sqrt{Q} \quad \text{and} \quad B = (\mathbf{1} - X)\sqrt{Q}, \quad (8.84)$$

for a given operator $X \in \text{L}(\mathcal{X})$, yields

$$XQ(\mathbf{1} - X)^* + (\mathbf{1} - X)QX^* \leq XQX^* + (\mathbf{1} - X)Q(\mathbf{1} - X)^*, \quad (8.85)$$

and therefore

$$\begin{aligned} Q &= XQX^* + XQ(\mathbf{1} - X)^* + (\mathbf{1} - X)QX^* + (\mathbf{1} - X)Q(\mathbf{1} - X)^* \\ &\leq 2XQX^* + 2(\mathbf{1} - X)Q(\mathbf{1} - X)^*. \end{aligned} \quad (8.86)$$

For the specific choice $X = \sqrt{P+Q}$, one obtains

$$Q \leq 2\sqrt{P+Q}Q\sqrt{P+Q} + 2(\mathbf{1} - \sqrt{P+Q})Q(\mathbf{1} - \sqrt{P+Q}), \quad (8.87)$$

and from the observation that $Q \leq P + Q$ it follows that

$$\begin{aligned} Q &\leq 2\sqrt{P+Q}Q\sqrt{P+Q} \\ &\quad + 2(\mathbb{1} - \sqrt{P+Q})(P+Q)(\mathbb{1} - \sqrt{P+Q}) \\ &= \sqrt{P+Q}(2\mathbb{1} + 4Q - 4\sqrt{P+Q} + 2P)\sqrt{P+Q}. \end{aligned} \quad (8.88)$$

Using the fact that $P \leq \mathbb{1}$ together with Lemma 8.24, one has

$$P \leq \sqrt{P} \leq \sqrt{P+Q}, \quad (8.89)$$

and therefore

$$Q \leq \sqrt{P+Q}(2\mathbb{1} - 2P + 4Q)\sqrt{P+Q}. \quad (8.90)$$

Conjugating both sides of this inequality by the Moore–Penrose pseudo-inverse of $\sqrt{P+Q}$ yields

$$\sqrt{(P+Q)^+}Q\sqrt{(P+Q)^+} \leq 2\Pi_{\text{im}(P+Q)} - 2P + 4Q. \quad (8.91)$$

It follows that

$$\begin{aligned} \mathbb{1} - \sqrt{(P+Q)^+}P\sqrt{(P+Q)^+} \\ &= \mathbb{1} - \Pi_{\text{im}(P+Q)} + \sqrt{(P+Q)^+}Q\sqrt{(P+Q)^+} \\ &\leq \mathbb{1} + \Pi_{\text{im}(P+Q)} - 2P + 4Q \\ &\leq 2(\mathbb{1} - P) + 4Q, \end{aligned} \quad (8.92)$$

as required. \square

An existence proof for classical-to-quantum product state channel codes

Returning to the discussion of classical-to-quantum product state channel codes, assume as before that an alphabet Σ , a complex Euclidean space \mathcal{X} , and a collection of states

$$\{\sigma_a : a \in \Sigma\} \subseteq \text{D}(\mathcal{X}) \quad (8.93)$$

has been fixed, and let $\Gamma = \{0, 1\}$ denote the binary alphabet. It is natural to ask, for any choice of a positive real number $\delta > 0$ and positive integers m and n , whether or not there exists a classical-to-quantum product state channel code (f, μ) for this collection, taking the form

$$f : \Gamma^m \rightarrow \Sigma^n \quad \text{and} \quad \mu : \Gamma^m \rightarrow \text{Pos}(\mathcal{X}^{\otimes n}) \quad (8.94)$$

and having error bounded by δ .

In general, one may expect that making such a determination is not tractable from a computational point of view. It is possible, however, to prove the existence of reasonably good classical-to-quantum product state channel codes through the probabilistic method: for suitable choices of n , m , and δ , a *random* choice of a function $f : \Gamma^m \rightarrow \Sigma^n$ and a well-chosen measurement $\mu : \Gamma^m \rightarrow \text{Pos}(\mathcal{X}^{\otimes n})$ are considered, and a coding scheme with error bounded by δ is obtained with a nonzero probability. The theorem that follows gives a precise statement regarding the parameters n , m , and δ through which this methodology proves the existence of classical-to-quantum product state channels codes.

Theorem 8.26 *Let Σ be an alphabet, let \mathcal{X} be a complex Euclidean space, let*

$$\{\sigma_a : a \in \Sigma\} \subseteq \text{D}(\mathcal{X}) \quad (8.95)$$

be a collection of states, and let $\Gamma = \{0, 1\}$ denote the binary alphabet. Also let $p \in \mathcal{P}(\Sigma)$ be a probability vector, let $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$ be the ensemble defined as

$$\eta(a) = p(a)\sigma_a \quad (8.96)$$

for each $a \in \Sigma$, assume α is a positive real number satisfying $\alpha < \chi(\eta)$, and let $\delta > 0$ be a positive real number. For all but finitely many positive integers n , and for $m = \lfloor \alpha n \rfloor$, there exists a function $f : \Gamma^m \rightarrow \Sigma^n$ and a measurement $\mu : \Gamma^m \rightarrow \text{Pos}(\mathcal{X}^{\otimes n})$ such that

$$\langle \mu(b_1 \cdots b_m), \sigma_{f(b_1 \cdots b_m)} \rangle > 1 - \delta \quad (8.97)$$

for every $b_1 \cdots b_m \in \Gamma^m$.

Proof It will first be assumed that n and m are arbitrary positive integers. As suggested previously, the proof makes use of the probabilistic method: a random function $g : \Gamma^{m+1} \rightarrow \Sigma^n$ is chosen from a particular probability distribution, a decoding measurement μ is defined for each possible choice of g , and the expected probability of a decoding error for the pair (g, μ) is analyzed. As is to be explained later in the proof, this analysis implies the existence of a channel coding scheme (f, μ) , where $f : \Gamma^m \rightarrow \Sigma^n$ is derived from g , satisfying the requirements theorem for all but finitely many n and for $m = \lfloor \alpha n \rfloor$.

The particular distribution from which g is to be chosen is one in which each individual output symbol of g is selected independently according to the probability vector p . Equivalently, for a random selection of g according

to the distribution being described, one has that

$$\Pr(g(b_1 \cdots b_{m+1}) = a_1 \cdots a_n) = p(a_1) \cdots p(a_n) \quad (8.98)$$

for every choice of $b_1 \cdots b_{m+1} \in \Gamma^{m+1}$ and $a_1 \cdots a_n \in \Sigma^n$, and moreover the outputs of a randomly chosen g on distinct choices of the input string $b_1 \cdots b_{m+1}$ are uncorrelated.

The specification of the decoding measurement μ that is to be associated with a given g is not chosen randomly; a unique measurement is defined for each g in a way that is dependent upon the ensemble η . First, let $\varepsilon > 0$ be a sufficiently small positive real number such that the inequality

$$\alpha < \chi(\eta) - 3\varepsilon \quad (8.99)$$

holds. For each string $a_1 \cdots a_n \in \Sigma^n$, let $\Lambda_{a_1 \cdots a_n}$ denote the projection onto the ε -typical subspace of $\mathcal{X}^{\otimes n}$ conditioned on $a_1 \cdots a_n$, with respect to the ensemble η , and let Π_n be the projection onto the ε -typical subspace of $\mathcal{X}^{\otimes n}$ with respect to the average state

$$\sigma = \sum_{a \in \Sigma} p(a) \sigma_a \quad (8.100)$$

of the ensemble η . (As ε has been fixed, the dependence of $\Lambda_{a_1 \cdots a_n}$ and Π_n on ε is not written explicitly, allowing for slightly less cluttered equations.) Next, for a given choice of a function $g : \Gamma^{m+1} \rightarrow \Sigma^n$, define an operator

$$Q = \sum_{b_1 \cdots b_{m+1} \in \Gamma^{m+1}} \Pi_n \Lambda_{g(b_1 \cdots b_{m+1})} \Pi_n, \quad (8.101)$$

and, for each binary string $b_1 \cdots b_{m+1} \in \Gamma^{m+1}$, define an operator

$$Q_{b_1 \cdots b_{m+1}} = \sqrt{Q^+} \Pi_n \Lambda_{g(b_1 \cdots b_{m+1})} \Pi_n \sqrt{Q^+}. \quad (8.102)$$

Each operator $Q_{b_1 \cdots b_{m+1}}$ is positive semidefinite, and moreover

$$\sum_{b_1 \cdots b_{m+1} \in \Gamma^{m+1}} Q_{b_1 \cdots b_{m+1}} = \Pi_{\text{im}(Q)}. \quad (8.103)$$

Finally, the measurement $\mu : \Gamma^{m+1} \rightarrow \text{Pos}(\mathcal{X}^{\otimes n})$ to be associated with g is defined as

$$\mu(b_1 \cdots b_{m+1}) = Q_{b_1 \cdots b_{m+1}} + \frac{1}{2^{m+1}} (\mathbb{1} - \Pi_{\text{im}(Q)}) \quad (8.104)$$

for each $b_1 \cdots b_{m+1} \in \Gamma^{m+1}$.

For each choice of g , the probability that the measurement μ associated with g errs in recovering a string $b_1 \cdots b_{m+1} \in \Gamma^{m+1}$ from its encoding is equal to

$$\langle \mathbb{1} - \mu(b_1 \cdots b_{m+1}), \sigma_{g(b_1 \cdots b_{m+1})} \rangle. \quad (8.105)$$

The next phase of the proof establishes an upper bound on the average error probability

$$\frac{1}{2^{m+1}} \sum_{b_1 \cdots b_{m+1} \in \Gamma^{m+1}} \langle \mathbb{1} - \mu(b_1 \cdots b_{m+1}), \sigma_{g(b_1 \cdots b_{m+1})} \rangle, \quad (8.106)$$

for a uniformly chosen string $b_1 \cdots b_{m+1} \in \Gamma^{m+1}$. To bound this average probability of error, one may first observe that Lemma 8.25 implies that

$$\begin{aligned} \mathbb{1} - Q_{b_1 \cdots b_{m+1}} \\ \leq 2(\mathbb{1} - \Pi_n \Lambda_{g(b_1 \cdots b_{m+1})} \Pi_n) + 4(Q - \Pi_n \Lambda_{g(b_1 \cdots b_{m+1})} \Pi_n) \end{aligned} \quad (8.107)$$

for each $b_1 \cdots b_{m+1} \in \Gamma^{m+1}$. For a fixed choice of g , the probability of an error in recovering a given string $b_1 \cdots b_{m+1}$ is therefore upper-bounded by

$$\begin{aligned} 2\langle \mathbb{1} - \Pi_n \Lambda_{g(b_1 \cdots b_{m+1})} \Pi_n, \sigma_{g(b_1 \cdots b_{m+1})} \rangle \\ + 4\langle Q - \Pi_n \Lambda_{g(b_1 \cdots b_{m+1})} \Pi_n, \sigma_{g(b_1 \cdots b_{m+1})} \rangle. \end{aligned} \quad (8.108)$$

The expected value of this expression will be shown to be small, under the additional assumption that $m = \lfloor \alpha n \rfloor$, when $b_1 \cdots b_{m+1} \in \Gamma^{m+1}$ is chosen uniformly and g is chosen according to the distribution described above.

The first term in the expression (8.108) will be considered first. To prove an upper bound on the expected value of this quantity, it is convenient to make use of the operator identity

$$ABA = AB + BA - B + (\mathbb{1} - A)B(\mathbb{1} - A). \quad (8.109)$$

In particular, for any choice of a string $a_1 \cdots a_n \in \Sigma^n$, this identity implies

$$\begin{aligned} & \langle \Pi_n \Lambda_{a_1 \cdots a_n} \Pi_n, \sigma_{a_1 \cdots a_n} \rangle \\ &= \langle \Pi_n \Lambda_{a_1 \cdots a_n}, \sigma_{a_1 \cdots a_n} \rangle + \langle \Lambda_{a_1 \cdots a_n} \Pi_n, \sigma_{a_1 \cdots a_n} \rangle - \langle \Lambda_{a_1 \cdots a_n}, \sigma_{a_1 \cdots a_n} \rangle \\ & \quad + \langle (\mathbb{1} - \Pi_n) \Lambda_{a_1 \cdots a_n} (\mathbb{1} - \Pi_n), \sigma_{a_1 \cdots a_n} \rangle \\ & \geq \langle \Pi_n \Lambda_{a_1 \cdots a_n}, \sigma_{a_1 \cdots a_n} \rangle + \langle \Lambda_{a_1 \cdots a_n} \Pi_n, \sigma_{a_1 \cdots a_n} \rangle - \langle \Lambda_{a_1 \cdots a_n}, \sigma_{a_1 \cdots a_n} \rangle. \end{aligned} \quad (8.110)$$

As $\Lambda_{a_1 \dots a_n}$ is a projection operator and commutes with $\sigma_{a_1 \dots a_n}$, it follows that

$$\begin{aligned} & \langle \Pi_n \Lambda_{a_1 \dots a_n}, \sigma_{a_1 \dots a_n} \rangle + \langle \Lambda_{a_1 \dots a_n} \Pi_n, \sigma_{a_1 \dots a_n} \rangle - \langle \Lambda_{a_1 \dots a_n}, \sigma_{a_1 \dots a_n} \rangle \\ &= \langle 2\Pi_n - \mathbf{1}, \Lambda_{a_1 \dots a_n} \sigma_{a_1 \dots a_n} \rangle \\ &= \langle 2\Pi_n - \mathbf{1}, \sigma_{a_1 \dots a_n} \rangle + \langle \mathbf{1} - 2\Pi_n, (\mathbf{1} - \Lambda_{a_1 \dots a_n}) \sigma_{a_1 \dots a_n} \rangle \quad (8.111) \\ &\geq \langle 2\Pi_n - \mathbf{1}, \sigma_{a_1 \dots a_n} \rangle - \langle \mathbf{1} - \Lambda_{a_1 \dots a_n}, \sigma_{a_1 \dots a_n} \rangle \\ &= 2\langle \Pi_n, \sigma_{a_1 \dots a_n} \rangle + \langle \Lambda_{a_1 \dots a_n}, \sigma_{a_1 \dots a_n} \rangle - 2. \end{aligned}$$

By combining the inequalities (8.110) and (8.111), and averaging over all choices of $a_1 \dots a_n \in \Sigma^n$, with each a_k selected independently according to the probability vector p , one finds that

$$\begin{aligned} & \sum_{a_1 \dots a_n \in \Sigma^n} p(a_1) \dots p(a_n) \langle \Pi_n \Lambda_{a_1 \dots a_n} \Pi_n, \sigma_{a_1 \dots a_n} \rangle \\ &\geq 2\langle \Pi_n, \sigma^{\otimes n} \rangle + \sum_{a_1 \dots a_n \in \Sigma^n} p(a_1) \dots p(a_n) \langle \Lambda_{a_1 \dots a_n}, \sigma_{a_1 \dots a_n} \rangle - 2. \quad (8.112) \end{aligned}$$

The right-hand side of the expression (8.112) approaches 1 in the limit as n goes to infinity by Propositions 5.42 and 8.23, from which it follows that

$$\sum_{a_1 \dots a_n \in \Sigma^n} p(a_1) \dots p(a_n) \langle \mathbf{1} - \Pi_n \Lambda_{a_1 \dots a_n} \Pi_n, \sigma_{a_1 \dots a_n} \rangle < \frac{\delta}{8} \quad (8.113)$$

for all but finitely many choices of a positive integer n . For any n for which the inequality (8.113) holds, and for a random selection of $g : \Gamma^{m+1} \rightarrow \Sigma^n$ as described above, it therefore holds that the expected value of the expression

$$2\langle \mathbf{1} - \Pi_n \Lambda_{g(b_1 \dots b_{m+1})} \Pi_n, \sigma_{g(b_1 \dots b_{m+1})} \rangle \quad (8.114)$$

is at most $\delta/4$ for an arbitrary choice of $b_1 \dots b_{m+1}$, and therefore the same bound holds for a uniformly selected binary string $b_1 \dots b_{m+1} \in \Gamma^{m+1}$.

The second term in the expression (8.108) will be considered next. It may first be observed that

$$Q - \Pi_n \Lambda_{g(b_1 \dots b_{m+1})} \Pi_n = \sum_{\substack{c_1 \dots c_{m+1} \in \Gamma^{m+1} \\ c_1 \dots c_{m+1} \neq b_1 \dots b_{m+1}}} \Pi_n \Lambda_{g(c_1 \dots c_{m+1})} \Pi_n, \quad (8.115)$$

so that

$$\begin{aligned} & \langle Q - \Pi_n \Lambda_{g(b_1 \dots b_{m+1})} \Pi_n, \sigma_{g(b_1 \dots b_{m+1})} \rangle \\ &= \sum_{\substack{c_1 \dots c_{m+1} \in \Gamma^{m+1} \\ c_1 \dots c_{m+1} \neq b_1 \dots b_{m+1}}} \langle \Pi_n \Lambda_{g(c_1 \dots c_{m+1})} \Pi_n, \sigma_{g(b_1 \dots b_{m+1})} \rangle. \quad (8.116) \end{aligned}$$

The value of the function g on each input string is chosen independently according to the probability vector $p^{\otimes n}$, so there is no correlation between $g(b_1 \dots b_{m+1})$ and $g(c_1 \dots c_{m+1})$ for $b_1 \dots b_{m+1} \neq c_1 \dots c_{m+1}$. It follows that the expected value of the above expression is given by

$$(2^{m+1} - 1) \sum_{a_1 \dots a_n \in \Sigma^n} p(a_1) \dots p(a_n) \langle \Lambda_{a_1 \dots a_n}, \Pi_n \sigma^{\otimes n} \Pi_n \rangle. \quad (8.117)$$

By Proposition 8.23 it holds that

$$\sum_{a_1 \dots a_n \in \Sigma^n} p(a_1) \dots p(a_n) \text{Tr}(\Lambda_{a_1 \dots a_n}) \leq 2^{n(\beta+\varepsilon)} \quad (8.118)$$

for

$$\beta = \sum_{a \in \Sigma} p(a) \text{H}(\sigma_a), \quad (8.119)$$

and by the definition of Π_n one has that

$$\lambda_1(\Pi_n \sigma^{\otimes n} \Pi_n) \leq 2^{-n(\text{H}(\sigma) - \varepsilon)}. \quad (8.120)$$

It follows that

$$\begin{aligned} & (2^{m+1} - 1) \sum_{a_1 \dots a_n \in \Sigma^n} p(a_1) \dots p(a_n) \langle \Lambda_{a_1 \dots a_n}, \Pi_n \sigma^{\otimes n} \Pi_n \rangle \\ &\leq 2^{m+1-n(\chi(\eta)-2\varepsilon)}, \quad (8.121) \end{aligned}$$

so that the expected value of the second term in the expression (8.108) is upper-bounded by

$$2^{m-n(\chi(\eta)-2\varepsilon)+3}. \quad (8.122)$$

Now assume that $m = \lfloor \alpha n \rfloor$. For $g : \Gamma^{m+1} \rightarrow \Sigma^n$ chosen according to the distribution specified earlier and $b_1 \dots b_{m+1} \in \Gamma^{m+1}$ chosen uniformly, one has that the expected value of the error probability (8.106) is at most

$$\frac{\delta}{4} + 2^{\alpha n - n(\chi(\eta) - 2\varepsilon) + 3} \leq \frac{\delta}{4} + 2^{-\varepsilon n + 3} \quad (8.123)$$

for all but finitely many choices of n . As

$$2^{-\varepsilon n} < \frac{\delta}{32} \quad (8.124)$$

for all sufficiently large n , it follows that the expected value of the error probability (8.106) is smaller than $\delta/2$ for all but finitely many choices of n . For all but finitely many choices of n , there must therefore exist at least one

choice of a function $g : \Gamma^{m+1} \rightarrow \Sigma^n$ such that, for μ being the measurement associated with g , it holds that

$$\frac{1}{2^{m+1}} \sum_{b_1 \cdots b_{m+1} \in \Gamma^{m+1}} \langle \mathbb{1} - \mu(b_1 \cdots b_{m+1}), \sigma_{g(b_1 \cdots b_{m+1})} \rangle < \frac{\delta}{2}. \quad (8.125)$$

Finally, for a given choice of n , $m = \lfloor \alpha n \rfloor$, g , and μ for which the bound (8.125) holds, consider the set

$$B = \left\{ b_1 \cdots b_{m+1} \in \Gamma^{m+1} : \langle \mathbb{1} - \mu(b_1 \cdots b_{m+1}), \sigma_{g(b_1 \cdots b_{m+1})} \rangle \geq \delta \right\} \quad (8.126)$$

of all strings whose encodings incur a decoding error with probability at least δ . It holds that

$$\frac{\delta |B|}{2^{m+1}} < \frac{\delta}{2}, \quad (8.127)$$

and therefore $|B| \leq 2^m$. By defining a function $f : \Gamma^m \rightarrow \Sigma^n$ as $f = gh$, for an arbitrarily chosen injection $h : \Gamma^m \rightarrow \Gamma^{m+1} \setminus B$, one has that

$$\langle \mu(b_1 \cdots b_m), \sigma_{f(b_1 \cdots b_m)} \rangle > 1 - \delta \quad (8.128)$$

for every choice of $b_1 \cdots b_m \in \Gamma^m$, which completes the proof. \square

Statement and proof of the Holevo–Schumacher–Westmoreland theorem

The Holevo–Schumacher–Westmoreland theorem will now be stated, and proved through the use of Theorem 8.26.

Theorem 8.27 (Holevo–Schumacher–Westmoreland theorem) *Let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces and let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ be a channel. The classical capacity of Φ is equal to its regularized Holevo capacity:*

$$C(\Phi) = \lim_{n \rightarrow \infty} \frac{\chi(\Phi^{\otimes n})}{n}. \quad (8.129)$$

Proof The first main step of the proof is to establish the inequality

$$\chi(\Phi) \leq C(\Phi) \quad (8.130)$$

through the use of Theorem 8.26. This inequality holds trivially if $\chi(\Phi) = 0$, so it will be assumed that $\chi(\Phi)$ is positive.

Consider an ensemble $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$, for any alphabet Σ , expressed as $\eta(a) = p(a)\rho_a$ for each $a \in \Sigma$, where

$$\{\rho_a : a \in \Sigma\} \subseteq \mathcal{D}(\mathcal{X}) \quad (8.131)$$

is a collection of states and $p \in \mathcal{P}(\Sigma)$ is a probability vector. Assume that $\chi(\Phi(\eta))$ is positive and fix a positive real number $\alpha < \chi(\Phi(\eta))$. Also define

$\sigma_a = \Phi(\rho_a)$ for each $a \in \Sigma$, let $\varepsilon > 0$ be a positive real number, let $\Gamma = \{0, 1\}$ denote the binary alphabet, and define $\mathcal{Z} = \mathbb{C}^\Gamma$.

By Theorem 8.26, for all but finitely many choices of a positive integer n , and for $m = \lfloor \alpha n \rfloor$, there exists a classical-to-quantum product state channel code (f, μ) of the form

$$f : \Gamma^m \rightarrow \Sigma^n \quad \text{and} \quad \mu : \Gamma^m \rightarrow \text{Pos}(\mathcal{Y}^{\otimes n}) \quad (8.132)$$

for the collection

$$\{\sigma_a : a \in \Sigma\} \subseteq \mathcal{D}(\mathcal{Y}) \quad (8.133)$$

that errs with probability strictly less than $\varepsilon/2$ on every binary string of length m . Assume that such a choice of n , m , and a code (f, μ) have been fixed, and define encoding and decoding channels

$$\Xi_E \in \mathcal{C}(\mathcal{Z}^{\otimes m}, \mathcal{X}^{\otimes n}) \quad \text{and} \quad \Xi_D \in \mathcal{C}(\mathcal{Y}^{\otimes n}, \mathcal{Z}^{\otimes m}) \quad (8.134)$$

as follows:

$$\begin{aligned} \Xi_E(Z) &= \sum_{b_1 \cdots b_m \in \Gamma^m} \langle E_{b_1 \cdots b_m, b_1 \cdots b_m}, Z \rangle \rho_{f(b_1 \cdots b_m)}, \\ \Xi_D(Y) &= \sum_{b_1 \cdots b_m \in \Gamma^m} \langle \mu(b_1 \cdots b_m), Y \rangle E_{b_1 \cdots b_m, b_1 \cdots b_m}, \end{aligned} \quad (8.135)$$

for all $Z \in \mathcal{L}(\mathcal{Z}^{\otimes m})$ and $Y \in \mathcal{L}(\mathcal{Y}^{\otimes n})$. It follows from the properties of the code (f, μ) suggested above that

$$\langle E_{b_1 \cdots b_m, b_1 \cdots b_m}, (\Xi_D \Phi^{\otimes n} \Xi_E)(E_{b_1 \cdots b_m, b_1 \cdots b_m}) \rangle > 1 - \frac{\varepsilon}{2} \quad (8.136)$$

for every $b_1 \cdots b_m \in \Gamma^m$. As Ξ_E is a classical-to-quantum channel and Ξ_D is quantum-to-classical, one finds that $\Xi_D \Phi^{\otimes n} \Xi_E$ is a ε -approximation to the completely dephasing channel $\Delta^{\otimes m} \in \mathcal{C}(\mathcal{Z}^{\otimes m})$.

It has been proved that, for any choice of positive real numbers $\alpha < \chi(\Phi)$ and $\varepsilon > 0$, the channel $\Phi^{\otimes n}$ emulates an ε -approximation to the completely dephasing channel $\Delta^{\otimes m}$ for all but finitely many positive integers n and for $m = \lfloor \alpha n \rfloor$. From this fact the inequality (8.130) follows. One may apply the same reasoning to the channel $\Phi^{\otimes n}$ in place of Φ , for any positive integer n , to obtain

$$\frac{\chi(\Phi^{\otimes n})}{n} \leq \frac{C(\Phi^{\otimes n})}{n} = C(\Phi). \quad (8.137)$$

The second main step of the proof establishes that the regularized Holevo capacity is an upper bound on the classical capacity of Φ . When combined with the inequality (8.137), one finds that the limit in (8.129) indeed exists

and that the equality holds. There is nothing to prove if $C(\Phi) = 0$, so it will be assumed hereafter that $C(\Phi) > 0$.

Let $\alpha > 0$ be an achievable rate for classical information transmission through Φ , and let $\varepsilon > 0$ be chosen arbitrarily. It must therefore hold, for all but finitely many positive integers n , and for $m = \lfloor \alpha n \rfloor$, that $\Phi^{\otimes n}$ emulates an ε -approximation to the completely dephasing channel $\Delta^{\otimes m} \in \mathcal{C}(\mathcal{Z}^{\otimes m})$. Let n be any positive integer for which this property holds and for which $m = \lfloor \alpha n \rfloor \geq 2$. The situation in which a sender generates a binary string of length m , uniformly at random, and transmits this string through the ε -approximation to $\Delta^{\otimes m}$ emulated by $\Phi^{\otimes n}$ will be considered.

Let \mathbf{X} and \mathbf{Z} be classical registers both having state set Γ^m ; the register \mathbf{X} corresponds to the randomly generated string selected by the sender and \mathbf{Z} corresponds to the string obtained by the receiver when a copy of the string stored in \mathbf{X} is transmitted through the ε -approximation to $\Delta^{\otimes m}$ emulated by $\Phi^{\otimes n}$. As $\Phi^{\otimes n}$ emulates an ε -approximation to $\Delta^{\otimes m}$, there must exist a collection of states

$$\{\rho_{b_1 \dots b_m} : b_1 \dots b_m \in \Gamma^m\} \subseteq \mathcal{D}(\mathcal{X}^{\otimes n}), \quad (8.138)$$

along with a measurement $\mu : \Gamma^m \rightarrow \text{Pos}(\mathcal{Y}^{\otimes n})$, such that

$$\langle \mu(b_1 \dots b_m), \Phi^{\otimes n}(\rho_{b_1 \dots b_m}) \rangle > 1 - \frac{\varepsilon}{2} \quad (8.139)$$

for every binary string $b_1 \dots b_m \in \Gamma^m$. With respect to the probability vector $p \in \mathcal{P}(\Gamma^m \times \Gamma^m)$ defined as

$$p(b_1 \dots b_m, c_1 \dots c_m) = \frac{1}{2^m} \langle \mu(c_1 \dots c_m), \Phi^{\otimes n}(\rho_{b_1 \dots b_m}) \rangle, \quad (8.140)$$

which represents the probabilistic state of (\mathbf{X}, \mathbf{Z}) suggested above, it follows from Holevo's theorem (Theorem 5.49) that

$$I(\mathbf{X} : \mathbf{Z}) \leq \chi(\Phi^{\otimes n}(\eta)), \quad (8.141)$$

where $\eta : \Gamma^m \rightarrow \text{Pos}(\mathcal{X}^{\otimes n})$ is the ensemble defined as

$$\eta(b_1 \dots b_m) = \frac{1}{2^m} \rho_{b_1 \dots b_m} \quad (8.142)$$

for each $b_1 \dots b_m \in \Gamma^m$.

A lower bound on the mutual information $I(\mathbf{X} : \mathbf{Z})$ will now be derived. The distribution represented by the marginal probability vector $p[\mathbf{X}]$ is uniform, and therefore $H(p[\mathbf{X}]) = m$. By (8.139), each entry of the probability vector $p[\mathbf{Z}]$ is lower-bounded by $(1 - \varepsilon/2)2^{-m}$. It is therefore possible to write

$$p[\mathbf{Z}] = \left(1 - \frac{\varepsilon}{2}\right)r + \frac{\varepsilon}{2}q \quad (8.143)$$

for $q \in \mathcal{P}(\Gamma^m)$ being some choice of a probability vector and $r \in \mathcal{P}(\Gamma^m)$ denoting the uniform probability vector, defined as $r(b_1 \dots b_m) = 2^{-m}$ for every $b_1 \dots b_m \in \Gamma^m$. The inequality

$$H(p[\mathbf{Z}]) \geq \left(1 - \frac{\varepsilon}{2}\right)H(r) + \frac{\varepsilon}{2}H(q) \geq \left(1 - \frac{\varepsilon}{2}\right)m \quad (8.144)$$

follows by the concavity of the Shannon entropy function (Proposition 5.5). On the other hand, because the probability vector p satisfies

$$p(b_1 \dots b_m, b_1 \dots b_m) \geq \left(1 - \frac{\varepsilon}{2}\right)2^{-m} \quad (8.145)$$

for every $b_1 \dots b_m \in \Gamma^m$, it must hold that

$$\begin{aligned} H(p) &\leq -\left(1 - \frac{\varepsilon}{2}\right) \log\left(\frac{1 - \varepsilon/2}{2^m}\right) - \frac{\varepsilon}{2} \log\left(\frac{\varepsilon/2}{2^{2m} - 2^m}\right) \\ &< \left(1 + \frac{\varepsilon}{2}\right)m + H\left(1 - \frac{\varepsilon}{2}, \frac{\varepsilon}{2}\right) \leq \left(1 + \frac{\varepsilon}{2}\right)m + 1; \end{aligned} \quad (8.146)$$

the first inequality is a consequence of the fact that the entropy of p subject to the constraint (8.145) is maximized when p is defined as follows:

$$p(b_1 \dots b_m, c_1 \dots c_m) = \begin{cases} \frac{1 - \varepsilon/2}{2^m} & b_1 \dots b_m = c_1 \dots c_m \\ \frac{\varepsilon/2}{2^m(2^m - 1)} & b_1 \dots b_m \neq c_1 \dots c_m. \end{cases} \quad (8.147)$$

It therefore follows that

$$\begin{aligned} \chi(\Phi^{\otimes n}) &\geq I(\mathbf{X} : \mathbf{Z}) = H(p[\mathbf{X}]) + H(p[\mathbf{Z}]) - H(p) \\ &\geq (1 - \varepsilon)m - 1 \geq (1 - \varepsilon)\alpha n - 2, \end{aligned} \quad (8.148)$$

and consequently

$$\frac{\chi(\Phi^{\otimes n})}{n} \geq (1 - \varepsilon)\alpha - \frac{2}{n}. \quad (8.149)$$

It has been proved, for any achievable rate $\alpha > 0$ for classical information transmission through Φ , and for any $\varepsilon > 0$, that the inequality (8.149) holds for all but finitely many positive integers n . Because the supremum over all achievable rates α for classical information transmission through Φ is equal to $C(\Phi)$, this inequality may be combined with (8.137) to obtain the required equality (8.129). \square

8.1.3 The entanglement-assisted classical capacity theorem

This section focuses on the *entanglement-assisted classical capacity theorem*, which characterizes the entanglement-assisted classical capacity of a given channel. It stands out among the capacity theorems presented in the present chapter, as no regularization is required by the characterization it provides.

Holevo–Schumacher–Westmoreland theorem with entanglement assistance

A preliminary step toward the proof of the entanglement-assisted classical capacity theorem is the observation that, when the classical capacity and Holevo capacity are replaced by their entanglement-assisted formulations, a statement analogous to the Holevo–Schumacher–Westmoreland theorem holds.

Theorem 8.28 *Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} . The entanglement-assisted classical capacity of Φ equals the regularized entanglement-assisted Holevo capacity of Φ :*

$$C_E(\Phi) = \lim_{n \rightarrow \infty} \frac{\chi_E(\Phi^{\otimes n})}{n}. \quad (8.150)$$

Proof The theorem is proved in essentially the same way as the Holevo–Schumacher–Westmoreland theorem (Theorem 8.27), with each step being modified to allow for the possibility of entanglement assistance.

In greater detail, let Σ be an alphabet, let \mathcal{W} be a complex Euclidean space, let η be an ensemble of the form $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X} \otimes \mathcal{W})$ that is homogeneous on \mathcal{W} , assume $\chi((\Phi \otimes \mathbb{1}_{L(\mathcal{W})})(\eta))$ is positive, and let α be a positive real number satisfying

$$\alpha < \chi((\Phi \otimes \mathbb{1}_{L(\mathcal{W})})(\eta)). \quad (8.151)$$

By Proposition 8.12, one may choose a complex Euclidean space \mathcal{V} , a state $\xi \in D(\mathcal{V} \otimes \mathcal{W})$, a probability vector $p \in \mathcal{P}(\Sigma)$, and a collection of channels

$$\{\Psi_a : a \in \Sigma\} \subseteq C(\mathcal{V}, \mathcal{X}) \quad (8.152)$$

such that

$$\eta(a) = p(a)(\Psi_a \otimes \mathbb{1}_{L(\mathcal{W})})(\xi) \quad (8.153)$$

for every $a \in \Sigma$. For each $a \in \Sigma$ let

$$\sigma_a = (\Phi \Psi_a \otimes \mathbb{1}_{L(\mathcal{W})})(\xi), \quad (8.154)$$

and also let $\varepsilon > 0$ be an arbitrarily chosen positive real number.

By Theorem 8.26, for all but finitely many choices of a positive integer n , and for $m = \lfloor \alpha n \rfloor$, there exists a classical-to-quantum product state channel code (f, μ) of the form

$$f : \Gamma^m \rightarrow \Sigma^n \quad \text{and} \quad \mu : \Gamma^m \rightarrow \text{Pos}((\mathcal{Y} \otimes \mathcal{W})^{\otimes n}) \quad (8.155)$$

for the collection $\{\sigma_a : a \in \Sigma\} \subseteq D(\mathcal{Y} \otimes \mathcal{W})$ that errs with probability strictly less than $\varepsilon/2$ on every binary string of length m . Assume that such a choice of n , m , and a code (f, μ) have been fixed.

It will now be proved that the channel $\Phi^{\otimes n}$ emulates a ε -approximation to the completely dephasing channel $\Delta^{\otimes m} \in C(\mathcal{Z}^{\otimes m})$ with the assistance of entanglement. The entangled state to be used to assist this emulation is

$$V\xi^{\otimes n}V^* \in D(\mathcal{V}^{\otimes n} \otimes \mathcal{W}^{\otimes n}), \quad (8.156)$$

where $V \in U((\mathcal{V} \otimes \mathcal{W})^{\otimes n}, \mathcal{V}^{\otimes n} \otimes \mathcal{W}^{\otimes n})$ represents a permutation of tensor factors:

$$\begin{aligned} V((v_1 \otimes w_1) \otimes \cdots \otimes (v_n \otimes w_n)) \\ = (v_1 \otimes \cdots \otimes v_n) \otimes (w_1 \otimes \cdots \otimes w_n) \end{aligned} \quad (8.157)$$

for all vectors $v_1, \dots, v_n \in \mathcal{V}$ and $w_1, \dots, w_n \in \mathcal{W}$.

The encoding channel $\Xi_E \in C(\mathcal{Z}^{\otimes m} \otimes \mathcal{V}^{\otimes n}, \mathcal{X}^{\otimes n})$ used to perform this emulation is defined as

$$\Xi_E = \sum_{b_1 \cdots b_m \in \Gamma^m} \Theta_{b_1 \cdots b_m} \otimes \Psi_{f(b_1 \cdots b_m)}, \quad (8.158)$$

where

$$\Psi_{a_1 \cdots a_n} = \Psi_{a_1} \otimes \cdots \otimes \Psi_{a_n} \quad (8.159)$$

for each $a_1 \cdots a_n \in \Sigma^n$, and where $\Theta_{b_1 \cdots b_m} \in \text{CP}(\mathcal{Z}^{\otimes m}, \mathbb{C})$ is given by

$$\Theta_{b_1 \cdots b_m}(Z) = Z(b_1 \cdots b_m, b_1 \cdots b_m) \quad (8.160)$$

for every $Z \in L(\mathcal{Z}^{\otimes m})$. Described in words, the encoding map Ξ_E takes as input a compound register $(Z_1, \dots, Z_m, V_1, \dots, V_n)$, measures (Z_1, \dots, Z_m) with respect to the standard basis measurement, and applies the channel $\Psi_{f(b_1 \cdots b_m)}$ to (V_1, \dots, V_n) , for $b_1 \cdots b_m$ being the string obtained from the standard basis measurement on (Z_1, \dots, Z_m) .

The decoding channel $\Xi_D \in C(\mathcal{Y}^{\otimes n} \otimes \mathcal{W}^{\otimes n}, \mathcal{Z}^{\otimes m})$ used to perform the emulation is defined as

$$\Xi_D(Y) = \sum_{b_1 \cdots b_m \in \Gamma^m} \langle W \mu(b_1 \cdots b_m) W^*, Y \rangle E_{b_1 \cdots b_m, b_1 \cdots b_m} \quad (8.161)$$

for all $Y \in L(\mathcal{Y}^{\otimes n} \otimes \mathcal{W}^{\otimes n})$, where $W \in U((\mathcal{Y} \otimes \mathcal{W})^{\otimes n}, \mathcal{Y}^{\otimes n} \otimes \mathcal{W}^{\otimes n})$ is an isometry representing a permutation of tensor factors that is similar to V , but with \mathcal{V} replaced by \mathcal{Y} :

$$\begin{aligned} W((y_1 \otimes w_1) \otimes \cdots \otimes (y_n \otimes w_n)) \\ = (y_1 \otimes \cdots \otimes y_n) \otimes (w_1 \otimes \cdots \otimes w_n) \end{aligned} \quad (8.162)$$

for all choices of vectors $y_1, \dots, y_n \in \mathcal{Y}$ and $w_1, \dots, w_n \in \mathcal{W}$.

Now, let $\Psi \in \mathcal{C}(\mathcal{Z}^{\otimes m})$ denote the channel that has been emulated with the assistance of entanglement by the above construction; this channel may be expressed as

$$\Psi(Z) = (\Xi_D(\Phi^{\otimes n} \Xi_E \otimes \mathbb{1}_{L(\mathcal{W})}^{\otimes n})) (Z \otimes V \xi^{\otimes n} V^*) \quad (8.163)$$

for every $Z \in \mathcal{L}(\mathcal{Z}^{\otimes m})$, and it may be observed that $\Psi = \Delta^{\otimes m} \Psi \Delta^{\otimes m}$. For every string $b_1 \cdots b_m \in \Gamma^m$ it holds that

$$(\Phi^{\otimes n} \Xi_E \otimes \mathbb{1}_{L(\mathcal{W})}^{\otimes n})(E_{b_1 \cdots b_m, b_1 \cdots b_m} \otimes V \xi^{\otimes n} V^*) = W \sigma_{f(b_1 \cdots b_m)} W^*, \quad (8.164)$$

and therefore

$$\langle E_{b_1 \cdots b_m, b_1 \cdots b_m}, \Psi(E_{b_1 \cdots b_m, b_1 \cdots b_m}) \rangle > 1 - \frac{\varepsilon}{2}. \quad (8.165)$$

It follows that Ψ is a ε -approximation to $\Delta^{\otimes m}$, as claimed.

In summary, for any choice of positive real numbers $\alpha < \chi_E(\Phi)$ and $\varepsilon > 0$, it holds that $\Phi^{\otimes n}$ emulates an ε -approximation to the completely dephasing channel $\Delta^{\otimes m}$ with the assistance of entanglement, for all but finitely many positive integers n and for $m = \lfloor \alpha n \rfloor$. From this fact one concludes that $\chi_E(\Phi) \leq C_E(\Phi)$. Applying the same argument to the channel $\Phi^{\otimes n}$ in place of Φ , for any choice of a positive integer n , yields

$$\frac{\chi_E(\Phi^{\otimes n})}{n} \leq \frac{C_E(\Phi^{\otimes n})}{n} = C_E(\Phi). \quad (8.166)$$

Next it will be proved that the entanglement-assisted classical capacity of Φ cannot exceed its regularized entanglement-assisted Holevo capacity. As in the proof of Theorem 8.27, it may be assumed that $C_E(\Phi) > 0$, and it suffices to consider the situation in which a sender transmits a uniformly generated binary string of length m to a receiver.

Suppose $\alpha > 0$ is an achievable rate for entanglement-assisted classical information transmission through Φ , and let $\varepsilon > 0$ be chosen arbitrarily. It must therefore hold, for all but finitely many positive integers n , and for $m = \lfloor \alpha n \rfloor$, that $\Phi^{\otimes n}$ emulates an ε -approximation to the completely dephasing channel $\Delta^{\otimes m}$ with the assistance of entanglement. Let n be an arbitrarily chosen positive integer for which this property holds and for which $m = \lfloor \alpha n \rfloor \geq 2$.

As before, let \mathbf{X} and \mathbf{Z} be classical registers both having state set Γ^m ; \mathbf{X} stores the randomly generated string selected by the sender and \mathbf{Z} represents the string obtained by the receiver when a copy of the string stored in \mathbf{X} is transmitted through the ε -approximation to $\Delta^{\otimes m}$ emulated by $\Phi^{\otimes n}$ with the assistance of entanglement. By the assumption that $\Phi^{\otimes n}$ emulates an ε -approximation to $\Delta^{\otimes m}$ with the assistance of entanglement, one may

conclude that there exists a choice of complex Euclidean spaces \mathcal{V} and \mathcal{W} , a state $\xi \in \mathcal{D}(\mathcal{V} \otimes \mathcal{W})$, a collection of channels

$$\{\Psi_{b_1 \cdots b_m} : b_1 \cdots b_m \in \Gamma^m\} \subset \mathcal{C}(\mathcal{V}, \mathcal{X}^{\otimes n}), \quad (8.167)$$

and a measurement $\mu : \Gamma^m \rightarrow \text{Pos}(\mathcal{Y}^{\otimes n} \otimes \mathcal{W})$, such that

$$\langle \mu(b_1 \cdots b_m), (\Phi^{\otimes n} \Psi_{b_1 \cdots b_m} \otimes \mathbb{1}_{L(\mathcal{W})})(\xi) \rangle > 1 - \frac{\varepsilon}{2} \quad (8.168)$$

for every string $b_1 \cdots b_m \in \Gamma^m$. With respect to $p \in \mathcal{P}(\Gamma^m \times \Gamma^m)$ defined as

$$\begin{aligned} & p(b_1 \cdots b_m, c_1 \cdots c_m) \\ &= \frac{1}{2^m} \langle \mu(c_1 \cdots c_m), (\Phi^{\otimes n} \Psi_{b_1 \cdots b_m} \otimes \mathbb{1}_{L(\mathcal{W})})(\xi) \rangle, \end{aligned} \quad (8.169)$$

which represents the probabilistic state of (\mathbf{X}, \mathbf{Z}) suggested above, it follows from Holevo's theorem (Theorem 5.49) that

$$I(\mathbf{X} : \mathbf{Z}) \leq \chi((\Phi^{\otimes n} \otimes \mathbb{1}_{L(\mathcal{W})})(\eta)), \quad (8.170)$$

for $\eta : \Gamma^m \rightarrow \text{Pos}(\mathcal{X}^{\otimes n} \otimes \mathcal{W})$ being the ensemble defined as

$$\eta(b_1 \cdots b_m) = \frac{1}{2^m} (\Psi_{b_1 \cdots b_m} \otimes \mathbb{1}_{L(\mathcal{W})})(\xi) \quad (8.171)$$

for each $b_1 \cdots b_m \in \Gamma^m$.

The same lower-bound on the quantity $I(\mathbf{X} : \mathbf{Z})$ derived in the proof of Theorem 8.27 holds in the present case, from which it follows that

$$\chi_E(\Phi^{\otimes n}) \geq I(\mathbf{X} : \mathbf{Z}) \geq (1 - \varepsilon)\alpha n - 2, \quad (8.172)$$

and therefore

$$\frac{\chi_E(\Phi^{\otimes n})}{n} \geq (1 - \varepsilon)\alpha - \frac{2}{n}. \quad (8.173)$$

Thus, for any achievable rate $\alpha > 0$ for entanglement-assisted classical information transmission through Φ , and for any positive real number $\varepsilon > 0$, the inequality (8.173) holds for all but finitely many positive integers n . Because the supremum over all achievable rates α for entanglement-assisted classical information transmission through Φ is equal to $C_E(\Phi)$, one may combine this inequality with the upper bound (8.166) to obtain the required equality (8.150). \square

Strongly typical strings and projections

The proof of the entanglement-assisted classical capacity theorem that is presented in this book will make use of a notion of typicality, known as *strong typicality*, that differs from the standard notion discussed previously

in Section 5.3.1. True to its name, strong typicality is the more restrictive of the two notions; every strongly typical string will necessarily be a typical string, up to a simple change of parameters, while some typical strings are not strongly typical.

Similar to the standard notion of typicality, one may define an ε -strongly typical subspace with respect to a spectral decomposition of a given state. Unlike the standard typical subspace, however, the strongly typical subspace is not always uniquely determined by a given state; it can depend on the particular choice of a spectral decomposition (in the sense of Corollary 1.4) with respect to which it is defined. Despite this apparent drawback, the notion of an ε -strongly typical subspace will be a useful tool when proving the entanglement-assisted classical capacity theorem.

The definition of strong-typicality to follow uses the following notation, for which it is to be assumed that Σ is an alphabet and n is a positive integer. For every string $a_1 \cdots a_n \in \Sigma^n$ and symbol $a \in \Sigma$, one writes

$$N(a | a_1 \cdots a_n) = |\{k \in \{1, \dots, n\} : a_k = a\}|, \quad (8.174)$$

which is the number of times the symbol a occurs in the string $a_1 \cdots a_n$.

Definition 8.29 Let Σ be an alphabet, let $p \in \mathcal{P}(\Sigma)$ be a probability vector, let n be a positive integer, and let $\varepsilon > 0$ be a positive real number. A string $a_1 \cdots a_n \in \Sigma^n$ is said to be ε -strongly typical with respect to p if

$$\left| \frac{N(a | a_1 \cdots a_n)}{n} - p(a) \right| \leq p(a)\varepsilon \quad (8.175)$$

for every $a \in \Sigma$. The set of all ε -strongly typical strings of length n with respect to p is denoted $S_{n,\varepsilon}(p)$ (or by $S_{n,\varepsilon}$ when p is implicit and can safely be omitted).

The average behavior of a nonnegative real-valued function defined on the individual symbols of a strongly typical string may be analyzed using the following elementary proposition.

Proposition 8.30 Let Σ be an alphabet, let $p \in \mathcal{P}(\Sigma)$ be a probability vector, let n be a positive integer, let $\varepsilon > 0$ be a positive real number, let $a_1 \cdots a_n \in S_{n,\varepsilon}(p)$ be an ε -strongly typical string with respect to p , and let $\phi : \Sigma \rightarrow [0, \infty)$ be a nonnegative real-valued function. It holds that

$$\left| \frac{\phi(a_1) + \cdots + \phi(a_n)}{n} - \sum_{a \in \Sigma} p(a)\phi(a) \right| \leq \varepsilon \sum_{a \in \Sigma} p(a)\phi(a). \quad (8.176)$$

Proof The inequality (8.176) follows from the definition of strong typicality together with the triangle inequality:

$$\begin{aligned} & \left| \frac{\phi(a_1) + \cdots + \phi(a_n)}{n} - \sum_{a \in \Sigma} p(a)\phi(a) \right| \\ &= \left| \sum_{a \in \Sigma} \left(\frac{N(a | a_1 \cdots a_n)\phi(a)}{n} - p(a)\phi(a) \right) \right| \\ &\leq \sum_{a \in \Sigma} \phi(a) \left| \frac{N(a | a_1 \cdots a_n)}{n} - p(a) \right| \leq \varepsilon \sum_{a \in \Sigma} p(a)\phi(a), \end{aligned} \quad (8.177)$$

as required. \square

As a corollary to Proposition 8.30, one has that every ε -strongly typical string, with respect to a given probability vector p , is necessarily δ -typical for every choice of $\delta > \varepsilon H(p)$.

Corollary 8.31 Let Σ be an alphabet, let $p \in \mathcal{P}(\Sigma)$ be a probability vector, let n be a positive integer, let $\varepsilon > 0$ be a positive real number, and let $a_1 \cdots a_n \in S_{n,\varepsilon}(p)$ be an ε -strongly typical string with respect to p . It holds that

$$2^{-n(1+\varepsilon)H(p)} \leq p(a_1) \cdots p(a_n) \leq 2^{-n(1-\varepsilon)H(p)}. \quad (8.178)$$

Proof Define a function $\phi : \Sigma \rightarrow [0, \infty)$ as

$$\phi(a) = \begin{cases} -\log(p(a)) & \text{if } p(a) > 0 \\ 0 & \text{if } p(a) = 0. \end{cases} \quad (8.179)$$

With respect to this function, the implication provided by Proposition 8.30 is equivalent to (8.178). \square

Strings that are obtained by independently selecting symbols at random according to a given probability vector are likely to be not only typical, but strongly typical, with the probability of strong typicality increasing with string length. The following lemma establishes a quantitative bound on this probability.

Lemma 8.32 Let Σ be an alphabet, let $p \in \mathcal{P}(\Sigma)$ be a probability vector, let n be a positive integer, and let $\varepsilon > 0$ be a positive real number. It holds that

$$\sum_{a_1 \cdots a_n \in S_{n,\varepsilon}(p)} p(a_1) \cdots p(a_n) \geq 1 - \zeta_{n,\varepsilon}(p) \quad (8.180)$$

for

$$\zeta_{n,\varepsilon}(p) = 2 \sum_{\substack{a \in \Sigma \\ p(a) > 0}} \exp(-2n\varepsilon^2 p(a)^2). \quad (8.181)$$

Proof Suppose first that $a \in \Sigma$ is fixed, and consider the probability that a string $a_1 \cdots a_n \in \Sigma^n$, randomly selected according to the probability vector $p^{\otimes n}$, satisfies

$$\left| \frac{N(a | a_1 \cdots a_n)}{n} - p(a) \right| > p(a)\varepsilon. \quad (8.182)$$

To bound this probability, one may define X_1, \dots, X_n to be independent and identically distributed random variables, taking value 1 with probability $p(a)$ and value 0 otherwise, so that the probability of the event (8.182) is equal to

$$\Pr\left(\left|\frac{X_1 + \cdots + X_n}{n} - p(a)\right| > p(a)\varepsilon\right). \quad (8.183)$$

If it is the case that $p(a) > 0$, then Hoeffding's inequality (Theorem 1.16) implies that

$$\Pr\left(\left|\frac{X_1 + \cdots + X_n}{n} - p(a)\right| > p(a)\varepsilon\right) \leq 2 \exp(-2n\varepsilon^2 p(a)^2), \quad (8.184)$$

while it holds that

$$\Pr\left(\left|\frac{X_1 + \cdots + X_n}{n} - p(a)\right| > p(a)\varepsilon\right) = 0 \quad (8.185)$$

in case $p(a) = 0$. The lemma follows from the union bound. \square

The next proposition establishes upper and lower bounds on the number of strings in an ε -strongly typical set for a given length.

Proposition 8.33 *Let Σ be an alphabet, let $p \in \mathcal{P}(\Sigma)$ be a probability vector, let n be a positive integer, and let $\varepsilon > 0$ be a positive real number. It holds that*

$$(1 - \zeta_{n,\varepsilon}(p)) 2^{n(1-\varepsilon)\mathbb{H}(p)} \leq |S_{n,\varepsilon}(p)| \leq 2^{n(1+\varepsilon)\mathbb{H}(p)}, \quad (8.186)$$

for $\zeta_{n,\varepsilon}(p)$ as defined in Lemma 8.32.

Proof By Corollary 8.31, one has

$$p(a_1) \cdots p(a_n) \geq 2^{-n(1+\varepsilon)\mathbb{H}(p)} \quad (8.187)$$

for every string $a_1 \cdots a_n \in S_{n,\varepsilon}(p)$. Consequently,

$$1 \geq \sum_{a_1 \cdots a_n \in S_{n,\varepsilon}(p)} p(a_1) \cdots p(a_n) \geq |S_{n,\varepsilon}(p)| 2^{-n(1+\varepsilon)\mathbb{H}(p)}, \quad (8.188)$$

and therefore

$$|S_{n,\varepsilon}(p)| \leq 2^{n(1+\varepsilon)\mathbb{H}(p)}. \quad (8.189)$$

Along similar lines, one has

$$p(a_1) \cdots p(a_n) \leq 2^{-n(1-\varepsilon)\mathbb{H}(p)} \quad (8.190)$$

for every string $a_1 \cdots a_n \in S_{n,\varepsilon}(p)$. By Lemma 8.32, it follows that

$$1 - \zeta_{n,\varepsilon}(p) \leq \sum_{a_1 \cdots a_n \in S_{n,\varepsilon}(p)} p(a_1) \cdots p(a_n) \leq |S_{n,\varepsilon}(p)| 2^{-n(1-\varepsilon)\mathbb{H}(p)}, \quad (8.191)$$

and therefore

$$|S_{n,\varepsilon}(p)| \geq (1 - \zeta_{n,\varepsilon}(p)) 2^{n(1-\varepsilon)\mathbb{H}(p)}, \quad (8.192)$$

as required. \square

The ε -strongly typical subspaces associated with a given density operator are defined as follows.

Definition 8.34 Let \mathcal{X} be a complex Euclidean space, let $\rho \in \mathbb{D}(\mathcal{X})$ be a density operator, let $\varepsilon > 0$ be a positive real number, and let n be a positive integer. Also let

$$\rho = \sum_{a \in \Sigma} p(a) x_a x_a^* \quad (8.193)$$

be a spectral decomposition of ρ , for Σ being an alphabet, $p \in \mathcal{P}(\Sigma)$ being a probability vector, and $\{x_a : a \in \Sigma\} \subset \mathcal{X}$ being an orthonormal set of vectors. The *projection operator onto the ε -strongly typical subspace of $\mathcal{X}^{\otimes n}$ with respect to the spectral decomposition (8.193)* is defined as

$$\Lambda = \sum_{a_1 \cdots a_n \in S_{n,\varepsilon}(p)} x_{a_1} x_{a_1}^* \otimes \cdots \otimes x_{a_n} x_{a_n}^*. \quad (8.194)$$

With respect to the decomposition (8.193), the *ε -strongly typical subspace of $\mathcal{X}^{\otimes n}$* is defined as the image of Λ .

Example 8.35 Let $\Sigma = \{0, 1\}$, let $\mathcal{X} = \mathbb{C}^\Sigma$, and let $\rho = \mathbb{1}/2 \in \mathbb{D}(\mathcal{X})$. With respect to the spectral decomposition

$$\rho = \frac{1}{2} e_0 e_0^* + \frac{1}{2} e_1 e_1^*, \quad (8.195)$$

for $n = 2$, and for any choice of $\varepsilon \in (0, 1)$, one has that the corresponding projection operator onto the ε -strongly typical subspace is given by

$$\Lambda_0 = E_{0,0} \otimes E_{1,1} + E_{1,1} \otimes E_{0,0}. \quad (8.196)$$

Replacing the spectral decomposition by

$$\rho = \frac{1}{2}x_0x_0^* + \frac{1}{2}x_1x_1^*, \quad (8.197)$$

for

$$x_0 = \frac{e_0 + e_1}{\sqrt{2}} \quad \text{and} \quad x_1 = \frac{e_0 - e_1}{\sqrt{2}}, \quad (8.198)$$

one obtains the corresponding projection operator

$$\Lambda_1 = x_0x_0^* \otimes x_1x_1^* + x_1x_1^* \otimes x_0x_0^* \neq \Lambda_0. \quad (8.199)$$

Two lemmas on the output entropy of channels

The proof of the entanglement-assisted classical capacity theorem appearing at the end of the present section will make use of multiple lemmas. The two lemmas that follow concern the output entropy of channels. The first of these two lemmas will also be used in the next section of the chapter, to prove that the coherent information lower-bounds the quantum capacity of a channel.

Lemma 8.36 *Let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces, let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, let $\rho \in D(\mathcal{X})$ be a density operator, let $\varepsilon > 0$ be a positive real number, and let n be a positive integer. Also let*

$$\rho = \sum_{a \in \Sigma} p(a)x_ax_a^* \quad (8.200)$$

be a spectral decomposition of ρ , for Σ being an alphabet, $\{x_a : a \in \Sigma\} \subset \mathcal{X}$ being an orthonormal set, and $p \in \mathcal{P}(\Sigma)$ being a probability vector, let $\Lambda_{n,\varepsilon}$ denote the projection operator onto the ε -strongly typical subspace of $\mathcal{X}^{\otimes n}$ with respect to the decomposition (8.200), and let

$$\omega_{n,\varepsilon} = \frac{\Lambda_{n,\varepsilon}}{\text{Tr}(\Lambda_{n,\varepsilon})}. \quad (8.201)$$

It holds that

$$\left| \frac{\text{H}(\Phi^{\otimes n}(\omega_{n,\varepsilon}))}{n} - \text{H}(\Phi(\rho)) \right| \leq 2\varepsilon \text{H}(\rho) + \varepsilon \text{H}(\Phi(\rho)) - \frac{\log(1 - \zeta_{n,\varepsilon}(p))}{n}, \quad (8.202)$$

for $\zeta_{n,\varepsilon}(p)$ being the quantity defined in Lemma 8.32.

Proof It may be verified that the equation

$$\begin{aligned} \text{H}(\Phi(\rho)) - \frac{1}{n} \text{H}(\Phi^{\otimes n}(\omega_{n,\varepsilon})) \\ = \frac{1}{n} \text{D}(\Phi^{\otimes n}(\omega_{n,\varepsilon}) \parallel \Phi^{\otimes n}(\rho^{\otimes n})) \\ + \frac{1}{n} \text{Tr}((\Phi^{\otimes n}(\omega_{n,\varepsilon}) - \Phi(\rho)^{\otimes n}) \log(\Phi(\rho)^{\otimes n})) \end{aligned} \quad (8.203)$$

holds for every positive integer n . Bounds on the absolute values of the two terms on the right-hand side of this equation will be established separately.

The first term on the right-hand side of (8.203) is nonnegative, and an upper bound on it may be obtained from the monotonicity of the quantum relative entropy under the action of channels (Theorem 5.35). Specifically, one has

$$\begin{aligned} \frac{1}{n} \text{D}(\Phi^{\otimes n}(\omega_{n,\varepsilon}) \parallel \Phi^{\otimes n}(\rho^{\otimes n})) &\leq \frac{1}{n} \text{D}(\omega_{n,\varepsilon} \parallel \rho^{\otimes n}) \\ &= -\frac{1}{n} \log(|S_{n,\varepsilon}|) - \frac{1}{n|S_{n,\varepsilon}|} \sum_{a_1 \dots a_n \in S_{n,\varepsilon}} \log(p(a_1) \dots p(a_n)), \end{aligned} \quad (8.204)$$

where $S_{n,\varepsilon}$ denotes the set of ε -strongly typical strings of length n with respect to p . By Corollary 8.31 it holds that

$$-\frac{1}{n|S_{n,\varepsilon}|} \sum_{a_1 \dots a_n \in S_{n,\varepsilon}} \log(p(a_1) \dots p(a_n)) \leq (1 + \varepsilon) \text{H}(\rho), \quad (8.205)$$

and by Proposition 8.33, one has

$$\frac{1}{n} \log(|S_{n,\varepsilon}|) \geq \frac{\log(1 - \zeta_{n,\varepsilon}(p))}{n} + (1 - \varepsilon) \text{H}(\rho). \quad (8.206)$$

It therefore holds that

$$\frac{1}{n} \text{D}(\Phi^{\otimes n}(\omega_{n,\varepsilon}) \parallel \Phi^{\otimes n}(\rho^{\otimes n})) \leq 2\varepsilon \text{H}(\rho) - \frac{\log(1 - \zeta_{n,\varepsilon}(p))}{n}. \quad (8.207)$$

To bound the absolute value of second term on the right-hand side of (8.203), one may first define a function $\phi : \Sigma \rightarrow [0, \infty)$ as

$$\phi(a) = \begin{cases} -\text{Tr}(\Phi(x_ax_a^*) \log(\Phi(\rho))) & \text{if } p(a) > 0 \\ 0 & \text{if } p(a) = 0 \end{cases} \quad (8.208)$$

for each $a \in \Sigma$. It is evident from its specification that $\phi(a)$ is nonnegative for each $a \in \Sigma$, and is finite by virtue of the fact that

$$\text{im}(\Phi(x_ax_a^*)) \subseteq \text{im}(\Phi(\rho)) \quad (8.209)$$

for each $a \in \Sigma$ with $p(a) > 0$. Using the identity

$$\log(P^{\otimes n}) = \sum_{k=1}^n \mathbf{1}^{\otimes(k-1)} \otimes \log(P) \otimes \mathbf{1}^{\otimes(n-k)}, \quad (8.210)$$

it may be verified that

$$\begin{aligned} & \text{Tr}(\Phi^{\otimes n}(\omega_{n,\varepsilon}) \log(\Phi(\rho)^{\otimes n})) \\ &= -\frac{1}{|S_{n,\varepsilon}|} \sum_{a_1 \cdots a_n \in S_{n,\varepsilon}} (\phi(a_1) + \cdots + \phi(a_n)). \end{aligned} \quad (8.211)$$

By combining Proposition 8.30 with the observation that

$$\mathbf{H}(\Phi(\rho)) = \sum_{a \in \Sigma} p(a) \phi(a), \quad (8.212)$$

one finds that

$$\begin{aligned} & \left| \frac{1}{n} \text{Tr}((\Phi^{\otimes n}(\omega_{n,\varepsilon}) - \Phi(\rho)^{\otimes n}) \log(\Phi(\rho)^{\otimes n})) \right| \\ & \leq \frac{1}{|S_{n,\varepsilon}|} \sum_{a_1 \cdots a_n \in S_{n,\varepsilon}} \left| \mathbf{H}(\Phi(\rho)) - \frac{\phi(a_1) + \cdots + \phi(a_n)}{n} \right| \\ & \leq \varepsilon \mathbf{H}(\Phi(\rho)). \end{aligned} \quad (8.213)$$

The inequalities (8.207) and (8.213) together imply the required inequality (8.202), which completes the proof. \square

Lemma 8.37 *Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} . The function $f : D(\mathcal{X}) \rightarrow \mathbb{R}$ defined by*

$$f(\rho) = \mathbf{H}(\rho) - \mathbf{H}(\Phi(\rho)) \quad (8.214)$$

is concave.

Proof Let \mathcal{Z} be an arbitrary complex Euclidean space, and consider first the function $g : D(\mathcal{Y} \otimes \mathcal{Z}) \rightarrow \mathbb{R}$ defined as

$$g(\sigma) = \mathbf{H}(\sigma) - \mathbf{H}(\text{Tr}_{\mathcal{Z}}(\sigma)) \quad (8.215)$$

for every $\sigma \in D(\mathcal{Y} \otimes \mathcal{Z})$. An alternative expression for g is

$$g(\sigma) = -D(\sigma \| \text{Tr}_{\mathcal{Z}}(\sigma) \otimes \mathbf{1}_{\mathcal{Z}}), \quad (8.216)$$

and the concavity of g therefore follows from the joint convexity of quantum relative entropy (Corollary 5.33).

For a suitable choice of a complex Euclidean space \mathcal{Z} , let $A \in U(\mathcal{X}, \mathcal{Y} \otimes \mathcal{Z})$ be an isometry that yields a Stinespring representation of Φ :

$$\Phi(X) = \text{Tr}_{\mathcal{Z}}(AXA^*) \quad (8.217)$$

for every $X \in L(\mathcal{X})$. The function f is given by $f(\rho) = g(A\rho A^*)$ for every $\rho \in D(\mathcal{X})$, and therefore the concavity of g implies that f is concave as well. \square

An additivity lemma concerning the coherent information

Another lemma that will be used in the proof of the entanglement-assisted capacity theorem is proved below. It states that the quantity

$$\max_{\sigma \in D(\mathcal{X})} (\mathbf{H}(\sigma) + \mathbf{I}_c(\sigma; \Phi)), \quad (8.218)$$

defined for each channel $\Phi \in C(\mathcal{X}, \mathcal{Y})$, is additive with respect to tensor products. It is precisely this quantity that the entanglement-assisted classical capacity theorem establishes is equal to the entanglement-assisted classical capacity of the channel Φ .

Lemma 8.38 (Adami–Cerf) *Let $\Phi_0 \in C(\mathcal{X}_0, \mathcal{Y}_0)$ and $\Phi_1 \in C(\mathcal{X}_1, \mathcal{Y}_1)$ be channels, for complex Euclidean spaces $\mathcal{X}_0, \mathcal{X}_1, \mathcal{Y}_0$, and \mathcal{Y}_1 . It holds that*

$$\begin{aligned} & \max_{\sigma \in D(\mathcal{X}_0 \otimes \mathcal{X}_1)} (\mathbf{H}(\sigma) + \mathbf{I}_c(\sigma; \Phi_0 \otimes \Phi_1)) \\ &= \max_{\sigma_0 \in D(\mathcal{X}_0)} (\mathbf{H}(\sigma_0) + \mathbf{I}_c(\sigma_0; \Phi_0)) + \max_{\sigma_1 \in D(\mathcal{X}_1)} (\mathbf{H}(\sigma_1) + \mathbf{I}_c(\sigma_1; \Phi_1)). \end{aligned} \quad (8.219)$$

Proof Choose isometries $A_0 \in U(\mathcal{X}_0, \mathcal{Y}_0 \otimes \mathcal{Z}_0)$ and $A_1 \in U(\mathcal{X}_1, \mathcal{Y}_1 \otimes \mathcal{Z}_1)$, for an appropriate choice of complex Euclidean spaces \mathcal{Z}_0 and \mathcal{Z}_1 , so that Stinespring representations of Φ_0 and Φ_1 are obtained:

$$\Phi_0(X_0) = \text{Tr}_{\mathcal{Z}_0}(A_0 X_0 A_0^*) \quad \text{and} \quad \Phi_1(X_1) = \text{Tr}_{\mathcal{Z}_1}(A_1 X_1 A_1^*) \quad (8.220)$$

for all $X_0 \in L(\mathcal{X}_0)$ and $X_1 \in L(\mathcal{X}_1)$. The channels $\Psi_0 \in C(\mathcal{X}_0, \mathcal{Z}_0)$ and $\Psi_1 \in C(\mathcal{X}_1, \mathcal{Z}_1)$ defined as

$$\Psi_0(X_0) = \text{Tr}_{\mathcal{Y}_0}(A_0 X_0 A_0^*) \quad \text{and} \quad \Psi_1(X_1) = \text{Tr}_{\mathcal{Y}_1}(A_1 X_1 A_1^*) \quad (8.221)$$

for all $X_0 \in L(\mathcal{X}_0)$ and $X_1 \in L(\mathcal{X}_1)$ are therefore complementary to Φ_0 and Φ_1 , respectively.

Now, consider registers $\mathcal{X}_0, \mathcal{X}_1, \mathcal{Y}_0, \mathcal{Y}_1, \mathcal{Z}_0$, and \mathcal{Z}_1 corresponding to the spaces $\mathcal{X}_0, \mathcal{X}_1, \mathcal{Y}_0, \mathcal{Y}_1, \mathcal{Z}_0$, and \mathcal{Z}_1 , respectively. Let $\sigma \in D(\mathcal{X}_0 \otimes \mathcal{X}_1)$ be an arbitrary density operator. With respect to the state

$$(A_0 \otimes A_1)\sigma(A_0 \otimes A_1)^* \in D(\mathcal{Y}_0 \otimes \mathcal{Z}_0 \otimes \mathcal{Y}_1 \otimes \mathcal{Z}_1) \quad (8.222)$$

of (Y_0, Z_0, Y_1, Z_1) , one has that

$$\begin{aligned} H(\sigma) + I_c(\sigma; \Phi_0 \otimes \Phi_1) \\ = H(Y_0, Z_0, Y_1, Z_1) + H(Y_0, Y_1) - H(Z_0, Z_1). \end{aligned} \quad (8.223)$$

For every state of (Y_0, Z_0, Y_1, Z_1) , including the state (8.222), it holds that

$$\begin{aligned} H(Y_0, Z_0, Y_1, Z_1) &\leq H(Z_0, Y_1, Z_1) + H(Y_0, Z_0) - H(Z_0) \\ &\leq H(Z_0, Z_1) + H(Y_1, Z_1) - H(Z_1) + H(Y_0, Z_0) - H(Z_0); \end{aligned} \quad (8.224)$$

both inequalities follow from the strong subadditivity of the von Neumann entropy (Theorem 5.36). The subadditivity of the von Neumann entropy (Theorem 5.24) implies $H(Y_0, Y_1) \leq H(Y_0) + H(Y_1)$, and therefore

$$\begin{aligned} H(Y_0, Z_0, Y_1, Z_1) + H(Y_0, Y_1) - H(Z_0, Z_1) \\ \leq (H(Y_0, Z_0) + H(Y_0) - H(Z_0)) \\ + (H(Y_1, Z_1) + H(Y_1) - H(Z_1)). \end{aligned} \quad (8.225)$$

For $\sigma_0 = \sigma[X_0]$ and $\sigma_1 = \sigma[X_1]$, one has the equations

$$\begin{aligned} H(Y_0, Z_0) + H(Y_0) - H(Z_0) &= H(\sigma_0) + I_c(\sigma_0; \Phi_0), \\ H(Y_1, Z_1) + H(Y_1) - H(Z_1) &= H(\sigma_1) + I_c(\sigma_1; \Phi_1). \end{aligned} \quad (8.226)$$

It follows that

$$\begin{aligned} H(\sigma) + I_c(\sigma; \Phi_0 \otimes \Phi_1) \\ \leq (H(\sigma_0) + I_c(\sigma_0; \Phi_0)) + (H(\sigma_1) + I_c(\sigma_1; \Phi_1)). \end{aligned} \quad (8.227)$$

Maximizing over all $\sigma \in D(\mathcal{X}_0 \otimes \mathcal{X}_1)$, one obtains the inequality

$$\begin{aligned} \max_{\sigma \in D(\mathcal{X}_0 \otimes \mathcal{X}_1)} (H(\sigma) + I_c(\sigma; \Phi_0 \otimes \Phi_1)) \\ \leq \max_{\sigma_0 \in D(\mathcal{X}_0)} (H(\sigma_0) + I_c(\sigma_0; \Phi_0)) + \max_{\sigma_1 \in D(\mathcal{X}_1)} (H(\sigma_1) + I_c(\sigma_1; \Phi_1)). \end{aligned} \quad (8.228)$$

For the reverse inequality, it suffices to observe that

$$\begin{aligned} H(\sigma_0 \otimes \sigma_1) + I_c(\sigma_0 \otimes \sigma_1; \Phi_0 \otimes \Phi_1) \\ = H(\sigma_0) + I_c(\sigma_0; \Phi_0) + H(\sigma_1) + I_c(\sigma_1; \Phi_1) \end{aligned} \quad (8.229)$$

for every choice of $\sigma_0 \in D(\mathcal{X}_0)$ and $\sigma_1 \in D(\mathcal{X}_1)$, and therefore

$$\begin{aligned} \max_{\sigma \in D(\mathcal{X}_0 \otimes \mathcal{X}_1)} (H(\sigma) + I_c(\sigma; \Phi_0 \otimes \Phi_1)) \\ \geq \max_{\sigma_0 \in D(\mathcal{X}_0)} (H(\sigma_0) + I_c(\sigma_0; \Phi_0)) + \max_{\sigma_1 \in D(\mathcal{X}_1)} (H(\sigma_1) + I_c(\sigma_1; \Phi_1)), \end{aligned} \quad (8.230)$$

which completes the proof. \square

A lower-bound on the Holevo capacity for flat states by dense coding

Next in the sequence of lemmas needed to prove the entanglement-assisted classical capacity theorem is the following lemma, which establishes a lower bound on the entanglement-assisted Holevo capacity of a given channel. Its proof may be viewed an application of dense coding (q.v. Section 6.3.1).

Lemma 8.39 *Let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces, let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, let $\Pi \in \text{Proj}(\mathcal{X})$ be a nonzero projection operator, and let $\omega = \Pi / \text{Tr}(\Pi)$. It holds that*

$$\chi_E(\Phi) \geq H(\omega) + I_c(\omega; \Phi). \quad (8.231)$$

Proof Let $m = \text{rank}(\Pi)$, let $\mathcal{W} = \mathbb{C}^{\mathbb{Z}_m}$, let $V \in U(\mathcal{W}, \mathcal{X})$ be any isometry satisfying $VV^* = \Pi$, and let

$$\tau = \frac{1}{m} \text{vec}(V) \text{vec}(V)^* \in D(\mathcal{X} \otimes \mathcal{W}). \quad (8.232)$$

Recall the collection of discrete Weyl operators

$$\{W_{a,b} : a, b \in \mathbb{Z}_m\} \subset U(\mathcal{W}), \quad (8.233)$$

as defined in Section 4.1.2 of Chapter 4, and define a collection of unitary channels

$$\{\Psi_{a,b} : a, b \in \mathbb{Z}_m\} \subseteq C(\mathcal{W}) \quad (8.234)$$

in correspondence with these operators:

$$\Psi_{a,b}(Y) = W_{a,b} Y W_{a,b}^* \quad (8.235)$$

for each $Y \in L(\mathcal{W})$. Finally, consider the ensemble

$$\eta : \mathbb{Z}_m \times \mathbb{Z}_m \rightarrow \text{Pos}(\mathcal{X} \otimes \mathcal{W}) \quad (8.236)$$

defined as

$$\eta(a, b) = \frac{1}{m^2} (\mathbb{1}_{L(\mathcal{X})} \otimes \Psi_{a,b})(\tau), \quad (8.237)$$

for all $(a, b) \in \mathbb{Z}_m \times \mathbb{Z}_m$.

It holds that

$$\begin{aligned} H\left(\frac{1}{m^2} \sum_{a,b \in \mathbb{Z}_m} (\Phi \otimes \Psi_{a,b})(\tau)\right) \\ = H\left(\Phi(\omega) \otimes \frac{\mathbb{1}_{\mathcal{W}}}{m}\right) = H(\Phi(\omega)) + H(\omega) \end{aligned} \quad (8.238)$$

and

$$\begin{aligned} \frac{1}{m^2} \sum_{a,b \in \mathbb{Z}_n} \mathbb{H}((\Phi \otimes \Psi_{a,b})(\tau)) &= \mathbb{H}((\Phi \otimes \mathbb{1}_{L(\mathcal{W})})(\tau)) \\ &= \mathbb{H}((\Phi \otimes \mathbb{1}_{L(\mathcal{X})})(\text{vec}(\sqrt{\omega}) \text{vec}(\sqrt{\omega})^*)), \end{aligned} \quad (8.239)$$

from which it follows that

$$\chi((\Phi \otimes \mathbb{1}_{L(\mathcal{W})})(\eta)) = \mathbb{H}(\omega) + \mathbb{I}_c(\omega; \Phi). \quad (8.240)$$

Moreover, η is homogeneous on \mathcal{W} , as is evident from the fact that

$$\text{Tr}_{\mathcal{X}}(\eta(a, b)) = \frac{1}{m^3} \mathbb{1}_{\mathcal{W}} \quad (8.241)$$

for each choice of $(a, b) \in \mathbb{Z}_m \times \mathbb{Z}_m$. It therefore holds that

$$\chi_{\mathbb{E}}(\Phi) \geq \chi((\Phi \otimes \mathbb{1}_{L(\mathcal{W})})(\eta)) = \mathbb{H}(\omega) + \mathbb{I}_c(\omega; \Phi), \quad (8.242)$$

which completes the proof. \square

An upper-bound on the Holevo capacity

The final lemma needed for the proof of the entanglement-assisted classical capacity theorem establishes an upper bound on the entanglement-assisted Holevo capacity of a channel.

Lemma 8.40 *Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} . Also let \mathcal{W} be a complex Euclidean space, let Σ be an alphabet, let $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X} \otimes \mathcal{W})$ be an ensemble that is homogeneous on \mathcal{W} , and let*

$$\sigma = \sum_{a \in \Sigma} \text{Tr}_{\mathcal{W}}(\eta(a)). \quad (8.243)$$

It holds that

$$\chi((\Phi \otimes \mathbb{1}_{L(\mathcal{W})})(\eta)) \leq \mathbb{H}(\sigma) + \mathbb{I}_c(\sigma; \Phi). \quad (8.244)$$

Proof Assume that \mathcal{Z} is a complex Euclidean space and $A \in U(\mathcal{X}, \mathcal{Y} \otimes \mathcal{Z})$ is an isometry for which

$$\Phi(X) = \text{Tr}_{\mathcal{Z}}(AXA^*) \quad (8.245)$$

for all $X \in L(\mathcal{X})$. The channel $\Psi \in C(\mathcal{X}, \mathcal{Z})$ defined by

$$\Psi(X) = \text{Tr}_{\mathcal{Y}}(AXA^*) \quad (8.246)$$

for all $X \in L(\mathcal{X})$ is therefore complementary to Φ , so that

$$\mathbb{I}_c(\sigma; \Phi) = \mathbb{H}(\Phi(\sigma)) - \mathbb{H}(\Psi(\sigma)). \quad (8.247)$$

It therefore suffices to prove that

$$\chi((\Phi \otimes \mathbb{1}_{L(\mathcal{W})})(\eta)) \leq \mathbb{H}(\sigma) + \mathbb{H}(\Phi(\sigma)) - \mathbb{H}(\Psi(\sigma)). \quad (8.248)$$

By the assumption that η is homogeneous on \mathcal{W} , Proposition 8.12 implies that there must exist a complex Euclidean space \mathcal{V} , a collection of channels

$$\{\Xi_a : a \in \Sigma\} \subseteq C(\mathcal{V}, \mathcal{X}), \quad (8.249)$$

a unit vector $u \in \mathcal{V} \otimes \mathcal{W}$, and a probability vector $p \in \mathcal{P}(\Sigma)$ such that

$$\eta(a) = p(a)(\Xi_a \otimes \mathbb{1}_{L(\mathcal{W})})(uu^*) \quad (8.250)$$

for every $a \in \Sigma$. Assume hereafter that such a choice for these objects has been fixed, and define states $\tau \in D(\mathcal{W})$ and $\xi \in D(\mathcal{V})$ as

$$\tau = \text{Tr}_{\mathcal{V}}(uu^*) \quad \text{and} \quad \xi = \text{Tr}_{\mathcal{W}}(uu^*). \quad (8.251)$$

It may be noted that

$$\sigma = \sum_{a \in \Sigma} p(a)\Xi_a(\xi). \quad (8.252)$$

Let \mathcal{U} be a complex Euclidean space such that $\dim(\mathcal{U}) = \dim(\mathcal{V} \otimes \mathcal{X})$, and select a collection of isometries $\{B_a : a \in \Sigma\} \subset U(\mathcal{V}, \mathcal{X} \otimes \mathcal{U})$ satisfying

$$\Xi_a(V) = \text{Tr}_{\mathcal{U}}(B_a V B_a^*) \quad (8.253)$$

for every $V \in L(\mathcal{V})$.

Assume momentarily that $a \in \Sigma$ has been fixed, and define a unit vector

$$v_a = (A \otimes \mathbb{1}_{\mathcal{U}} \otimes \mathbb{1}_{\mathcal{W}})(B_a \otimes \mathbb{1}_{\mathcal{W}})u \in \mathcal{Y} \otimes \mathcal{Z} \otimes \mathcal{U} \otimes \mathcal{W}. \quad (8.254)$$

Let \mathcal{Y} , \mathcal{Z} , \mathcal{U} , and \mathcal{W} be registers having corresponding complex Euclidean spaces \mathcal{Y} , \mathcal{Z} , \mathcal{U} , and \mathcal{W} , and consider the situation in which the compound register $(\mathcal{Y}, \mathcal{Z}, \mathcal{U}, \mathcal{W})$ is in the pure state $v_a v_a^*$. The following equalities may be verified:

$$\begin{aligned} \mathbb{H}(\mathcal{W}) &= \mathbb{H}(\tau), \\ \mathbb{H}(\mathcal{Y}, \mathcal{W}) &= \mathbb{H}((\Phi \Xi_a \otimes \mathbb{1}_{L(\mathcal{W})})(uu^*)), \\ \mathbb{H}(\mathcal{U}, \mathcal{W}) &= \mathbb{H}(\mathcal{Y}, \mathcal{Z}) = \mathbb{H}(\Xi_a(\xi)), \\ \mathbb{H}(\mathcal{Y}, \mathcal{U}, \mathcal{W}) &= \mathbb{H}(\mathcal{Z}) = \mathbb{H}((\Psi \Xi_a)(\xi)). \end{aligned} \quad (8.255)$$

By the strong subadditivity of the von Neumann entropy (Theorem 5.36), it holds that

$$\mathbb{H}(\mathcal{W}) - \mathbb{H}(\mathcal{Y}, \mathcal{W}) \leq \mathbb{H}(\mathcal{U}, \mathcal{W}) - \mathbb{H}(\mathcal{Y}, \mathcal{U}, \mathcal{W}), \quad (8.256)$$

and therefore

$$\mathbb{H}(\tau) - \mathbb{H}((\Phi \Xi_a \otimes \mathbb{1}_{L(\mathcal{W})})(uu^*)) \leq \mathbb{H}(\Xi_a(\xi)) - \mathbb{H}((\Psi \Xi_a)(\xi)). \quad (8.257)$$

Finally, in accordance with the probability vector p , one may average the two sides of (8.257) over all $a \in \Sigma$ and apply Lemma 8.37, obtaining

$$\begin{aligned} \mathbb{H}(\tau) - \sum_{a \in \Sigma} p(a) \mathbb{H}((\Phi \Xi_a \otimes \mathbb{1}_{L(\mathcal{W})})(uu^*)) \\ \leq \sum_{a \in \Sigma} p(a) (\mathbb{H}(\Xi_a(\xi)) - \mathbb{H}((\Psi \Xi_a)(\xi))) \leq \mathbb{H}(\sigma) - \mathbb{H}(\Psi(\sigma)). \end{aligned} \quad (8.258)$$

By the subadditivity of the von Neumann entropy (Proposition 5.9) one has

$$\mathbb{H}\left(\sum_{a \in \Sigma} p(a) (\Phi \Xi_a \otimes \mathbb{1}_{L(\mathcal{W})})(uu^*)\right) \leq \mathbb{H}(\Phi(\sigma)) + \mathbb{H}(\tau). \quad (8.259)$$

The inequality (8.248) follows from (8.258) and (8.259), which completes the proof. \square

The entanglement-assisted classical capacity theorem

Finally, the entanglement-assisted classical capacity theorem will be stated, and proved through the use of the lemmas presented above.

Theorem 8.41 (Entanglement-assisted classical capacity theorem) *Let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces and let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ be a channel. It holds that*

$$C_E(\Phi) = \max_{\sigma \in \mathcal{D}(\mathcal{X})} (\mathbb{H}(\sigma) + I_C(\sigma; \Phi)). \quad (8.260)$$

Proof By applying Lemma 8.40, followed by Lemma 8.38, one may conclude that

$$\begin{aligned} \chi_E(\Phi^{\otimes n}) &\leq \max_{\sigma \in \mathcal{D}(\mathcal{X}^{\otimes n})} (\mathbb{H}(\sigma) + I_C(\sigma; \Phi^{\otimes n})) \\ &= n \max_{\sigma \in \mathcal{D}(\mathcal{X})} (\mathbb{H}(\sigma) + I_C(\sigma; \Phi)) \end{aligned} \quad (8.261)$$

for every positive integer n . By Theorem 8.28, it therefore follows that

$$C_E(\Phi) = \lim_{n \rightarrow \infty} \frac{\chi_E(\Phi^{\otimes n})}{n} \leq \max_{\sigma \in \mathcal{D}(\mathcal{X})} (\mathbb{H}(\sigma) + I_C(\sigma; \Phi)). \quad (8.262)$$

For the reverse inequality, one may first choose a complex Euclidean space \mathcal{Z} and an isometry $A \in \mathcal{U}(\mathcal{X}, \mathcal{Y} \otimes \mathcal{Z})$ such that

$$\Phi(X) = \text{Tr}_{\mathcal{Z}}(AXA^*) \quad (8.263)$$

for all $X \in \mathcal{L}(\mathcal{X})$. It holds that the channel $\Psi \in \mathcal{C}(\mathcal{X}, \mathcal{Z})$, defined by

$$\Psi(X) = \text{Tr}_{\mathcal{Y}}(AXA^*) \quad (8.264)$$

for all $X \in \mathcal{L}(\mathcal{X})$, is complementary to Φ , so that Proposition 8.17 implies

$$I_C(\sigma; \Phi) = \mathbb{H}(\Phi(\sigma)) - \mathbb{H}(\Psi(\sigma)) \quad (8.265)$$

for all $\sigma \in \mathcal{D}(\mathcal{X})$.

Next, let $\sigma \in \mathcal{D}(\mathcal{X})$ be any density operator, let $\delta > 0$ be chosen arbitrarily, and choose $\varepsilon > 0$ to be sufficiently small so that

$$(7\mathbb{H}(\sigma) + \mathbb{H}(\Phi(\sigma)) + \mathbb{H}(\Psi(\sigma)))\varepsilon < \delta. \quad (8.266)$$

Also let

$$\omega_{n,\varepsilon} = \frac{\Lambda_{n,\varepsilon}}{\text{Tr}(\Lambda_{n,\varepsilon})} \quad (8.267)$$

for $\Lambda_{n,\varepsilon}$ denoting the ε -strongly typical projection with respect to any fixed spectral decomposition of σ , for each positive integer n .

By Lemma 8.36, one may conclude that the following three inequalities hold simultaneously for all but finitely many positive integers n :

$$\begin{aligned} \mathbb{H}(\sigma) - \frac{\mathbb{H}(\omega_{n,\varepsilon})}{n} &\leq 3\mathbb{H}(\sigma)\varepsilon + \delta, \\ \mathbb{H}(\Phi(\sigma)) - \frac{\mathbb{H}(\Phi^{\otimes n}(\omega_{n,\varepsilon}))}{n} &\leq (2\mathbb{H}(\sigma) + \mathbb{H}(\Phi(\sigma)))\varepsilon + \delta, \\ \frac{\mathbb{H}(\Phi^{\otimes n}(\omega_{n,\varepsilon}))}{n} - \mathbb{H}(\Psi(\sigma)) &\leq (2\mathbb{H}(\sigma) + \mathbb{H}(\Psi(\sigma)))\varepsilon + \delta. \end{aligned} \quad (8.268)$$

By Lemma 8.39, it therefore holds that

$$\begin{aligned} \frac{\chi_E(\Phi^{\otimes n})}{n} &\geq \frac{1}{n} (\mathbb{H}(\omega_{n,\varepsilon}) + \mathbb{H}(\Phi^{\otimes n}(\omega_{n,\varepsilon})) - \mathbb{H}(\Psi^{\otimes n}(\omega_{n,\varepsilon}))) \\ &\geq \mathbb{H}(\sigma) + \mathbb{H}(\Phi(\sigma)) - \mathbb{H}(\Psi(\sigma)) - 4\delta \end{aligned} \quad (8.269)$$

for all but finitely many positive integers n , and consequently

$$C_E(\Phi) = \lim_{n \rightarrow \infty} \frac{\chi_E(\Phi^{\otimes n})}{n} \geq \mathbb{H}(\sigma) + \mathbb{H}(\Phi(\sigma)) - \mathbb{H}(\Psi(\sigma)) - 4\delta. \quad (8.270)$$

As this inequality holds for all $\delta > 0$, one has

$$C_E(\Phi) \geq \mathbb{H}(\sigma) + \mathbb{H}(\Phi(\sigma)) - \mathbb{H}(\Psi(\sigma)) = \mathbb{H}(\sigma) + I_C(\sigma; \Phi), \quad (8.271)$$

and maximizing over all $\sigma \in \mathcal{D}(\mathcal{X})$ completes the proof. \square

8.2 Quantum information over quantum channels

This section is concerned with the capacity of quantum channels to transmit quantum information from a sender to a receiver. Along similar lines to the classical capacities considered in the previous section, one may consider the quantum capacity of a channel both when the sender and receiver share prior entanglement, used to assist with the information transmission, and when they do not.

As it turns out, the capacity of a channel to transmit quantum information with the assistance of entanglement is, in all cases, equal to one-half of the entanglement-assisted classical capacity of the same channel. This fact is proved below through a combination of the teleportation and dense coding protocols discussed in Section 6.3.1. As the entanglement-assisted classical capacity has already been characterized by Theorem 8.41, a characterization of the capacity of a quantum channel to transmit quantum information with the assistance of entanglement follows directly. For this reason, the primary focus of the section is on an analysis of the capacity of quantum channels to transmit quantum information without the assistance of entanglement.

The first subsection below presents a definition of the quantum capacity of a channel, together with the closely related notion of a channel's capacity to generate shared entanglement. The second subsection presents a proof of the quantum capacity theorem, which characterizes the capacity of a given channel to transmit quantum information.

8.2.1 Definitions of quantum capacity and related notions

Definitions of the *quantum capacity* and *entanglement-generation capacity* of a channel are presented below, and it is proved that the two quantities coincide. The *entanglement-assisted quantum capacity* of a channel is also defined, and its simple relationship to the entanglement-assisted classical capacity of a channel is clarified.

The quantum capacity of a channel

Informally speaking, the quantum capacity of a channel is the number of qubits, on average, that can be accurately transmitted with each use of that channel. Like the capacities discussed in the previous section, the quantum capacity of a channel is defined in information-theoretic terms, referring to a situation in which an asymptotically large number of channel uses, acting on a collection of possibly entangled registers, is made available.

The definition of quantum capacity that follows makes use of the same notions of an emulation of one channel by another (Definition 8.1) and of an ε -approximation of one channel by another (Definition 8.2) that were used in the previous section.

Definition 8.42 (Quantum capacity of a channel) Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} , and also let $\mathcal{Z} = \mathbb{C}^\Gamma$ for $\Gamma = \{0, 1\}$ denoting the binary alphabet.

1. A value $\alpha \geq 0$ is an *achievable rate* for the transmission of quantum information through Φ if (i) $\alpha = 0$, or (ii) $\alpha > 0$ and the following holds for every choice of a positive real number $\varepsilon > 0$: for all but finitely many positive integers n , and for $m = \lfloor \alpha n \rfloor$, the channel $\Phi^{\otimes n}$ emulates an ε -approximation to the identity channel $\mathbb{1}_{L(\mathcal{Z})}^{\otimes m}$.
2. The *quantum capacity* of Φ , which is denoted $Q(\Phi)$, is defined as the supremum of all achievable rates for quantum information transmission through Φ .

The argument through which Proposition 8.4 in the previous section was proved yields the following analogous proposition for the quantum capacity.

Proposition 8.43 Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} . It holds that $Q(\Phi^{\otimes k}) = k Q(\Phi)$ for every positive integer k .

The entanglement generation capacity of a channel

The *entanglement generation capacity* of a channel is defined in a similar way to the quantum capacity, except that the associated task is more narrowly focused: by means of multiple, independent uses of a channel, a sender and receiver aim to establish a state, shared between them, having high fidelity with a maximally entangled state.

Definition 8.44 (Entanglement generation capacity of a channel) Let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces, let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, and let $\mathcal{Z} = \mathbb{C}^\Gamma$ for $\Gamma = \{0, 1\}$ denoting the binary alphabet.

1. A value $\alpha \geq 0$ is an *achievable rate* for entanglement generation through Φ if (i) $\alpha = 0$, or (ii) $\alpha > 0$ and the following holds for every positive real number $\varepsilon > 0$: for all but finitely many positive integers n , and for $m = \lfloor \alpha n \rfloor$, there exists a state $\rho \in D(\mathcal{X}^{\otimes n} \otimes \mathcal{Z}^{\otimes m})$ and a channel $\Xi \in C(\mathcal{Y}^{\otimes n}, \mathcal{Z}^{\otimes m})$ such that

$$F\left(2^{-m} \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m}) \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m})^*, (\Xi \Phi^{\otimes n} \otimes \mathbb{1}_{L(\mathcal{Z})}^{\otimes m})(\rho)\right) \geq 1 - \varepsilon. \quad (8.272)$$

2. The *entanglement generation capacity* of Φ , denoted $Q_{\text{EG}}(\Phi)$, is defined as the supremum of all achievable rates for entanglement generation through Φ .

Remark For any choice of complex Euclidean spaces \mathcal{X} and \mathcal{Y} , a unit vector $y \in \mathcal{Y}$, and a channel $\Psi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$, the maximum value for the fidelity $F(yy^*, \Psi(\rho))$ over $\rho \in \mathcal{D}(\mathcal{X})$ is achieved when ρ is a pure state. It follows from this observation that the quantity $Q_{\text{EG}}(\Phi)$ would not change if the states $\rho \in \mathcal{D}(\mathcal{X}^{\otimes n} \otimes \mathcal{Z}^{\otimes m})$ considered in the specification of achievable rates in Definition 8.44 are constrained to be pure states.

Equivalence of quantum capacity and entanglement generation capacity

The task associated with entanglement generation capacity would seem to be more specialized than the one associated with quantum capacity. That is, the emulation of a close approximation to an identity channel evidently allows a sender and receiver to generate a shared state having high fidelity with a maximally entangled state, but it is not immediate that the ability of a channel to generate near-maximally entangled states should allow it to accurately transmit quantum information at a similar rate. One may note, in particular, that the teleportation protocol discussed in Section 6.3.1 is not immediately applicable in this situation, as the protocol requires classical communication that must be considered in the calculation of transmission rates. Nevertheless, the relationship between entanglement generation and identity channel emulation provided by the following theorem allows one to prove that the quantum capacity and entanglement generation capacity of any given channel do indeed coincide.

Theorem 8.45 *Let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces, let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ be a channel, and let $u \in \mathcal{X} \otimes \mathcal{Y}$ be a unit vector. Also let $n = \dim(\mathcal{Y})$ and let $\delta \geq 0$ be a nonnegative real number such that*

$$F\left(\frac{1}{n} \text{vec}(\mathbb{1}_{\mathcal{Y}}) \text{vec}(\mathbb{1}_{\mathcal{Y}})^*, (\Phi \otimes \mathbb{1}_{L(\mathcal{Y})})(uu^*)\right) \geq 1 - \delta. \quad (8.273)$$

For any complex Euclidean space \mathcal{Z} satisfying $\dim(\mathcal{Z}) \leq n/2$, it holds that Φ emulates an ε -approximation to the identity channel $\mathbb{1}_{L(\mathcal{Z})}$ for $\varepsilon = 4\delta^{\frac{1}{4}}$.

Proof Let $A \in L(\mathcal{Y}, \mathcal{X})$ be the operator defined by the equation $\text{vec}(A) = u$, let $r = \text{rank}(A)$, and let

$$A = \sum_{k=1}^r \sqrt{p_k} x_k y_k^* \quad (8.274)$$

be a singular value decomposition of A , so that (p_1, \dots, p_r) is a probability

vector and $\{x_1, \dots, x_r\} \subset \mathcal{X}$ and $\{y_1, \dots, y_r\} \subset \mathcal{Y}$ are orthonormal sets. Also define $W \in L(\mathcal{Y}, \mathcal{X})$ as

$$W = \sum_{k=1}^r x_k y_k^*, \quad (8.275)$$

and define a unit vector $v \in \mathcal{X} \otimes \mathcal{Y}$ as

$$v = \frac{1}{\sqrt{r}} \text{vec}(W). \quad (8.276)$$

By the monotonicity of the fidelity function under partial tracing, one has

$$\begin{aligned} \frac{1}{\sqrt{n}} \sum_{k=1}^r \sqrt{p_k} &= F\left(\frac{1}{n} \mathbb{1}_{\mathcal{Y}}, \text{Tr}_{\mathcal{X}}(uu^*)\right) \\ &\geq F\left(\frac{1}{n} \text{vec}(\mathbb{1}_{\mathcal{Y}}) \text{vec}(\mathbb{1}_{\mathcal{Y}})^*, (\Phi \otimes \mathbb{1}_{L(\mathcal{Y})})(uu^*)\right) \geq 1 - \delta, \end{aligned} \quad (8.277)$$

and therefore

$$F(uu^*, vv^*) = \frac{1}{\sqrt{r}} \sum_{k=1}^r \sqrt{p_k} \geq \frac{1}{\sqrt{n}} \sum_{k=1}^r \sqrt{p_k} \geq 1 - \delta. \quad (8.278)$$

Consequently, by Theorems 3.27 and 3.29, one has

$$\begin{aligned} &F\left(\frac{1}{n} \text{vec}(\mathbb{1}_{\mathcal{Y}}) \text{vec}(\mathbb{1}_{\mathcal{Y}})^*, (\Phi \otimes \mathbb{1}_{L(\mathcal{Y})})(vv^*)\right) + 1 \\ &\geq F\left(\frac{1}{n} \text{vec}(\mathbb{1}_{\mathcal{Y}}) \text{vec}(\mathbb{1}_{\mathcal{Y}})^*, (\Phi \otimes \mathbb{1}_{L(\mathcal{Y})})(uu^*)\right)^2 + F(vv^*, uu^*)^2 \\ &\geq 2(1 - \delta)^2, \end{aligned} \quad (8.279)$$

and therefore

$$F\left(\frac{1}{n} \text{vec}(\mathbb{1}_{\mathcal{Y}}) \text{vec}(\mathbb{1}_{\mathcal{Y}})^*, (\Phi \otimes \mathbb{1}_{L(\mathcal{Y})})(vv^*)\right) \geq 1 - 4\delta. \quad (8.280)$$

Next, define a projection operator $\Pi_r = W^*W \in \text{Proj}(\mathcal{Y})$ and define $\mathcal{V}_r = \text{im}(\Pi_r)$. For each choice of k beginning with r and decreasing to 1, choose $w_k \in \mathcal{V}_k$ to be a unit vector that minimizes the quantity

$$\alpha_k = \langle w_k w_k^*, \Phi(W w_k w_k^* W^*) \rangle, \quad (8.281)$$

and define

$$\mathcal{V}_{k-1} = \{z \in \mathcal{V}_k : \langle w_k, z \rangle = 0\}. \quad (8.282)$$

Observe that $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_r$ and that $\{w_1, \dots, w_k\}$ is an orthonormal

basis for \mathcal{V}_k , for each $k \in \{1, \dots, r\}$. In particular, it holds that

$$v = \frac{1}{\sqrt{r}}(W \otimes \mathbf{1}_{\mathcal{Y}}) \text{vec}(\Pi_r) = \frac{1}{\sqrt{r}} \sum_{k=1}^r W w_k \otimes \bar{w}_k. \quad (8.283)$$

At this point, a calculation reveals that

$$\begin{aligned} & \mathbb{F}\left(\frac{1}{n} \text{vec}(\mathbf{1}_{\mathcal{Y}}) \text{vec}(\mathbf{1}_{\mathcal{Y}})^*, (\Phi \otimes \mathbf{1}_{L(\mathcal{Y})})(vv^*)\right)^2 \\ &= \frac{1}{nr} \sum_{j,k \in \{1, \dots, r\}} \langle w_j w_k^*, \Phi(W w_j w_k^* W^*) \rangle. \end{aligned} \quad (8.284)$$

By the complete positivity of Φ , one may conclude that

$$\begin{aligned} & |\langle w_j w_k^*, \Phi(W w_j w_k^* W^*) \rangle| \\ & \leq \sqrt{\langle w_j w_j^*, \Phi(W w_j w_j^* W^*) \rangle} \sqrt{\langle w_k w_k^*, \Phi(W w_k w_k^* W^*) \rangle} \\ & = \sqrt{\alpha_j \alpha_k}, \end{aligned} \quad (8.285)$$

for each choice of $j, k \in \{1, \dots, r\}$. Therefore, by the triangle inequality, it holds that

$$\mathbb{F}\left(\frac{1}{n} \text{vec}(\mathbf{1}_{\mathcal{Y}}) \text{vec}(\mathbf{1}_{\mathcal{Y}})^*, (\Phi \otimes \mathbf{1}_{L(\mathcal{Y})})(vv^*)\right) \leq \frac{1}{\sqrt{nr}} \sum_{k=1}^r \sqrt{\alpha_k}. \quad (8.286)$$

Applying the Cauchy–Schwarz inequality, one obtains

$$\frac{1}{\sqrt{nr}} \sum_{k=1}^r \sqrt{\alpha_k} \leq \sqrt{\frac{1}{n} \sum_{k=1}^r \alpha_k}, \quad (8.287)$$

and therefore

$$\frac{1}{n} \sum_{k=1}^r \alpha_k \geq (1 - 4\delta)^2 \geq 1 - 8\delta. \quad (8.288)$$

Now let

$$m = \max\{k \in \{1, \dots, r\} : \alpha_k \geq 1 - 16\delta\}. \quad (8.289)$$

It follows from (8.288) that

$$1 - 8\delta \leq \frac{m}{n} + \frac{n-m}{n}(1 - 16\delta), \quad (8.290)$$

and therefore $m \geq n/2$. By the definition of the values $\alpha_1, \dots, \alpha_r$, one may conclude that

$$\langle w w^*, \Phi(W w w^* W^*) \rangle \geq 1 - 16\delta \quad (8.291)$$

for every unit vector $w \in \mathcal{V}_m$.

Finally, let $V \in \mathcal{U}(\mathcal{Z}, \mathcal{Y})$ be any isometry for which $\text{im}(V) \subseteq \mathcal{V}_m$. Such an isometry exists by the assumption that $\dim(\mathcal{Z}) \leq n/2$ together with the fact that $n/2 \leq m = \dim(\mathcal{V}_m)$. Let $\Xi_{\mathbb{E}} \in \mathcal{C}(\mathcal{Z}, \mathcal{X})$ and $\Xi_{\mathbb{D}} \in \mathcal{C}(\mathcal{Y}, \mathcal{Z})$ be channels of the form

$$\begin{aligned} \Xi_{\mathbb{E}}(Z) &= W V Z V^* W^* + \Psi_{\mathbb{E}}(Z), \\ \Xi_{\mathbb{D}}(Y) &= V^* Y V + \Psi_{\mathbb{D}}(Y), \end{aligned} \quad (8.292)$$

for all $Z \in L(\mathcal{Z})$ and $Y \in L(\mathcal{Y})$, where $\Psi_{\mathbb{E}} \in \mathcal{CP}(\mathcal{Z}, \mathcal{X})$ and $\Psi_{\mathbb{D}} \in \mathcal{CP}(\mathcal{Y}, \mathcal{Z})$ are completely positive maps that cause $\Xi_{\mathbb{E}}$ and $\Xi_{\mathbb{D}}$ to be trace preserving. For every unit vector $z \in \mathcal{Z}$ it holds that

$$\begin{aligned} & \langle z z^*, (\Xi_{\mathbb{D}} \Phi \Xi_{\mathbb{E}})(z z^*) \rangle \\ & \geq \langle V z z^* V^*, \Phi(W V z z^* V^* W^*) \rangle \geq 1 - 16\delta, \end{aligned} \quad (8.293)$$

and therefore

$$\|z z^* - (\Xi_{\mathbb{D}} \Phi \Xi_{\mathbb{E}})(z z^*)\|_1 \leq 8\sqrt{\delta} \quad (8.294)$$

by one of the Fuchs–van de Graaf inequalities (Theorem 3.33). Applying Theorem 3.56, one therefore finds that

$$\|\|\Xi_{\mathbb{D}} \Phi \Xi_{\mathbb{E}} - \mathbf{1}_{L(\mathcal{Z})}\|\|_1 \leq 4\delta^{\frac{1}{4}}, \quad (8.295)$$

which completes the proof. \square

Theorem 8.46 *Let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} . The entanglement generation capacity and the quantum capacity of Φ are equal: $\mathbb{Q}(\Phi) = \mathbb{Q}_{\text{EG}}(\Phi)$.*

Proof It will first be proved that $\mathbb{Q}(\Phi) \leq \mathbb{Q}_{\text{EG}}(\Phi)$, which is straightforward. If the quantum capacity of Φ is zero, there is nothing to prove, so it will be assumed that $\mathbb{Q}(\Phi) > 0$. Let $\alpha > 0$ be an achievable rate for quantum information transmission through Φ , and let $\varepsilon > 0$ be chosen arbitrarily.

Setting $\Gamma = \{0, 1\}$ and $\mathcal{Z} = \mathbb{C}^{\Gamma}$, one therefore has that the channel $\Phi^{\otimes n}$ emulates an ε -approximation to the identity channel $\mathbf{1}_{L(\mathcal{Z})}^{\otimes m}$ for all but finitely many positive integers n and for $m = \lfloor \alpha n \rfloor$. That is, for all but finitely many positive integers n , and for $m = \lfloor \alpha n \rfloor$, there must exist channels $\Xi_{\mathbb{E}} \in \mathcal{C}(\mathcal{Z}^{\otimes m}, \mathcal{X}^{\otimes n})$ and $\Xi_{\mathbb{D}} \in \mathcal{C}(\mathcal{Y}^{\otimes n}, \mathcal{Z}^{\otimes m})$ such that

$$\|\|\Xi_{\mathbb{D}} \Phi^{\otimes n} \Xi_{\mathbb{E}} - \mathbf{1}_{L(\mathcal{Z})}^{\otimes m}\|\|_1 < \varepsilon. \quad (8.296)$$

Supposing that n and m are positive integers for which such channels exist, one may consider the density operators

$$\tau = 2^{-m} \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m}) \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m})^* \quad \text{and} \quad \rho = (\Xi_{\mathbb{E}} \otimes \mathbb{1}_{L(\mathcal{Z})}^{\otimes m})(\tau), \quad (8.297)$$

along with the channel $\Xi = \Xi_{\mathbb{D}}$. One of the Fuchs–van de Graaf inequalities (Theorem 3.33) implies that

$$\begin{aligned} F\left(\tau, (\Xi \Phi^{\otimes n} \otimes \mathbb{1}_{L(\mathcal{Z})}^{\otimes m})(\rho)\right) &= F\left(\tau, (\Xi_{\mathbb{D}} \Phi^{\otimes n} \Xi_{\mathbb{E}} \otimes \mathbb{1}_{L(\mathcal{Z})}^{\otimes m})(\tau)\right) \\ &\geq 1 - \frac{1}{2} \left\| (\Xi_{\mathbb{D}} \Phi^{\otimes n} \Xi_{\mathbb{E}} \otimes \mathbb{1}_{L(\mathcal{Z})}^{\otimes m})(\tau) - \tau \right\|_1 > 1 - \frac{\varepsilon}{2}. \end{aligned} \quad (8.298)$$

Because this is so for all but finitely many positive integers n and for $m = \lfloor \alpha n \rfloor$, it holds that α is an achievable rate for entanglement generation through Φ . Taking the supremum over all achievable rates α for quantum communication through Φ , one obtains $Q(\Phi) \leq Q_{\text{EG}}(\Phi)$.

It remains to prove that $Q_{\text{EG}}(\Phi) \leq Q(\Phi)$. As for the reverse inequality just proved, there is nothing to prove if $Q_{\text{EG}}(\Phi) = 0$, so it will be assumed that $Q_{\text{EG}}(\Phi) > 0$. Let $\alpha > 0$ be an achievable rate for entanglement generation through Φ and let $\beta \in (0, \alpha)$ be chosen arbitrarily. It will be proved that β is an achievable rate for quantum communication through Φ . The required relation $Q_{\text{EG}}(\Phi) \leq Q(\Phi)$ follows by taking the supremum over all achievable rates α for entanglement generation through Φ and over all $\beta \in (0, \alpha)$.

Let $\varepsilon > 0$ be chosen arbitrarily and let $\delta = \varepsilon^4/256$, so that $\varepsilon = 4\delta^{1/4}$. For all but finitely many positive integers n , and for $m = \lfloor \alpha n \rfloor$, there exists a state $\rho \in D(\mathcal{X}^{\otimes n} \otimes \mathcal{Z}^{\otimes m})$ and a channel $\Xi \in C(\mathcal{Y}^{\otimes n}, \mathcal{Z}^{\otimes m})$ such that

$$F(2^{-m} \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m}) \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m})^*, (\Xi \Phi^{\otimes n} \otimes \mathbb{1}_{L(\mathcal{Z})}^{\otimes m})(\rho)) \geq 1 - \delta. \quad (8.299)$$

Note that the existence of a state ρ for which (8.299) holds implies the existence of a pure state $\rho = uu^*$ for which the same inequality holds, by virtue of the fact that the function

$$\begin{aligned} \rho \mapsto F(2^{-m} \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m}) \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m})^*, (\Xi \Phi^{\otimes n} \otimes \mathbb{1}_{L(\mathcal{Z})}^{\otimes m})(\rho))^2 \\ = \langle 2^{-m} \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m}) \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m})^*, (\Xi \Phi^{\otimes n} \otimes \mathbb{1}_{L(\mathcal{Z})}^{\otimes m})(\rho) \rangle \end{aligned} \quad (8.300)$$

must achieve its maximum value (over all density operators) on a pure state. By Theorem 8.45, it follows that $\Phi^{\otimes n}$ emulates an ε -approximation to the identity channel $\mathbb{1}_{L(\mathcal{Z})}^{\otimes k}$ for $k = m - 1$.

Under the assumption $n \geq 1/(\alpha - \beta)$, one has that $\beta n \leq \alpha n - 1$. Thus, for all but finitely many positive integers n and for $k = \lfloor \beta n \rfloor$, it holds that $\Phi^{\otimes n}$ emulates an ε -approximation to the identity channel $\mathbb{1}_{L(\mathcal{Z})}^{\otimes k}$. As $\varepsilon > 0$ has been chosen arbitrarily, it follows that β is an achievable rate for quantum communication through Φ , which completes the proof. \square

The entanglement-assisted quantum capacity of a channel

The entanglement-assisted quantum capacity of a channel, which will be proved is equal to one-half of its entanglement-assisted classical capacity, may be formally defined as follows.

Definition 8.47 (Entanglement-assisted quantum capacity of a channel) Let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces and let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel. Also let $\Gamma = \{0, 1\}$ denote the binary alphabet, and let $\mathcal{Z} = \mathbb{C}^{\Gamma}$.

1. A value $\alpha \geq 0$ is an *achievable rate* for entanglement-assisted quantum information transmission through Φ if (i) $\alpha = 0$, or (ii) $\alpha > 0$ and the following holds for every choice of a positive real number $\varepsilon > 0$: for all but finitely many positive integers n , and for $m = \lfloor \alpha n \rfloor$, the channel $\Phi^{\otimes n}$ emulates an ε -approximation to the identity channel $\mathbb{1}_{L(\mathcal{Z})}^{\otimes m}$ with the assistance of entanglement.
2. The *entanglement-assisted quantum capacity* of Φ , denoted $Q_{\text{E}}(\Phi)$, is the supremum of all achievable rates for entanglement-assisted quantum information transmission through Φ .

Proposition 8.48 Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} . It holds that

$$Q_{\text{E}}(\Phi) = \frac{C_{\text{E}}(\Phi)}{2}. \quad (8.301)$$

Proof Assume α is an achievable rate for entanglement-assisted classical communication through Φ . It will be proved that $\alpha/2$ is an achievable rate for entanglement-assisted quantum information transmission through Φ . Taking the supremum over all achievable rates α for entanglement-assisted classical communication through Φ , one obtains

$$Q_{\text{E}}(\Phi) \geq \frac{C_{\text{E}}(\Phi)}{2}. \quad (8.302)$$

As the case $\alpha = 0$ is trivial, it will be assumed that $\alpha > 0$.

Suppose n and $m = \lfloor \alpha n \rfloor$ are positive integers and $\varepsilon > 0$ is a positive real number such that $\Phi^{\otimes n}$ emulates an ε -approximation to the channel $\Delta^{\otimes m}$, where $\Delta \in C(\mathcal{Z})$ denotes the completely dephasing channel as usual. Let $k = \lfloor m/2 \rfloor$, and consider the maximally entangled state

$$\tau = 2^{-k} \text{vec}(\mathbb{1}_{\mathcal{Z}^k}) \text{vec}(\mathbb{1}_{\mathcal{Z}^k})^*. \quad (8.303)$$

By tensoring τ with the state ξ used for the emulation of an ε -approximation to $\Delta^{\otimes m}$ by $\Phi^{\otimes n}$, one may define a new channel $\Psi \in C(\mathcal{Z}^{\otimes k})$ through the use of the traditional teleportation protocol (q.v. Example 6.50 in Section 6.3.1),

but where the classical communication channel required for teleportation is replaced by the ε -approximation to the channel $\Delta^{\otimes m}$ emulated by $\Phi^{\otimes n}$. It holds that Ψ is an ε -approximation to the identity channel $\mathbb{1}_{L(\mathcal{Z})}^{\otimes k}$.

One therefore has that, for all $\varepsilon > 0$, for all but finitely many positive integers n , and for

$$k = \left\lfloor \frac{\lfloor \alpha n \rfloor}{2} \right\rfloor = \left\lfloor \frac{\alpha n}{2} \right\rfloor, \quad (8.304)$$

the channel $\Phi^{\otimes n}$ emulates an ε -approximation to the identity channel $\mathbb{1}_{L(\mathcal{Z})}^{\otimes k}$ through the assistance of entanglement. It is therefore the case that $\alpha/2$ is an achievable rate for entanglement-assisted quantum communication through Φ , as required.

Now assume α is an achievable rate for entanglement-assisted quantum communication through Φ . It will be proved that 2α is an achievable rate for entanglement-assisted classical communication through Φ . This statement is trivial in the case $\alpha = 0$, so it will be assumed that $\alpha > 0$. The proof is essentially the same as the reverse direction just considered, with dense coding replacing teleportation.

Suppose that n and $m = \lfloor \alpha n \rfloor$ are positive integers and $\varepsilon > 0$ is a positive real number such that $\Phi^{\otimes n}$ emulates an ε -approximation to $\mathbb{1}_{L(\mathcal{Z})}^{\otimes m}$. Using the maximally entangled state

$$\tau = 2^{-m} \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m}) \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m})^*, \quad (8.305)$$

tensored with the state ξ used for the emulation of $\mathbb{1}_{L(\mathcal{Z})}^{\otimes m}$ by $\Phi^{\otimes n}$, one may define a new channel $\Psi \in C(\mathcal{Z}^{\otimes 2m})$ through the traditional dense coding protocol (q.v. Example 6.55 in Section 6.3.1), where the quantum channel required for dense coding is replaced by the ε -approximation to the channel $\mathbb{1}_{L(\mathcal{Z})}^{\otimes m}$ emulated by $\Phi^{\otimes n}$. It holds that Ψ is an ε -approximation to $\Delta^{\otimes 2m}$.

It therefore holds that, for all $\varepsilon > 0$, for all but finitely many values of n , and for $m = \lfloor \alpha n \rfloor$, that $\Phi^{\otimes n}$ emulates an ε -approximation to the channel $\Delta^{\otimes 2m}$, which implies that 2α is an achievable rate for entanglement-assisted classical communication through Φ . The inequality

$$C_E(\Phi) \geq 2Q_E(\Phi) \quad (8.306)$$

is obtained when one takes the supremum over all achievable rates α for entanglement-assisted quantum communication through Φ .

The equality (8.301) therefore holds, which completes the proof. \square

8.2.2 The quantum capacity theorem

The purpose of the present subsection is to state and prove the quantum capacity theorem, which yields an expression for the quantum capacity of a given channel. Similar to the Holevo–Schumacher–Westmoreland theorem (Theorem 8.27), the expression that is obtained from the quantum capacity theorem includes a regularization over an increasing number of uses of a given channel.

The subsections that follow include statements and proofs of lemmas that will be used to prove the quantum capacity theorem, as well as the statement and proof of the theorem itself.

A decoupling lemma

The first of several lemmas that will be used to prove the quantum capacity theorem concerns a phenomenon known as *decoupling*. Informally speaking, this is the phenomenon whereby the action of a sufficiently noisy channel on a randomly chosen subspace of its input space can be expected not only to destroy entanglement with a secondary system, but to destroy classical correlations as well. The lemma that follows proves a fact along these lines that is specialized to the task at hand.

Lemma 8.49 *Let \mathcal{X} , \mathcal{Y} , \mathcal{W} , and \mathcal{Z} be complex Euclidean spaces such that $\dim(\mathcal{Z}) \leq \dim(\mathcal{X}) \leq \dim(\mathcal{Y} \otimes \mathcal{W})$, and let $A \in U(\mathcal{X}, \mathcal{Y} \otimes \mathcal{W})$ and $V \in U(\mathcal{Z}, \mathcal{X})$ be isometries. Define a state $\xi \in D(\mathcal{W} \otimes \mathcal{X})$ as*

$$\xi = \frac{1}{n} \text{Tr}_{\mathcal{Y}}(\text{vec}(A) \text{vec}(A)^*), \quad (8.307)$$

and for each unitary operator $U \in U(\mathcal{X})$ define a state $\rho_U \in D(\mathcal{W} \otimes \mathcal{Z})$ as

$$\rho_U = \frac{1}{m} \text{Tr}_{\mathcal{Y}}(\text{vec}(AUV) \text{vec}(AUV)^*), \quad (8.308)$$

where $n = \dim(\mathcal{X})$ and $m = \dim(\mathcal{Z})$. It holds that

$$\int \|\rho_U - \text{Tr}_{\mathcal{Z}}(\rho_U) \otimes \omega\|_2^2 d\eta(U) \leq \text{Tr}(\xi^2), \quad (8.309)$$

for $\omega = \mathbb{1}_{\mathcal{Z}}/m$ and η denoting the Haar measure on $U(\mathcal{X})$.

Proof Observe first that

$$\|\rho_U - \text{Tr}_{\mathcal{Z}}(\rho_U) \otimes \omega\|_2^2 = \text{Tr}(\rho_U^2) - \frac{1}{m} \text{Tr}((\text{Tr}_{\mathcal{Z}}(\rho_U))^2). \quad (8.310)$$

The lemma requires a bound on the integral of the expression represented by (8.310) over all U , and toward this goal the two terms on the right-hand side of that equation will be integrated separately.

To integrate the first term on the right-hand side of (8.310), let Γ be the alphabet for which $\mathcal{Y} = \mathbb{C}^\Gamma$, define $B_a = (e_a^* \otimes \mathbb{1}_{\mathcal{W}})A$ for each $a \in \Gamma$, and observe that

$$\rho_U = \frac{1}{m} \sum_{a \in \Gamma} \text{vec}(B_a UV) \text{vec}(B_a UV)^*. \quad (8.311)$$

It therefore holds that

$$\begin{aligned} \text{Tr}(\rho_U^2) &= \frac{1}{m^2} \sum_{a,b \in \Gamma} |\text{Tr}(V^* U^* B_a^* B_b UV)|^2 \\ &= \frac{1}{m^2} \sum_{a,b \in \Gamma} \text{Tr}(V^* U^* B_a^* B_b UV \otimes V^* U^* B_b^* B_a UV) \\ &= \left\langle UVV^* U^* \otimes UVV^* U^*, \frac{1}{m^2} \sum_{a,b \in \Gamma} B_a^* B_b \otimes B_b^* B_a \right\rangle. \end{aligned} \quad (8.312)$$

Integrating over all $U \in \mathcal{U}(\mathcal{X})$ yields

$$\int \text{Tr}(\rho_U^2) d\eta(U) = \left\langle \Xi(VV^* \otimes VV^*), \frac{1}{m^2} \sum_{a,b \in \Gamma} B_a^* B_b \otimes B_b^* B_a \right\rangle, \quad (8.313)$$

for $\Xi \in \mathcal{C}(\mathcal{X} \otimes \mathcal{X})$ denoting the Werner twirling channel (q.v. Example 7.25 in the previous chapter). Making use of the expression

$$\Xi(X) = \frac{2}{n(n+1)} \langle \Pi_{\mathcal{X} \otimes \mathcal{X}}, X \rangle \Pi_{\mathcal{X} \otimes \mathcal{X}} + \frac{2}{n(n-1)} \langle \Pi_{\mathcal{X} \otimes \mathcal{X}}, X \rangle \Pi_{\mathcal{X} \otimes \mathcal{X}}, \quad (8.314)$$

which holds for every $X \in \mathcal{L}(\mathcal{X} \otimes \mathcal{X})$, and observing the equations

$$\langle \Pi_{\mathcal{X} \otimes \mathcal{X}}, VV^* \otimes VV^* \rangle = \frac{m(m+1)}{2}, \quad (8.315)$$

$$\langle \Pi_{\mathcal{X} \otimes \mathcal{X}}, VV^* \otimes VV^* \rangle = \frac{m(m-1)}{2}, \quad (8.316)$$

it follows that

$$\begin{aligned} &\int \text{Tr}(\rho_U^2) d\eta(U) \\ &= \frac{1}{nm} \left\langle \frac{m+1}{n+1} \Pi_{\mathcal{X} \otimes \mathcal{X}} + \frac{m-1}{n-1} \Pi_{\mathcal{X} \otimes \mathcal{X}}, \sum_{a,b \in \Gamma} B_a^* B_b \otimes B_b^* B_a \right\rangle. \end{aligned} \quad (8.317)$$

A similar methodology can be used to integrate the second term on the right-hand side of (8.310). In particular, one has

$$\text{Tr}_{\mathcal{Z}}(\rho_U) = \frac{1}{m} \sum_{a \in \Gamma} B_a UVV^* U^* B_a^*, \quad (8.318)$$

and therefore

$$\begin{aligned} &\text{Tr}\left(\left(\text{Tr}_{\mathcal{Z}}(\rho_U)\right)^2\right) \\ &= \frac{1}{m^2} \sum_{a,b \in \Gamma} \text{Tr}(V^* U^* B_a^* B_b UVV^* U^* B_b^* B_a UV) \\ &= \left\langle W_{\mathcal{Z}}, \frac{1}{m^2} \sum_{a,b \in \Gamma} V^* U^* B_a^* B_b UV \otimes V^* U^* B_b^* B_a UV \right\rangle \\ &= \left\langle (UV \otimes UV) W_{\mathcal{Z}} (UV \otimes UV)^*, \frac{1}{m^2} \sum_{a,b \in \Gamma} B_a^* B_b \otimes B_b^* B_a \right\rangle, \end{aligned} \quad (8.319)$$

where $W_{\mathcal{Z}} \in \mathcal{U}(\mathcal{Z} \otimes \mathcal{Z})$ denotes the swap operator on $\mathcal{Z} \otimes \mathcal{Z}$, and the second equality has used the identity $\langle W_{\mathcal{Z}}, X \otimes Y \rangle = \text{Tr}(XY)$. Integrating over all $U \in \mathcal{U}(\mathcal{X})$ yields

$$\begin{aligned} &\int \text{Tr}\left(\left(\text{Tr}_{\mathcal{Z}}(\rho_U)\right)^2\right) d\eta(U) \\ &= \left\langle \Xi((V \otimes V) W_{\mathcal{Z}} (V \otimes V)^*), \frac{1}{m^2} \sum_{a,b \in \Gamma} B_a^* B_b \otimes B_b^* B_a \right\rangle. \end{aligned} \quad (8.320)$$

By making use of the equations

$$\langle \Pi_{\mathcal{X} \otimes \mathcal{X}}, (V \otimes V) W_{\mathcal{Z}} (V \otimes V)^* \rangle = \frac{m(m+1)}{2}, \quad (8.321)$$

$$\langle \Pi_{\mathcal{X} \otimes \mathcal{X}}, (V \otimes V) W_{\mathcal{Z}} (V \otimes V)^* \rangle = -\frac{m(m-1)}{2},$$

and performing a similar calculation to the one above, one finds that

$$\begin{aligned} &\int \text{Tr}\left(\left(\text{Tr}_{\mathcal{Z}}(\rho_U)\right)^2\right) d\eta(U) \\ &= \frac{1}{nm} \left\langle \frac{m+1}{n+1} \Pi_{\mathcal{X} \otimes \mathcal{X}} - \frac{m-1}{n-1} \Pi_{\mathcal{X} \otimes \mathcal{X}}, \sum_{a,b \in \Gamma} B_a^* B_b \otimes B_b^* B_a \right\rangle. \end{aligned} \quad (8.322)$$

Combining (8.310), (8.317), and (8.322), together with some algebra, it follows that

$$\begin{aligned} &\int \|\rho_U - \text{Tr}_{\mathcal{Z}}(\rho_U) \otimes \omega\|_2^2 d\eta(U) \\ &= \frac{m^2 - 1}{m^2(n^2 - 1)} \left\langle \mathbb{1}_{\mathcal{X}} \otimes \mathbb{1}_{\mathcal{X}} - \frac{1}{n} W_{\mathcal{X}}, \sum_{a,b \in \Gamma} B_a^* B_b \otimes B_b^* B_a \right\rangle, \end{aligned} \quad (8.323)$$

where $W_{\mathcal{X}}$ denotes the swap operator on $\mathcal{X} \otimes \mathcal{X}$. By similar calculations to

(8.312) and (8.319) above, but replacing U and V by $\mathbf{1}_{\mathcal{X}}$, it may be verified that

$$\mathrm{Tr}(\xi^2) = \frac{1}{n^2} \mathrm{Tr} \left(\sum_{a,b \in \Gamma} B_a^* B_b \otimes B_b^* B_a \right) \quad (8.324)$$

and

$$\mathrm{Tr} \left((\mathrm{Tr}_{\mathcal{X}}(\xi))^2 \right) = \frac{1}{n^2} \left\langle W_{\mathcal{X}}, \sum_{a,b \in \Gamma} B_a^* B_b \otimes B_b^* B_a \right\rangle. \quad (8.325)$$

Consequently,

$$\begin{aligned} & \int \|\rho_U - \mathrm{Tr}_{\mathcal{Z}}(\rho_U) \otimes \omega\|_2^2 d\eta(U) \\ &= \frac{1-m^{-2}}{1-n^{-2}} \left(\mathrm{Tr}(\xi^2) - \frac{1}{n} \mathrm{Tr} \left((\mathrm{Tr}_{\mathcal{X}}(\xi))^2 \right) \right) \leq \mathrm{Tr}(\xi^2), \end{aligned} \quad (8.326)$$

as required. \square

A lower-bound on entanglement generation decoding fidelity

The next lemma is used, within the proof of the quantum capacity theorem, to infer the existence of a decoding channel for the task of entanglement generation. This inference is based on a calculation involving a Stinespring representation of the channel through which entanglement generation is to be considered.

Lemma 8.50 *Let \mathcal{X} , \mathcal{Y} , \mathcal{W} , and \mathcal{Z} be complex Euclidean spaces such that $\dim(\mathcal{Z}) \leq \dim(\mathcal{X}) \leq \dim(\mathcal{Y} \otimes \mathcal{W})$, and let $A \in \mathrm{U}(\mathcal{X}, \mathcal{Y} \otimes \mathcal{W})$ and $W \in \mathrm{U}(\mathcal{Z}, \mathcal{X})$ be isometries. Define a channel $\Phi \in \mathrm{C}(\mathcal{X}, \mathcal{Y})$ as*

$$\Phi(X) = \mathrm{Tr}_{\mathcal{W}}(AXA^*) \quad (8.327)$$

for all $X \in \mathrm{L}(\mathcal{X})$, and define a state $\rho \in \mathrm{D}(\mathcal{W} \otimes \mathcal{Z})$ as

$$\rho = \frac{1}{m} \mathrm{Tr}_{\mathcal{Y}}(\mathrm{vec}(AW) \mathrm{vec}(AW)^*), \quad (8.328)$$

where $m = \dim(\mathcal{Z})$. There exists a channel $\Xi \in \mathrm{C}(\mathcal{Y}, \mathcal{Z})$ such that

$$\begin{aligned} & \mathrm{F} \left(\frac{1}{m} \mathrm{vec}(\mathbf{1}_{\mathcal{Z}}) \mathrm{vec}(\mathbf{1}_{\mathcal{Z}})^*, \frac{1}{m} (\Xi \Phi \otimes \mathbf{1}_{\mathrm{L}(\mathcal{Z})})(\mathrm{vec}(W) \mathrm{vec}(W)^*) \right) \\ & \geq \mathrm{F}(\rho, \mathrm{Tr}_{\mathcal{Z}}(\rho) \otimes \omega), \end{aligned} \quad (8.329)$$

where $\omega = \mathbf{1}_{\mathcal{Z}}/m$.

Proof Let \mathcal{V} be a complex Euclidean space of sufficiently large dimension that the inequalities $\dim(\mathcal{V}) \geq \dim(\mathcal{W})$ and $\dim(\mathcal{V} \otimes \mathcal{Z}) \geq \dim(\mathcal{Y})$ hold, and let $B \in \mathrm{L}(\mathcal{W}, \mathcal{V})$ be an operator such that $\mathrm{Tr}_{\mathcal{V}}(\mathrm{vec}(B) \mathrm{vec}(B)^*) = \mathrm{Tr}_{\mathcal{Z}}(\rho)$. For the vector

$$u = \frac{1}{\sqrt{m}} \mathrm{vec}(B \otimes \mathbf{1}_{\mathcal{Z}}) \in (\mathcal{V} \otimes \mathcal{Z}) \otimes (\mathcal{W} \otimes \mathcal{Z}), \quad (8.330)$$

one has that $\mathrm{Tr}_{\mathcal{V} \otimes \mathcal{Z}}(uu^*) = \mathrm{Tr}_{\mathcal{Z}}(\rho) \otimes \omega$. It is evident that the vector

$$v = \frac{1}{\sqrt{m}} \mathrm{vec}(AW) \in \mathcal{Y} \otimes \mathcal{W} \otimes \mathcal{Z} \quad (8.331)$$

satisfies $\mathrm{Tr}_{\mathcal{Y}}(vv^*) = \rho$, so it follows by Uhlmann's theorem (Theorem 3.22) that there exists an isometry $V \in \mathrm{U}(\mathcal{Y}, \mathcal{V} \otimes \mathcal{Z})$ such that

$$\mathrm{F}(\rho, \mathrm{Tr}_{\mathcal{Z}}(\rho) \otimes \omega) = \mathrm{F}(uu^*, (V \otimes \mathbf{1}_{\mathcal{W} \otimes \mathcal{Z}})vv^*(V \otimes \mathbf{1}_{\mathcal{W} \otimes \mathcal{Z}})^*). \quad (8.332)$$

Define a channel $\Xi \in \mathrm{C}(\mathcal{Y}, \mathcal{Z})$ as

$$\Xi(Y) = \mathrm{Tr}_{\mathcal{V}}(VYV^*) \quad (8.333)$$

for every $Y \in \mathrm{L}(\mathcal{Y})$. It holds that

$$\mathrm{Tr}_{\mathcal{V}}(\mathrm{Tr}_{\mathcal{W}}(uu^*)) = \frac{1}{m} \mathrm{vec}(\mathbf{1}_{\mathcal{Z}}) \mathrm{vec}(\mathbf{1}_{\mathcal{Z}})^* \quad (8.334)$$

and

$$\begin{aligned} & \mathrm{Tr}_{\mathcal{V}}(\mathrm{Tr}_{\mathcal{W}}((V \otimes \mathbf{1}_{\mathcal{W} \otimes \mathcal{Z}})vv^*(V \otimes \mathbf{1}_{\mathcal{W} \otimes \mathcal{Z}})^*)) \\ &= \frac{1}{m} (\Xi \Phi \otimes \mathbf{1}_{\mathrm{L}(\mathcal{Z})})(\mathrm{vec}(W) \mathrm{vec}(W)^*), \end{aligned} \quad (8.335)$$

and therefore

$$\begin{aligned} & \mathrm{F}(uu^*, (V \otimes \mathbf{1}_{\mathcal{W} \otimes \mathcal{Z}})vv^*(V \otimes \mathbf{1}_{\mathcal{W} \otimes \mathcal{Z}})^*) \\ & \leq \mathrm{F} \left(\frac{1}{m} \mathrm{vec}(\mathbf{1}_{\mathcal{Z}}) \mathrm{vec}(\mathbf{1}_{\mathcal{Z}})^*, \frac{1}{m} (\Xi \Phi \otimes \mathbf{1}_{\mathrm{L}(\mathcal{Z})})(\mathrm{vec}(W) \mathrm{vec}(W)^*) \right) \end{aligned} \quad (8.336)$$

by the monotonicity of the fidelity under partial tracing (which is a special case of Theorem 3.27). The channel Ξ therefore satisfies the requirement of the lemma. \square

Two additional lemmas needed for the quantum capacity theorem

The two lemmas that follow represent technical facts that will be utilized in the proof of the quantum capacity theorem. The first lemma concerns the approximation of one isometry by another isometry that meets certain spectral requirements, and the second lemma is a general fact regarding Haar measure.

Lemma 8.51 *Let \mathcal{X} , \mathcal{Y} , and \mathcal{W} be complex Euclidean spaces such that $\dim(\mathcal{X}) \leq \dim(\mathcal{Y} \otimes \mathcal{W})$, let $A \in \mathbf{U}(\mathcal{X}, \mathcal{Y} \otimes \mathcal{W})$ be an isometry, let $\Lambda \in \mathbf{Proj}(\mathcal{Y})$ and $\Pi \in \mathbf{Proj}(\mathcal{W})$ be projection operators, and let $\varepsilon \in (0, 1/4)$ be a positive real number. Also let $n = \dim(\mathcal{X})$, and assume that the constraints*

$$\langle \Lambda \otimes \Pi, AA^* \rangle \geq (1 - \varepsilon)n \quad (8.337)$$

and

$$2 \operatorname{rank}(\Pi) \leq \dim(\mathcal{W}) \quad (8.338)$$

are satisfied. There exists an isometry $B \in \mathbf{U}(\mathcal{X}, \mathcal{Y} \otimes \mathcal{W})$ such that

1. $\|A - B\|_2 < 3\varepsilon^{1/4}\sqrt{n}$,
2. $\operatorname{Tr}_{\mathcal{W}}(BB^*) \leq 4\Lambda \operatorname{Tr}_{\mathcal{W}}(AA^*)\Lambda$, and
3. $\operatorname{rank}(\operatorname{Tr}_{\mathcal{Y}}(BB^*)) \leq 2 \operatorname{rank}(\Pi)$.

Proof By means of the singular value theorem, one may write

$$(\Lambda \otimes \Pi)A = \sum_{k=1}^n s_k u_k x_k^* \quad (8.339)$$

for an orthonormal basis $\{x_1, \dots, x_n\}$ of \mathcal{X} , an orthonormal set $\{u_1, \dots, u_n\}$ of vectors in $\mathcal{Y} \otimes \mathcal{W}$, and a collection $\{s_1, \dots, s_n\} \subset [0, 1]$ of nonnegative real numbers. It holds that

$$\sum_{k=1}^n s_k^2 = \langle \Lambda \otimes \Pi, AA^* \rangle \geq (1 - \varepsilon)n. \quad (8.340)$$

Define $\Gamma \subseteq \{1, \dots, n\}$ as

$$\Gamma = \{k \in \{1, \dots, n\} : s_k^2 \geq 1 - \sqrt{\varepsilon}\}, \quad (8.341)$$

and observe the inequality

$$\sum_{k=1}^n s_k^2 \leq |\Gamma| + (n - |\Gamma|)(1 - \sqrt{\varepsilon}). \quad (8.342)$$

From (8.340) and (8.342) it follows that

$$|\Gamma| \geq (1 - \sqrt{\varepsilon})n > \frac{n}{2}. \quad (8.343)$$

There must therefore exist an injective function $f : \{1, \dots, n\} \setminus \Gamma \rightarrow \Gamma$; this function may be chosen arbitrarily, but will be fixed for the remainder of the proof.

Next, let $W \in \mathbf{U}(\mathcal{W})$ be any unitary operator satisfying $\Pi W \Pi = 0$. The assumption that $2 \operatorname{rank}(\Pi) \leq \dim(\mathcal{W})$ guarantees the existence of such an operator W . As for the function f , the unitary operator W may be chosen arbitrarily, subject to the condition $\Pi W \Pi = 0$, but is understood to be fixed for the remainder of the proof.

Finally, define an isometry $B \in \mathbf{U}(\mathcal{X}, \mathcal{Y} \otimes \mathcal{W})$ as follows:

$$B = \sum_{k \in \Gamma} u_k x_k^* + \sum_{k \in \{1, \dots, n\} \setminus \Gamma} (\mathbf{1}_{\mathcal{Y}} \otimes W) u_{f(k)} x_k^*. \quad (8.344)$$

It remains to prove that B has the properties required by the statement of the lemma.

First, it will be verified that B is indeed an isometry. The set $\{u_k : k \in \Gamma\}$ is evidently orthonormal, as is the set

$$\{(\mathbf{1}_{\mathcal{Y}} \otimes W) u_{f(k)} : k \in \{1, \dots, n\} \setminus \Gamma\}. \quad (8.345)$$

For every choice of $k \in \{1, \dots, n\}$ one has

$$s_k u_k \in \operatorname{im}((\Lambda \otimes \Pi)A) \subseteq \operatorname{im}(\mathbf{1}_{\mathcal{Y}} \otimes \Pi), \quad (8.346)$$

and therefore $s_k u_k = s_k (\mathbf{1}_{\mathcal{Y}} \otimes \Pi) u_k$. It follows that

$$\begin{aligned} s_j s_k \langle u_j, (\mathbf{1}_{\mathcal{Y}} \otimes W) u_k \rangle &= s_j s_k \langle (\mathbf{1}_{\mathcal{Y}} \otimes \Pi) u_j, (\mathbf{1}_{\mathcal{Y}} \otimes \Pi W) u_k \rangle \\ &= s_j s_k \langle u_j, (\mathbf{1}_{\mathcal{Y}} \otimes \Pi W \Pi) u_k \rangle = 0 \end{aligned} \quad (8.347)$$

for every choice of $j, k \in \{1, \dots, n\}$, by virtue of the fact that $\Pi W \Pi = 0$. For $j, k \in \Gamma$, it must hold that $s_j s_k > 0$, and therefore $u_j \perp (\mathbf{1}_{\mathcal{Y}} \otimes W) u_k$. This implies that the set

$$\{u_k : k \in \Gamma\} \cup \{(\mathbf{1}_{\mathcal{Y}} \otimes W) u_{f(k)} : k \in \{1, \dots, n\} \setminus \Gamma\} \quad (8.348)$$

is orthonormal, and therefore B is an isometry.

Next, observe that

$$\|A - B\|_2 \leq \|A - (\Lambda \otimes \Pi)A\|_2 + \|(\Lambda \otimes \Pi)A - B\|_2. \quad (8.349)$$

The first term in this expression is bounded as

$$\|A - (\Lambda \otimes \Pi)A\|_2 = \sqrt{\langle \mathbf{1} - \Lambda \otimes \Pi, AA^* \rangle} \leq \sqrt{\varepsilon n}. \quad (8.350)$$

For the second term, it holds that

$$\begin{aligned} \|(\Lambda \otimes \Pi)A - B\|_2^2 &= \sum_{k \in \Gamma} (s_k - 1)^2 + \sum_{k \in \{1, \dots, n\} \setminus \Gamma} (s_k^2 + 1) \\ &= n + \sum_{k=1}^n s_k^2 - 2 \sum_{k \in \Gamma} s_k \leq 2n - 2|\Gamma|(1 - \sqrt{\varepsilon})^{\frac{1}{2}}. \end{aligned} \quad (8.351)$$

To obtain the first equality in the previous equation, it is helpful to observe that

$$s_k u_k \perp (\mathbb{1}_{\mathcal{Y}} \otimes W) u_{f(k)} \quad (8.352)$$

for $k \in \{1, \dots, n\} \setminus \Gamma$, which makes use of the equation (8.347) along with the inclusion $f(k) \in \Gamma$. By the inequality (8.343) it therefore holds that

$$\|(\Lambda \otimes \Pi)A - B\|_2^2 \leq 2n - 2(1 - \sqrt{\varepsilon})^{\frac{3}{2}}n < 3n\sqrt{\varepsilon}, \quad (8.353)$$

from which it follows that

$$\|A - B\|_2 < 3\varepsilon^{1/4}\sqrt{n}. \quad (8.354)$$

The first requirement on B listed in the statement of the lemma is therefore fulfilled.

The second requirement on B may be verified as follows:

$$\begin{aligned} \operatorname{Tr}_{\mathcal{W}}(BB^*) &\leq 2 \sum_{k \in \Gamma} \operatorname{Tr}_{\mathcal{W}}(u_k u_k^*) \\ &\leq \frac{2}{1 - \sqrt{\varepsilon}} \operatorname{Tr}_{\mathcal{W}}((\Lambda \otimes \Pi)AA^*(\Lambda \otimes \Pi)) \leq 4\Lambda \operatorname{Tr}_{\mathcal{W}}(AA^*)\Lambda. \end{aligned} \quad (8.355)$$

Finally, to verify that the third requirement on B is satisfied, one may again use the observation that $(\mathbb{1} \otimes \Pi)u_k = u_k$, which implies that

$$\operatorname{im}(\operatorname{Tr}_{\mathcal{Y}}(u_k u_k^*)) \subseteq \operatorname{im}(\Pi), \quad (8.356)$$

for each $k \in \Gamma$. As

$$\operatorname{Tr}_{\mathcal{Y}}(BB^*) = \sum_{k \in \Gamma} \operatorname{Tr}_{\mathcal{Y}}(u_k u_k^*) + \sum_{k \in \{1, \dots, n\} \setminus \Gamma} W(\operatorname{Tr}_{\mathcal{Y}}(u_{f(k)} u_{f(k)}^*))W^*, \quad (8.357)$$

it follows that

$$\operatorname{im}(\operatorname{Tr}_{\mathcal{Y}}(BB^*)) \subseteq \operatorname{im}(\Pi) + \operatorname{im}(W\Pi) \quad (8.358)$$

and therefore

$$\operatorname{rank}(\operatorname{Tr}_{\mathcal{Y}}(BB^*)) \leq 2\operatorname{rank}(\Pi), \quad (8.359)$$

as required. \square

Lemma 8.52 *Let \mathcal{X} , \mathcal{W} , and \mathcal{Z} be complex Euclidean spaces such that $\dim(\mathcal{Z}) \leq \dim(\mathcal{X})$, let $V \in \mathcal{U}(\mathcal{Z}, \mathcal{X})$ be an isometry, and let $Z \in \mathcal{L}(\mathcal{W} \otimes \mathcal{X})$ be an operator. It holds that*

$$\int \|(\mathbb{1}_{\mathcal{W}} \otimes V^*U^*)Z(\mathbb{1}_{\mathcal{W}} \otimes UV)\|_1 d\eta(U) \leq \frac{m}{n} \|Z\|_1 \quad (8.360)$$

where $m = \dim(\mathcal{Z})$, $n = \dim(\mathcal{X})$, and η denotes the Haar measure on $\mathcal{U}(\mathcal{X})$.

Proof Let $\{W_1, \dots, W_{n^2}\} \subset \mathcal{U}(\mathcal{X})$ be an orthogonal collection of unitary operators. (The discrete Weyl operators, defined in Section 4.1.2, provide an explicit choice for such a collection.) It therefore holds that the completely depolarizing channel $\Omega \in \mathcal{C}(\mathcal{X})$ may be expressed as

$$\Omega(X) = \frac{1}{n^2} \sum_{k=1}^{n^2} W_k X W_k^* \quad (8.361)$$

for all $X \in \mathcal{L}(\mathcal{X})$. Define $\mathcal{Y} = \mathbb{C}^{n^2}$, and define a channel $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Z} \otimes \mathcal{Y})$ as

$$\Phi(X) = \frac{1}{nm} \sum_{k=1}^{n^2} V^* W_k^* X W_k V \otimes E_{k,k} \quad (8.362)$$

for every $X \in \mathcal{L}(\mathcal{X})$. The fact that Φ is a channel follows from Corollary 2.27 together with the calculation

$$\frac{1}{nm} \sum_{k=1}^{n^2} W_k V V^* W_k^* = \frac{n}{m} \Omega(VV^*) = \mathbb{1}_{\mathcal{X}}. \quad (8.363)$$

Next, by the right unitary invariance of the Haar measure, it holds that

$$\begin{aligned} &\int \|(\mathbb{1}_{\mathcal{W}} \otimes V^*U^*)Z(\mathbb{1}_{\mathcal{W}} \otimes UV)\|_1 d\eta(U) \\ &= \int \|(\mathbb{1}_{\mathcal{W}} \otimes V^*W_k^*U^*)Z(\mathbb{1}_{\mathcal{W}} \otimes UW_k V)\|_1 d\eta(U) \end{aligned} \quad (8.364)$$

for every choice of $k \in \{1, \dots, n^2\}$, and therefore

$$\begin{aligned} &\int \|(\mathbb{1}_{\mathcal{W}} \otimes UV)^*Z(\mathbb{1}_{\mathcal{W}} \otimes UV)\|_1 d\eta(U) \\ &= \frac{1}{n^2} \sum_{k=1}^{n^2} \int \|(\mathbb{1}_{\mathcal{W}} \otimes UW_k V)^*Z(\mathbb{1}_{\mathcal{W}} \otimes UW_k V)\|_1 d\eta(U) \\ &= \frac{1}{n^2} \int \left\| \sum_{k=1}^{n^2} (\mathbb{1}_{\mathcal{W}} \otimes UW_k V)^*Z(\mathbb{1}_{\mathcal{W}} \otimes UW_k V) \otimes E_{k,k} \right\|_1 d\eta(U) \\ &= \frac{m}{n} \int \|(\mathbb{1}_{\mathcal{L}(\mathcal{W})} \otimes \Phi)((\mathbb{1}_{\mathcal{W}} \otimes U^*)Z(\mathbb{1}_{\mathcal{W}} \otimes U))\|_1 d\eta(U). \end{aligned} \quad (8.365)$$

As the trace norm is non-increasing under the action of channels, as well as unitary invariant, it follows that

$$\begin{aligned} &\frac{m}{n} \int \|(\mathbb{1}_{\mathcal{L}(\mathcal{W})} \otimes \Phi)((\mathbb{1}_{\mathcal{W}} \otimes U^*)Z(\mathbb{1}_{\mathcal{W}} \otimes U))\|_1 d\eta(U) \\ &\leq \frac{m}{n} \int \|(\mathbb{1}_{\mathcal{W}} \otimes U^*)Z(\mathbb{1}_{\mathcal{W}} \otimes U)\|_1 d\eta(U) = \frac{m}{n} \|Z\|_1, \end{aligned} \quad (8.366)$$

which completes the proof. \square

The quantum capacity theorem

As the following theorem establishes, the entanglement-generation capacity of a given channel is always at least as large as the coherent information of the completely mixed state through that channel. This fact, which will be generalized to arbitrary states in place of the completely mixed state in a corollary to the theorem, lies at the heart of the proof of the quantum capacity theorem.

Theorem 8.53 *Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} . The entanglement generation capacity of Φ is at least the coherent information of the completely mixed state $\omega \in D(\mathcal{X})$ through Φ :*

$$I_c(\omega; \Phi) \leq Q_{\text{EG}}(\Phi). \quad (8.367)$$

Proof Let \mathcal{W} be a complex Euclidean space such that

$$\dim(\mathcal{W}) = 2 \dim(\mathcal{X} \otimes \mathcal{Y}), \quad (8.368)$$

and let $A \in U(\mathcal{X}, \mathcal{Y} \otimes \mathcal{W})$ be an isometry for which

$$\Phi(X) = \text{Tr}_{\mathcal{W}}(AXA^*) \quad (8.369)$$

for all $X \in L(\mathcal{X})$. The somewhat unusual factor of 2 on the right-hand side of (8.368) will guarantee that the assumptions required by Lemma 8.51 are met, as is mentioned later in the proof. Define a channel $\Psi \in C(\mathcal{X}, \mathcal{W})$ as

$$\Psi(X) = \text{Tr}_{\mathcal{Y}}(AXA^*) \quad (8.370)$$

for all $X \in L(\mathcal{X})$, so that Ψ is complementary to Φ . It therefore holds that

$$I_c(\omega; \Phi) = H(\Phi(\omega)) - H(\Psi(\omega)). \quad (8.371)$$

The theorem is vacuous in the case that $I_c(\omega; \Phi) \leq 0$, so hereafter it will be assumed that $I_c(\omega; \Phi)$ is positive. To prove the theorem, it suffices to demonstrate that every positive real number smaller than $I_c(\omega; \Phi)$ is an achievable rate for entanglement generation through Φ . Toward this goal, assume that an arbitrary positive real number α satisfying $\alpha < I_c(\omega; \Phi)$ has been fixed, and that $\varepsilon > 0$ is a positive real number chosen to be sufficiently small so that the inequality

$$\alpha < I_c(\omega; \Phi) - 2\varepsilon(H(\Phi(\omega)) + H(\Psi(\omega))) \quad (8.372)$$

is satisfied. The remainder of the proof is devoted to proving that α is an achievable rate for entanglement generation through Φ .

Consider an arbitrary positive integer $n \geq 1/\alpha$, and let $m = \lfloor \alpha n \rfloor$. Also let $\Gamma = \{0, 1\}$ denote the binary alphabet, and let $\mathcal{Z} = \mathbb{C}^\Gamma$. The task in which a state having high fidelity with the maximally entangled state

$$2^{-m} \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m}) \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m})^* \quad (8.373)$$

is established between a sender and receiver through the channel $\Phi^{\otimes n}$ is to be considered. Note that the quantity $I_c(\omega; \Phi)$ is at most $\log(\dim(\mathcal{X}))$, and therefore $\alpha < \log(\dim(\mathcal{X}))$, and this implies $\dim(\mathcal{Z}^{\otimes m}) \leq \dim(\mathcal{X}^{\otimes n})$. For any isometry $W \in U(\mathcal{Z}^{\otimes m}, \mathcal{X}^{\otimes n})$ and a channel $\Xi \in C(\mathcal{Y}^{\otimes n}, \mathcal{Z}^{\otimes m})$, the state

$$2^{-m} (\Xi \Phi^{\otimes n} \otimes \mathbb{1}_{L(\mathcal{Z})}^{\otimes m}) (\text{vec}(W) \text{vec}(W)^*) \quad (8.374)$$

may be established through the channel $\Phi^{\otimes n}$, so one may aim to prove that there exists a choice of Ξ and W for which the fidelity between the states (8.373) and (8.374) is high.

It is helpful at this point to let $A_n \in U(\mathcal{X}^{\otimes n}, \mathcal{Y}^{\otimes n} \otimes \mathcal{W}^{\otimes n})$ be the isometry defined by the equation

$$\begin{aligned} & \langle y_1 \otimes \cdots \otimes y_n \otimes w_1 \otimes \cdots \otimes w_n, A_n(x_1 \otimes \cdots \otimes x_n) \rangle \\ &= \langle y_1 \otimes w_1, Ax_1 \rangle \cdots \langle y_n \otimes w_n, Ax_n \rangle \end{aligned} \quad (8.375)$$

holding for every choice of vectors $x_1, \dots, x_n \in \mathcal{X}$, $y_1, \dots, y_n \in \mathcal{Y}$, and $w_1, \dots, w_n \in \mathcal{W}$. In effect, A_n is equivalent to $A^{\otimes n}$, except that the tensor factors in its output space have been permuted, so that the output space becomes $\mathcal{Y}^{\otimes n} \otimes \mathcal{W}^{\otimes n}$ rather than $(\mathcal{Y} \otimes \mathcal{W})^{\otimes n}$. It may be noted that

$$\Phi^{\otimes n}(X) = \text{Tr}_{\mathcal{W}^{\otimes n}}(A_n X A_n^*) \quad \text{and} \quad \Psi^{\otimes n}(X) = \text{Tr}_{\mathcal{Y}^{\otimes n}}(A_n X A_n^*) \quad (8.376)$$

for every $X \in L(\mathcal{X}^{\otimes n})$.

Now, under the assumption that the decoding channel $\Xi \in C(\mathcal{Y}^{\otimes n}, \mathcal{Z}^{\otimes m})$ has been selected optimally, Lemma 8.50 implies that the fidelity between the states (8.373) and (8.374) is lower-bounded by

$$F(\rho, \text{Tr}_{\mathcal{Z}^{\otimes m}}(\rho) \otimes \omega_{\mathcal{Z}}^{\otimes m}) \quad (8.377)$$

for $\rho \in D(\mathcal{W}^{\otimes n} \otimes \mathcal{Z}^{\otimes m})$ defined as

$$\rho = 2^{-m} \text{Tr}_{\mathcal{Y}^{\otimes n}}(\text{vec}(A_n W) \text{vec}(A_n W)^*) \quad (8.378)$$

and for $\omega_{\mathcal{Z}} \in D(\mathcal{Z})$ denoting the completely mixed state on \mathcal{Z} .

The probabilistic method will be employed to prove the existence of an isometry W for which the expression (8.377) is close to 1, provided that n is sufficiently large. In particular, one may fix $V \in U(\mathcal{Z}^{\otimes m}, \mathcal{X}^{\otimes n})$ to be an

arbitrary isometry, and let $W = UV$ for U chosen at random with respect to the Haar measure on $U(\mathcal{X}^{\otimes n})$. The analysis that follows demonstrates that, for an operator W chosen in this way, one expects the quantity (8.377) to be close to 1, for sufficiently large n , which proves the existence of a choice of W for which this is true.

Let $k = \dim(\mathcal{X})$ and define $\xi \in D(\mathcal{W}^{\otimes n} \otimes \mathcal{X}^{\otimes n})$ as

$$\xi = \frac{1}{k^n} \text{Tr}_{\mathcal{Y}^{\otimes n}}(\text{vec}(A_n) \text{vec}(A_n)^*). \quad (8.379)$$

Also define $\rho_U \in D(\mathcal{W}^{\otimes n} \otimes \mathcal{Z}^{\otimes m})$ as

$$\rho_U = \frac{1}{2^m} \text{Tr}_{\mathcal{Y}^{\otimes n}}(\text{vec}(A_n UV) \text{vec}(A_n UV)^*), \quad (8.380)$$

for each unitary operator $U \in U(\mathcal{X}^{\otimes n})$, and observe that

$$\rho_U = \frac{k^n}{2^m} (\mathbb{1}_{\mathcal{W}}^{\otimes n} \otimes V^\top U^\top) \xi (\mathbb{1}_{\mathcal{W}}^{\otimes n} \otimes V^\top U^\top)^*. \quad (8.381)$$

For the isometry $W = UV$, the fidelity between the states (8.373) and (8.374) is lower-bounded by

$$F(\rho_U, \text{Tr}_{\mathcal{Z}^{\otimes m}}(\rho_U) \otimes \omega_{\mathcal{Z}}^{\otimes m}), \quad (8.382)$$

for a suitable choice of the decoding channel Ξ .

Let $\Lambda_{n,\varepsilon} \in \text{Proj}(\mathcal{Y}^{\otimes n})$ and $\Pi_{n,\varepsilon} \in \text{Proj}(\mathcal{W}^{\otimes n})$ be the projection operators onto the ε -strongly typical subspaces of $\mathcal{Y}^{\otimes n}$ and $\mathcal{W}^{\otimes n}$, with respect to any fixed choice of spectral decompositions of $\Phi(\omega)$ and $\Psi(\omega)$, respectively. One may observe that because $\varepsilon > 0$ and $\text{rank}(\Psi(\omega)) \leq \dim(\mathcal{X} \otimes \mathcal{Y})$, it holds that

$$\text{rank}(\Pi_{n,\varepsilon}) \leq \frac{1}{2^n} \dim(\mathcal{W}^{\otimes n}) \leq \frac{1}{2} \dim(\mathcal{W}^{\otimes n}). \quad (8.383)$$

This is a very coarse bound that will nevertheless be required in order to utilize Lemma 8.51, and explains the factor of 2 in (8.368).

By Lemma 8.32, there must exist positive real numbers K and δ , both independent of n and ε , and both assumed to be fixed for the remainder of the proof, such that for

$$\zeta_{n,\varepsilon} = K \exp(-\delta n \varepsilon^2), \quad (8.384)$$

one has these inequalities:

$$\begin{aligned} \frac{1}{k^n} \langle \Lambda_{n,\varepsilon} \otimes \mathbb{1}_{\mathcal{W}}^{\otimes n} \otimes \mathbb{1}_{\mathcal{X}}^{\otimes n}, \text{vec}(A_n) \text{vec}(A_n)^* \rangle \\ &= \langle \Lambda_{n,\varepsilon}, (\Phi(\omega))^{\otimes n} \rangle \geq 1 - \frac{\zeta_{n,\varepsilon}}{2}, \\ \frac{1}{k^n} \langle \mathbb{1}_{\mathcal{Y}}^{\otimes n} \otimes \Pi_{n,\varepsilon} \otimes \mathbb{1}_{\mathcal{X}}^{\otimes n}, \text{vec}(A_n) \text{vec}(A_n)^* \rangle \\ &= \langle \Pi_{n,\varepsilon}, (\Psi(\omega))^{\otimes n} \rangle \geq 1 - \frac{\zeta_{n,\varepsilon}}{2}. \end{aligned} \quad (8.385)$$

It follows that

$$\frac{1}{k^n} \langle \Lambda_{n,\varepsilon} \otimes \Pi_{n,\varepsilon} \otimes \mathbb{1}_{\mathcal{X}}^{\otimes n}, \text{vec}(A_n) \text{vec}(A_n)^* \rangle \geq 1 - \zeta_{n,\varepsilon}, \quad (8.386)$$

which is equivalent to

$$\langle \Lambda_{n,\varepsilon} \otimes \Pi_{n,\varepsilon}, A_n A_n^* \rangle \geq (1 - \zeta_{n,\varepsilon}) k^n. \quad (8.387)$$

If n is sufficiently large so that $\zeta_{n,\varepsilon} < 1/4$, it follows by Lemma 8.51 that there exists an isometry $B_n \in U(\mathcal{X}^{\otimes n}, \mathcal{Y}^{\otimes n} \otimes \mathcal{W}^{\otimes n})$ satisfying these three conditions:

$$\begin{aligned} \|A_n - B_n\|_2 &\leq 3 \zeta_{n,\varepsilon}^{1/4} k^{n/2}, \\ \text{Tr}_{\mathcal{W}^{\otimes n}}(B_n B_n^*) &\leq 4 \Lambda_{n,\varepsilon} \text{Tr}_{\mathcal{W}^{\otimes n}}(A_n A_n^*) \Lambda_{n,\varepsilon}, \\ \text{rank}(\text{Tr}_{\mathcal{Y}^{\otimes n}}(B_n B_n^*)) &\leq 2 \text{rank}(\Pi_{n,\varepsilon}). \end{aligned} \quad (8.388)$$

By Proposition 8.33, the third condition implies that

$$\text{rank}(\text{Tr}_{\mathcal{Y}^{\otimes n}}(B_n B_n^*)) \leq 2^{n(1+\varepsilon) \text{H}(\Psi(\omega))+1}. \quad (8.389)$$

Using the second condition, together with Corollary 8.31 and the inequality $\text{Tr}(P^2) \leq \lambda_1(P) \text{Tr}(P)$, which holds for all $P \geq 0$, one obtains

$$\begin{aligned} \text{Tr} \left(\left(\frac{1}{k^n} \text{Tr}_{\mathcal{W}^{\otimes n}}(B_n B_n^*) \right)^2 \right) \\ &\leq \text{Tr} \left(\left(\frac{4}{k^n} \Lambda_{n,\varepsilon} \text{Tr}_{\mathcal{W}^{\otimes n}}(A_n A_n^*) \Lambda_{n,\varepsilon} \right)^2 \right) \\ &= 16 \text{Tr} \left((\Lambda_{n,\varepsilon} \Phi(\omega)^{\otimes n} \Lambda_{n,\varepsilon})^2 \right) \\ &\leq 2^{-n(1-\varepsilon) \text{H}(\Phi(\omega))+4}. \end{aligned} \quad (8.390)$$

Finally, define

$$\sigma = \frac{1}{k^n} \text{Tr}_{\mathcal{Y}^{\otimes n}}(\text{vec}(B_n) \text{vec}(B_n)^*), \quad (8.391)$$

and also define

$$\begin{aligned}\tau_U &= \frac{1}{2^m} \text{Tr}_{\mathcal{Y}^{\otimes n}}(\text{vec}(B_n UV) \text{vec}(B_n UV)^*) \\ &= \frac{k^n}{2^m} (\mathbf{1}_{\mathcal{W}}^{\otimes n} \otimes V^T U^T) \sigma(\mathbf{1}_{\mathcal{W}}^{\otimes n} \otimes V^T U^T)^*\end{aligned}\quad (8.392)$$

for each $U \in \text{U}(\mathcal{X}^{\otimes n})$. It holds that

$$\begin{aligned}\|\rho_U - \text{Tr}_{\mathcal{Z}^{\otimes m}}(\rho_U) \otimes \omega_{\mathcal{Z}}^{\otimes m}\|_1 &\leq \|\rho_U - \tau_U\|_1 + \|\tau_U - \text{Tr}_{\mathcal{Z}^{\otimes m}}(\tau_U) \otimes \omega_{\mathcal{Z}}^{\otimes m}\|_1 \\ &\quad + \left\| (\text{Tr}_{\mathcal{Z}^{\otimes m}}(\tau_U) - \text{Tr}_{\mathcal{Z}^{\otimes m}}(\rho_U)) \otimes \omega_{\mathcal{Z}}^{\otimes m} \right\|_1 \\ &\leq \|\tau_U - \text{Tr}_{\mathcal{Z}^{\otimes m}}(\tau_U) \otimes \omega_{\mathcal{Z}}^{\otimes m}\|_1 + 2\|\rho_U - \tau_U\|_1,\end{aligned}\quad (8.393)$$

and so it remains to consider the average value of the two terms in the final expression of this inequality. When considering the first term in the final expression of (8.393), it may be noted that

$$\text{im}(\tau_U) \subseteq \text{im}\left(\text{Tr}_{\mathcal{Z}^{\otimes m}}(\tau_U) \otimes \omega_{\mathcal{Z}}^{\otimes m}\right) \quad (8.394)$$

and therefore

$$\begin{aligned}\text{rank}\left(\tau_U - \text{Tr}_{\mathcal{Z}^{\otimes m}}(\tau_U) \otimes \omega_{\mathcal{Z}}^{\otimes m}\right) &\leq \text{rank}\left(\text{Tr}_{\mathcal{Z}^{\otimes m}}(\tau_U) \otimes \omega_{\mathcal{Z}}^{\otimes m}\right) \\ &\leq 2^m \text{rank}(\text{Tr}_{\mathcal{Y}^{\otimes n}}(B_n B_n^*)) \leq 2^{n(1+\varepsilon)\text{H}(\Psi(\omega))+m+1}.\end{aligned}\quad (8.395)$$

In addition, one has

$$\text{Tr}(\sigma^2) = \text{Tr}\left(\left(\frac{1}{k^n} \text{Tr}_{\mathcal{W}^{\otimes n}}(B_n B_n^*)\right)^2\right) \leq 2^{-n(1-\varepsilon)\text{H}(\Phi(\omega))+4}. \quad (8.396)$$

Making use of Lemma 8.49, it therefore follows that

$$\begin{aligned}\int \|\tau_U - \text{Tr}_{\mathcal{Z}^{\otimes m}}(\tau_U) \otimes \omega_{\mathcal{Z}}^{\otimes m}\|_1^2 d\eta(U) &\leq 2^{n(1+\varepsilon)\text{H}(\Psi(\omega))+m+1} \int \|\tau_U - \text{Tr}_{\mathcal{Z}^{\otimes m}}(\tau_U) \otimes \omega_{\mathcal{Z}}^{\otimes m}\|_2^2 d\eta(U) \\ &\leq 2^{n((1+\varepsilon)\text{H}(\Psi(\omega)) - (1-\varepsilon)\text{H}(\Phi(\omega)) + m + 5)} \\ &= 2^{-n(\text{I}_c(\omega; \Phi) - 2\varepsilon(\text{H}(\Phi(\omega)) + \text{H}(\Psi(\omega))) + m + 5)}.\end{aligned}\quad (8.397)$$

By the assumption (8.372), and using the fact that $m = \lfloor \alpha n \rfloor$, one has that this quantity approaches 0 in the limit as n approaches infinity. It therefore

holds (by Jensen's inequality) that the quantity

$$\int \|\tau_U - \text{Tr}_{\mathcal{Z}^{\otimes m}}(\tau_U) \otimes \omega_{\mathcal{Z}}^{\otimes m}\|_1 d\eta(U) \quad (8.398)$$

also approaches 0 in the limit as n approaches infinity. The average value of the second term in the final expression of (8.393) may be upper-bounded as

$$\begin{aligned}\int \|\rho_U - \tau_U\|_1 d\eta(U) &= \frac{k^n}{2^m} \int \left\| (\mathbf{1}_{\mathcal{W}^{\otimes n}} \otimes V^T U^T) (\xi - \sigma) (\mathbf{1}_{\mathcal{W}^{\otimes n}} \otimes V^T U^T)^* \right\|_1 d\eta(U) \\ &\leq \|\xi - \sigma\|_1 \leq \frac{1}{k^n} \|\text{vec}(A_n) \text{vec}(A_n)^* - \text{vec}(B_n) \text{vec}(B_n)^*\|_1 \\ &\leq \frac{2}{k^{n/2}} \|A_n - B_n\|_2 \leq 6 \zeta_{n,\varepsilon}^{1/4}\end{aligned}\quad (8.399)$$

by Lemma 8.52. Once again, this quantity approaches 0 in the limit as n approaches infinity. It follows that the entanglement generation capacity of Φ is at least α , which completes the proof. \square

Corollary 8.54 *Let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces, let $\Phi \in \text{C}(\mathcal{X}, \mathcal{Y})$ be a channel, and let $\sigma \in \text{D}(\mathcal{X})$ be a density operator. The quantum capacity of Φ is lower-bounded by the coherent information of σ through Φ :*

$$\text{I}_c(\sigma; \Phi) \leq \text{Q}(\Phi). \quad (8.400)$$

Proof Observe first that it is a consequence of Theorem 8.53 that

$$\text{I}_c(\omega_{\mathcal{V}}; \Phi) \leq \text{Q}(\Phi) \quad (8.401)$$

for every nontrivial subspace $\mathcal{V} \subseteq \mathcal{X}$, where

$$\omega_{\mathcal{V}} = \frac{\Pi_{\mathcal{V}}}{\text{dim}(\mathcal{V})} \quad (8.402)$$

is the flat state corresponding to the subspace \mathcal{V} . To verify that this is so, let \mathcal{Z} be any complex Euclidean space with $\text{dim}(\mathcal{Z}) = \text{dim}(\mathcal{V})$, let $V \in \text{U}(\mathcal{Z}, \mathcal{X})$ be an isometry such that $VV^* = \Pi_{\mathcal{V}}$, and define a channel $\Xi \in \text{C}(\mathcal{Z}, \mathcal{Y})$ as

$$\Xi(Z) = \Phi(VZV^*) \quad (8.403)$$

for all $Z \in \text{L}(\mathcal{Z})$. It is evident that $\text{Q}(\Xi) \leq \text{Q}(\Phi)$; the channel Φ emulates Ξ , so for every positive integer n it holds that $\Phi^{\otimes n}$ emulates every channel that can be emulated by $\Xi^{\otimes n}$. It follows that

$$\begin{aligned}\text{Q}(\Phi) &\geq \text{Q}(\Xi) = \text{Q}_{\text{EG}}(\Xi) \geq \text{I}_c(\omega_{\mathcal{Z}}; \Xi) \\ &= \text{I}_c(V\omega_{\mathcal{Z}}V^*; \Phi) = \text{I}_c(\omega_{\mathcal{V}}; \Phi),\end{aligned}\quad (8.404)$$

as claimed.

Now, let $A \in \mathcal{U}(\mathcal{X}, \mathcal{Y} \otimes \mathcal{W})$ be an isometry such that

$$\Phi(X) = \text{Tr}_{\mathcal{W}}(AXA^*) \quad (8.405)$$

for all $X \in \mathcal{L}(\mathcal{X})$, for a suitable choice of a complex Euclidean space \mathcal{W} , and define a channel $\Psi \in \mathcal{C}(\mathcal{X}, \mathcal{W})$ as

$$\Psi(X) = \text{Tr}_{\mathcal{Y}}(AXA^*) \quad (8.406)$$

for all $X \in \mathcal{L}(\mathcal{X})$. It therefore holds that Ψ is complementary to Φ , so that

$$\mathcal{I}_c(\sigma; \Phi) = \mathcal{H}(\Phi(\sigma)) - \mathcal{H}(\Psi(\sigma)). \quad (8.407)$$

Let

$$\sigma = \sum_{a \in \Sigma} p(a) x_a x_a^* \quad (8.408)$$

be a spectral decomposition of σ , and let

$$\omega_{n,\varepsilon} = \frac{\Lambda_{n,\varepsilon}}{\text{Tr}(\Lambda_{n,\varepsilon})} \in \mathcal{D}(\mathcal{X}^{\otimes n}) \quad (8.409)$$

for each positive integer n and each positive real number $\varepsilon > 0$, for $\Lambda_{n,\varepsilon}$ denoting the projection onto the ε -strongly typical subspace of $\mathcal{X}^{\otimes n}$, with respect to the spectral decomposition (8.408).

Next, let $\varepsilon > 0$ be a positive real number, to be chosen arbitrarily. By Lemma 8.36, it follows that there must exist a positive integer n_0 such that, for all $n \geq n_0$, one has

$$\left| \frac{1}{n} \mathcal{H}(\Phi^{\otimes n}(\omega_{n,\varepsilon})) - \mathcal{H}(\Phi(\sigma)) \right| \leq (2\mathcal{H}(\sigma) + \mathcal{H}(\Phi(\sigma)) + 1)\varepsilon. \quad (8.410)$$

Along similar lines, there must exist a positive integer n_1 such that, for all $n \geq n_1$, one has

$$\left| \frac{1}{n} \mathcal{H}(\Psi^{\otimes n}(\omega_{n,\varepsilon})) - \mathcal{H}(\Psi(\sigma)) \right| \leq (2\mathcal{H}(\sigma) + \mathcal{H}(\Psi(\sigma)) + 1)\varepsilon. \quad (8.411)$$

There must therefore exist a positive integer n such that

$$\left| \frac{1}{n} \mathcal{I}_c(\omega_{n,\varepsilon}; \Phi^{\otimes n}) - \mathcal{I}_c(\sigma; \Phi) \right| \leq (4\mathcal{H}(\sigma) + \mathcal{H}(\Phi(\sigma)) + \mathcal{H}(\Psi(\sigma)) + 2)\varepsilon. \quad (8.412)$$

By the argument presented at the beginning of the proof, it holds that

$$\frac{\mathcal{I}_c(\omega_{n,\varepsilon}; \Phi^{\otimes n})}{n} \leq \frac{\mathcal{Q}(\Phi^{\otimes n})}{n} = \mathcal{Q}(\Phi), \quad (8.413)$$

and therefore

$$\mathcal{Q}(\Phi) \geq \mathcal{I}_c(\sigma; \Phi) - (4\mathcal{H}(\sigma) + \mathcal{H}(\Phi(\sigma)) + \mathcal{H}(\Psi(\sigma)) + 2)\varepsilon. \quad (8.414)$$

As ε has been chosen to be an arbitrary positive real number, it follows that

$$\mathcal{Q}(\Phi) \geq \mathcal{I}_c(\sigma; \Phi), \quad (8.415)$$

which completes the proof. \square

Finally, the quantum capacity theorem may be stated and proved.

Theorem 8.55 (Quantum capacity theorem) *Let \mathcal{X} and \mathcal{Y} be complex Euclidean spaces and let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ be a channel. It holds that*

$$\mathcal{Q}(\Phi) = \lim_{n \rightarrow \infty} \frac{\mathcal{I}_c(\Phi^{\otimes n})}{n} \quad (8.416)$$

Proof For every positive integer n and every density operator $\sigma \in \mathcal{D}(\mathcal{X}^{\otimes n})$, one has

$$\mathcal{I}_c(\sigma; \Phi^{\otimes n}) \leq \mathcal{Q}(\Phi^{\otimes n}) = n\mathcal{Q}(\Phi) \quad (8.417)$$

by Corollary 8.54, and therefore

$$\frac{\mathcal{I}_c(\Phi^{\otimes n})}{n} \leq \mathcal{Q}(\Phi). \quad (8.418)$$

If it holds that $\mathcal{Q}(\Phi) = 0$, then the theorem evidently follows, so it will be assumed that $\mathcal{Q}(\Phi) > 0$ for the remainder of the proof.

Suppose that $\alpha > 0$ is an achievable rate for entanglement generation through Φ , let $\delta \in (0, 1)$ be chosen arbitrarily, and set $\varepsilon = \delta^2/2$. Also let $\Gamma = \{0, 1\}$ and $\mathcal{Z} = \mathbb{C}^\Gamma$. As α is an achievable rate for entanglement generation through Φ , it holds, for all but finitely many positive integers n and for $m = \lfloor \alpha n \rfloor$, that there must exist a unit vector $u \in \mathcal{X}^{\otimes n} \otimes \mathcal{Z}^{\otimes m}$ and a channel $\Xi \in \mathcal{C}(\mathcal{Y}^{\otimes n}, \mathcal{Z}^{\otimes m})$ such that

$$\mathbb{F}\left(2^{-m} \text{vec}(\mathbf{1}_{\mathcal{Z}^{\otimes m}}) \text{vec}(\mathbf{1}_{\mathcal{Z}^{\otimes m}})^*, (\Xi \Phi^{\otimes n} \otimes \mathbf{1}_{\mathcal{L}(\mathcal{Z})}^{\otimes m})(uu^*)\right) > 1 - \varepsilon, \quad (8.419)$$

and therefore

$$\left\| 2^{-m} \text{vec}(\mathbf{1}_{\mathcal{Z}^{\otimes m}}) \text{vec}(\mathbf{1}_{\mathcal{Z}^{\otimes m}})^* - (\Xi \Phi^{\otimes n} \otimes \mathbf{1}_{\mathcal{L}(\mathcal{Z})}^{\otimes m})(uu^*) \right\|_1 < 2\delta \quad (8.420)$$

by one of the Fuchs–van de Graaf inequalities (Theorem 3.33). For any unit vector $u \in \mathcal{X}^{\otimes n} \otimes \mathcal{Z}^{\otimes m}$ for which the inequality (8.420) holds, one concludes from the Fannes–Audenaert inequality (Theorem 5.26) that for

$$\rho = \text{Tr}_{\mathcal{Z}^{\otimes m}}(uu^*) \quad (8.421)$$

the inequalities

$$H((\Xi\Phi^{\otimes n} \otimes \mathbf{1}_{L(z)}^{\otimes m})(uu^*)) \leq 2\delta m + 1 \quad (8.422)$$

and

$$m - H(\Xi\Phi^{\otimes n}(\rho)) \leq \delta m + 1 \quad (8.423)$$

are satisfied. Together with Proposition 8.15, these inequalities imply that

$$I_c(\rho; \Phi^{\otimes n}) \geq I_c(\rho; \Xi\Phi^{\otimes n}) \geq (1 - 3\delta)m - 2. \quad (8.424)$$

As $m = \lfloor \alpha n \rfloor \geq \alpha n - 1$, it follows that

$$\frac{I_c(\rho; \Phi^{\otimes n})}{n} \geq (1 - 3\delta)\alpha - \frac{3}{n} \quad (8.425)$$

It has been proved that for any achievable rate $\alpha > 0$ for entanglement generation through Φ , and for any $\delta > 0$, that

$$(1 - 3\delta)\alpha - \frac{3}{n} \leq \frac{I_c(\rho; \Phi^{\otimes n})}{n} \leq Q(\Phi) \quad (8.426)$$

for all but finitely many positive integers n . Because $Q(\Phi)$ is equal to the supremum value of all achievable rates for entanglement generation through Φ , and $\delta > 0$ may be chosen to be arbitrarily small, the required equality (8.416) follows. \square

8.3 Non-additivity and super-activation

Expressions for the classical and quantum capacities of a quantum channel are given by regularizations of the Holevo capacity and maximum coherent information,

$$C(\Phi) = \lim_{n \rightarrow \infty} \frac{\chi(\Phi^{\otimes n})}{n} \quad \text{and} \quad Q(\Psi) = \lim_{n \rightarrow \infty} \frac{I_c(\Psi^{\otimes n})}{n}, \quad (8.427)$$

as has been established by the Holevo–Schumacher–Westmoreland theorem and quantum capacity theorem (Theorems 8.27 and 8.55). Non-regularized analogues of these formulas do not, in general, hold. In particular, the strict inequalities

$$\chi(\Phi \otimes \Phi) > 2\chi(\Phi) \quad \text{and} \quad I_c(\Psi \otimes \Psi) > 2I_c(\Psi) \quad (8.428)$$

hold for a suitable choice of channels Φ and Ψ , as is demonstrated in the subsections that follow. These examples reveal that the Holevo capacity does not coincide directly with the classical capacity, and likewise for the maximum coherent information and quantum capacity.

With respect to the Holevo capacity, the fact that a strict inequality may hold for some channels Φ in (8.428) will be demonstrated in Section 8.3.1, through the use of Theorem 7.49 from the previous chapter. The existence of such channels is far from obvious, and no explicit examples are known at the time of this book's writing—it is only the existence of such channels that is known. The now falsified conjecture that the equality

$$\chi(\Phi_0 \otimes \Phi_1) = \chi(\Phi_0) + \chi(\Phi_1) \quad (8.429)$$

should hold for all choices of channels Φ_0 and Φ_1 was known for some time as the *additivity conjecture*.

In contrast, it is not difficult to find an example of a channel Ψ for which a strict inequality in (8.428) holds. There are, in fact, very striking examples of channels that go beyond the demonstration of non-additivity of maximum coherent information. In particular, one may find channels Ψ_0 and Ψ_1 such that both Ψ_0 and Ψ_1 have zero quantum capacity, and therefore

$$I_c(\Psi_0) = I_c(\Psi_1) = 0, \quad (8.430)$$

but for which

$$I_c(\Psi_0 \otimes \Psi_1) > 0, \quad (8.431)$$

and therefore $\Psi_0 \otimes \Psi_1$ has nonzero quantum capacity. This phenomenon is known as *super-activation*, and is discussed in Section 8.3.2. From such a choice of channels Ψ_0 and Ψ_1 , the construction of a channel Ψ for which the strict inequality (8.428) holds is possible.

8.3.1 Non-additivity of the Holevo capacity

The fact that there exists a channel Φ for which

$$\chi(\Phi \otimes \Phi) > 2\chi(\Phi) \quad (8.432)$$

is demonstrated below. The proof makes use of Theorem 7.49, together with two basic ideas: one concerns the *direct sum* of two channels, and the other is a construction that relates the minimum output entropy of a given channel to the Holevo capacity of a channel constructed from the one given.

Direct sums of channels and their minimum output entropy

The direct sum of two maps is defined as follows. (One may also consider direct sums of more than two maps, but it is sufficient for the needs of the present section to consider the case of just two maps.)

Definition 8.56 Let $\mathcal{X}_0, \mathcal{X}_1, \mathcal{Y}_0$, and \mathcal{Y}_1 be complex Euclidean spaces and let $\Phi_0 \in \mathcal{T}(\mathcal{X}_0, \mathcal{Y}_0)$ and $\Phi_1 \in \mathcal{T}(\mathcal{X}_1, \mathcal{Y}_1)$ be maps. The direct sum of Φ_0 and Φ_1 is the map $\Phi_0 \oplus \Phi_1 \in \mathcal{T}(\mathcal{X}_0 \oplus \mathcal{X}_1, \mathcal{Y}_0 \oplus \mathcal{Y}_1)$ defined as

$$(\Phi_0 \oplus \Phi_1) \begin{pmatrix} X_0 & \cdot \\ \cdot & X_1 \end{pmatrix} = \begin{pmatrix} \Phi_0(X_0) & 0 \\ 0 & \Phi_1(X_1) \end{pmatrix} \quad (8.433)$$

for every $X_0 \in \mathcal{L}(\mathcal{X}_0)$ and $X_1 \in \mathcal{L}(\mathcal{X}_1)$. The dots in (8.433) indicate arbitrary operators in $\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)$ and $\mathcal{L}(\mathcal{X}_0, \mathcal{X}_1)$ that have no influence on the output of the map $\Phi_0 \oplus \Phi_1$.

The direct sum of two channels is also a channel, as is established by the following straightforward proposition.

Proposition 8.57 Let $\mathcal{X}_0, \mathcal{X}_1, \mathcal{Y}_0$, and \mathcal{Y}_1 be complex Euclidean spaces and let $\Phi_0 \in \mathcal{C}(\mathcal{X}_0, \mathcal{Y}_0)$ and $\Phi_1 \in \mathcal{C}(\mathcal{X}_1, \mathcal{Y}_1)$ be channels. The direct sum of Φ_0 and Φ_1 is a channel: $\Phi_0 \oplus \Phi_1 \in \mathcal{C}(\mathcal{X}_0 \oplus \mathcal{X}_1, \mathcal{Y}_0 \oplus \mathcal{Y}_1)$.

Proof It is immediate from the definition of the direct sum of Φ_0 and Φ_1 that $\Phi_0 \oplus \Phi_1$ is trace preserving, so it suffices to prove that $\Phi_0 \oplus \Phi_1$ is completely positive. Because Φ_0 and Φ_1 are completely positive, Kraus representations of the form

$$\Phi_0(X_0) = \sum_{a \in \Sigma} A_a X_0 A_a^* \quad \text{and} \quad \Phi_1(X_1) = \sum_{b \in \Gamma} B_b X_1 B_b^* \quad (8.434)$$

of these maps must exist. Through a direct computation, one may verify that

$$\begin{aligned} (\Phi_0 \oplus \Phi_1)(X) &= \sum_{a \in \Sigma} \begin{pmatrix} A_a & 0 \\ 0 & 0 \end{pmatrix} X \begin{pmatrix} A_a & 0 \\ 0 & 0 \end{pmatrix}^* \\ &\quad + \sum_{b \in \Gamma} \begin{pmatrix} 0 & 0 \\ 0 & B_b \end{pmatrix} X \begin{pmatrix} 0 & 0 \\ 0 & B_b \end{pmatrix}^* \end{aligned} \quad (8.435)$$

for all $X \in \mathcal{L}(\mathcal{X}_0 \oplus \mathcal{X}_1)$. It follows that $\Phi_0 \oplus \Phi_1$ is completely positive, as required. \square

By Theorem 7.49, there exist channels Ψ_0 and Ψ_1 such that

$$\mathbb{H}_{\min}(\Phi_0 \otimes \Phi_1) < \mathbb{H}_{\min}(\Phi_0) + \mathbb{H}_{\min}(\Phi_1). \quad (8.436)$$

It is possible to obtain, from this fact, an example of a single channel Φ such that

$$\mathbb{H}_{\min}(\Phi \otimes \Phi) < 2 \mathbb{H}_{\min}(\Phi). \quad (8.437)$$

The following corollary (to Theorem 7.49) establishes that this is so.

Corollary 8.58 There exists a channel $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$, for some choice of complex Euclidean spaces \mathcal{X} and \mathcal{Y} , such that

$$\mathbb{H}_{\min}(\Phi \otimes \Phi) < 2 \mathbb{H}_{\min}(\Phi). \quad (8.438)$$

Proof By Theorem 7.49, there exist complex Euclidean spaces \mathcal{Z} and \mathcal{W} and channels $\Psi_0, \Psi_1 \in \mathcal{C}(\mathcal{Z}, \mathcal{W})$ such that

$$\mathbb{H}_{\min}(\Psi_0 \otimes \Psi_1) < \mathbb{H}_{\min}(\Psi_0) + \mathbb{H}_{\min}(\Psi_1). \quad (8.439)$$

Assume that such a choice of channels has been fixed for the remainder of the proof.

Let $\sigma_0, \sigma_1 \in \mathcal{D}(\mathcal{Z})$ be density operators satisfying

$$\mathbb{H}(\Psi_0(\sigma_0)) = \mathbb{H}_{\min}(\Psi_0) \quad \text{and} \quad \mathbb{H}(\Psi_1(\sigma_1)) = \mathbb{H}_{\min}(\Psi_1), \quad (8.440)$$

and define channels $\Phi_0, \Phi_1 \in \mathcal{C}(\mathcal{Z}, \mathcal{W} \otimes \mathcal{W})$ as

$$\Phi_0(Z) = \Psi_0(Z) \otimes \Psi_1(\sigma_1) \quad \text{and} \quad \Phi_1(Z) = \Psi_0(\sigma_0) \otimes \Psi_1(Z) \quad (8.441)$$

for all $Z \in \mathcal{L}(\mathcal{Z})$. Observe that

$$\mathbb{H}_{\min}(\Phi_0) = \mathbb{H}_{\min}(\Psi_0) + \mathbb{H}_{\min}(\Psi_1) = \mathbb{H}_{\min}(\Phi_1) \quad (8.442)$$

and

$$\begin{aligned} \mathbb{H}_{\min}(\Phi_0 \otimes \Phi_1) &= \mathbb{H}_{\min}(\Psi_0 \otimes \Psi_1) + \mathbb{H}_{\min}(\Psi_0) + \mathbb{H}_{\min}(\Psi_1) \\ &< 2 \mathbb{H}_{\min}(\Psi_0) + 2 \mathbb{H}_{\min}(\Psi_1) = \mathbb{H}_{\min}(\Phi_0) + \mathbb{H}_{\min}(\Phi_1). \end{aligned} \quad (8.443)$$

Finally, let $\mathcal{X} = \mathcal{Z} \oplus \mathcal{Z}$ and $\mathcal{Y} = (\mathcal{W} \otimes \mathcal{W}) \oplus (\mathcal{W} \otimes \mathcal{W})$, and define $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ as

$$\Phi = \Phi_0 \oplus \Phi_1. \quad (8.444)$$

It remains to verify that $\mathbb{H}_{\min}(\Phi \otimes \Phi) < 2 \mathbb{H}_{\min}(\Phi)$.

For any state $\rho \in \mathcal{D}(\mathcal{Z} \oplus \mathcal{Z})$, one may write

$$\rho = \begin{pmatrix} \lambda \rho_0 & Z \\ Z^* & (1 - \lambda) \rho_1 \end{pmatrix} \quad (8.445)$$

for some choice of $\lambda \in [0, 1]$, $\rho_0, \rho_1 \in \mathcal{D}(\mathcal{Z})$, and $Z \in \mathcal{L}(\mathcal{Z})$. Evaluating Φ on such a state ρ yields

$$\Phi(\rho) = \begin{pmatrix} \lambda \Phi_0(\rho_0) & 0 \\ 0 & (1 - \lambda) \Phi_1(\rho_1) \end{pmatrix}, \quad (8.446)$$

so that

$$\mathbb{H}(\Phi(\rho)) = \lambda \mathbb{H}(\Phi_0(\rho_0)) + (1 - \lambda) \mathbb{H}(\Phi_1(\rho_1)) + \mathbb{H}(\lambda, 1 - \lambda). \quad (8.447)$$

One concludes that

$$H_{\min}(\Phi) = H_{\min}(\Phi_0) = H_{\min}(\Phi_1). \quad (8.448)$$

Finally, define an isometry $V \in U(\mathcal{Z} \otimes \mathcal{Z}, (\mathcal{Z} \oplus \mathcal{Z}) \otimes (\mathcal{Z} \oplus \mathcal{Z}))$ by the equation

$$V(z_0 \otimes z_1) = (z_0 \oplus 0) \otimes (0 \oplus z_1) \quad (8.449)$$

holding for all $z_0, z_1 \in \mathcal{Z}$. For every choice of operators $Z_0, Z_1 \in L(\mathcal{Z})$ it therefore holds that

$$V(Z_0 \otimes Z_1)V^* = \begin{pmatrix} Z_0 & 0 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 0 & Z_1 \end{pmatrix}, \quad (8.450)$$

so that

$$(\Phi \otimes \Phi)(V(Z_0 \otimes Z_1)V^*) = \begin{pmatrix} \Phi_0(Z_0) & 0 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 0 & \Phi_1(Z_1) \end{pmatrix}. \quad (8.451)$$

One concludes that

$$H((\Phi \otimes \Phi)(V\xi V^*)) = H((\Phi_0 \otimes \Phi_1)(\xi)) \quad (8.452)$$

for every density operator $\xi \in D(\mathcal{Z} \otimes \mathcal{Z})$, and therefore

$$H_{\min}(\Phi \otimes \Phi) \leq H_{\min}(\Phi_0 \otimes \Phi_1) < H_{\min}(\Phi_0) + H_{\min}(\Phi_1) = 2H_{\min}(\Phi), \quad (8.453)$$

as required. \square

From low minimum output entropy to high Holevo capacity

The construction to be described below allows one to conclude that there exists a channel Ψ for which the Holevo capacity is super-additive, meaning that

$$\chi(\Psi \otimes \Psi) > 2\chi(\Psi), \quad (8.454)$$

by means of Corollary 8.58.

Suppose that \mathcal{X} and \mathcal{Y} are complex Euclidean spaces and $\Phi \in C(\mathcal{X}, \mathcal{Y})$ is an arbitrary channel. Suppose further that Σ is an alphabet and

$$\{U_a : a \in \Sigma\} \subset U(\mathcal{Y}) \quad (8.455)$$

is a collection of unitary operators with the property that the completely depolarizing channel $\Omega \in C(\mathcal{Y})$ is given by

$$\Omega(Y) = \frac{1}{|\Sigma|} \sum_{a \in \Sigma} U_a Y U_a^* \quad (8.456)$$

for all $Y \in L(\mathcal{Y})$. (Such a collection may, for instance, be derived from the

discrete Weyl operators defined in Section 4.1.2.) Let $\mathcal{Z} = \mathbb{C}^\Sigma$ and define a new channel $\Psi \in C(\mathcal{Z} \otimes \mathcal{X}, \mathcal{Y})$ by the equation

$$\Psi(E_{a,b} \otimes X) = \begin{cases} U_a \Phi(X) U_a^* & \text{if } a = b \\ 0 & \text{otherwise} \end{cases} \quad (8.457)$$

holding for all $a, b \in \Sigma$ and $X \in L(\mathcal{X})$.

The action of the channel Ψ may alternatively be described as follows. A pair of registers (Z, X) is taken as input, and a measurement of the register Z with respect to the standard basis of \mathcal{Z} is made, yielding a symbol $a \in \Sigma$. The channel Φ is applied to X , resulting in a register Y , and the unitary channel described by U_a is applied to Y . The measurement outcome a is discarded and Y is taken to be the output of the channel.

As the following proposition shows, the Holevo capacity of the channel Ψ constructed in this way is determined by the minimum output entropy of the channel Φ .

Proposition 8.59 *Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} , let Σ be an alphabet, let $\{U_a : a \in \Sigma\} \subset U(\mathcal{Y})$ be a collection of unitary operators for which the equation (8.456) holds for all $Y \in L(\mathcal{Y})$, let $\mathcal{Z} = \mathbb{C}^\Sigma$, and let $\Psi \in C(\mathcal{Z} \otimes \mathcal{X}, \mathcal{Y})$ be a channel defined by the equation (8.457) holding for all $a, b \in \Sigma$ and $X \in L(\mathcal{X})$. It holds that*

$$\chi(\Psi) = \log(\dim(\mathcal{Y})) - H_{\min}(\Phi). \quad (8.458)$$

Proof Consider first the ensemble $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{Z} \otimes \mathcal{X})$ defined as

$$\eta(a) = \frac{1}{|\Sigma|} E_{a,a} \otimes \rho \quad (8.459)$$

for all $a \in \Sigma$, where $\rho \in D(\mathcal{X})$ is any state for which

$$H_{\min}(\Phi) = H(\Phi(\rho)). \quad (8.460)$$

One has

$$\begin{aligned} \chi(\Psi(\eta)) &= H\left(\frac{1}{|\Sigma|} \sum_{a \in \Sigma} U_a \Phi(\rho) U_a^*\right) - \frac{1}{|\Sigma|} \sum_{a \in \Sigma} H(U_a \Phi(\rho) U_a^*) \\ &= H(\Omega(\rho)) - H(\Phi(\rho)) \\ &= \log(\dim(\mathcal{Y})) - H_{\min}(\Phi). \end{aligned} \quad (8.461)$$

It therefore holds that

$$\chi(\Psi) \geq \log(\dim(\mathcal{Y})) - H_{\min}(\Phi). \quad (8.462)$$

Next, consider an arbitrary state $\sigma \in \mathcal{D}(\mathcal{Z} \otimes \mathcal{X})$. For $\Delta \in \mathcal{C}(\mathcal{Z})$ denoting the completely dephasing channel, one may write

$$(\Delta \otimes \mathbf{1}_{L(\mathcal{X})})(\sigma) = \sum_{a \in \Sigma} q(a) E_{a,a} \otimes \xi_a, \quad (8.463)$$

for some choice of a probability vector $q \in \mathcal{P}(\Sigma)$ and a collection of states

$$\{\xi_a : a \in \Sigma\} \subseteq \mathcal{D}(\mathcal{X}). \quad (8.464)$$

It holds that

$$\Psi(\sigma) = \sum_{a \in \Sigma} q(a) U_a \Phi(\xi_a) U_a^*, \quad (8.465)$$

and therefore

$$\mathsf{H}(\Psi(\sigma)) \geq \sum_{a \in \Sigma} q(a) \mathsf{H}(\Phi(\xi_a)) \geq \mathsf{H}_{\min}(\Phi) \quad (8.466)$$

by the concavity of the von Neumann entropy function (Theorem 5.23).

Finally, consider an arbitrary ensemble $\eta : \Gamma \rightarrow \text{Pos}(\mathcal{Z} \otimes \mathcal{X})$, written as

$$\eta(b) = p(b) \sigma_b \quad (8.467)$$

for each $b \in \Gamma$, for $p \in \mathcal{P}(\Gamma)$ being a probability vector and

$$\{\sigma_b : b \in \Gamma\} \subseteq \mathcal{D}(\mathcal{Z} \otimes \mathcal{X}) \quad (8.468)$$

being a collection of states. It holds that

$$\begin{aligned} \chi(\Psi(\eta)) &= \mathsf{H}\left(\sum_{b \in \Gamma} p(b) \Psi(\sigma_b)\right) - \sum_{b \in \Gamma} p(b) \mathsf{H}(\Psi(\sigma_b)) \\ &\leq \log(\dim(\mathcal{Y})) - \mathsf{H}_{\min}(\Phi). \end{aligned} \quad (8.469)$$

The ensemble η was chosen arbitrarily, and therefore

$$\chi(\Psi) \leq \log(\dim(\mathcal{Y})) - \mathsf{H}_{\min}(\Phi), \quad (8.470)$$

which completes the proof. \square

Theorem 8.60 *There exists a channel $\Psi \in \mathcal{C}(\mathcal{W}, \mathcal{Y})$, for some choice of complex Euclidean spaces \mathcal{W} and \mathcal{Y} , such that*

$$\chi(\Psi \otimes \Psi) > 2\chi(\Psi). \quad (8.471)$$

Proof By Corollary 8.58 there exist complex Euclidean spaces \mathcal{X} and \mathcal{Y} and a channel $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ for which the inequality

$$\mathsf{H}_{\min}(\Phi \otimes \Phi) < 2\mathsf{H}_{\min}(\Phi) \quad (8.472)$$

holds. Let Σ be an alphabet and let

$$\{U_a : a \in \Sigma\} \subset \mathcal{U}(\mathcal{Y}) \quad (8.473)$$

be a collection of unitary operators for which

$$\Omega(Y) = \frac{1}{|\Sigma|} \sum_{a \in \Sigma} U_a Y U_a^* \quad (8.474)$$

for all $Y \in \mathcal{L}(\mathcal{Y})$. Also let $\mathcal{Z} = \mathbb{C}^\Sigma$ and let $\Psi \in \mathcal{C}(\mathcal{Z} \otimes \mathcal{X}, \mathcal{Y})$ be the channel defined by the equation (8.457) above for all $a, b \in \Sigma$ and $X \in \mathcal{L}(\mathcal{X})$.

Up to a permutation of the tensor factors of its input space, $\Psi \otimes \Psi$ is equivalent to the channel $\Xi \in \mathcal{C}((\mathcal{Z} \otimes \mathcal{Z}) \otimes (\mathcal{X} \otimes \mathcal{X}), \mathcal{Y} \otimes \mathcal{Y})$ that would be obtained from the channel $\Phi \otimes \Phi$ by means of a similar construction, using the collection of unitary operators

$$\{U_a \otimes U_b : (a, b) \in \Sigma \times \Sigma\} \subset \mathcal{U}(\mathcal{Y} \otimes \mathcal{Y}). \quad (8.475)$$

It therefore follows from Proposition 8.59 that

$$\chi(\Psi) = \log(\dim(\mathcal{Y})) - \mathsf{H}_{\min}(\Phi) \quad (8.476)$$

while

$$\chi(\Psi \otimes \Psi) = \log(\dim(\mathcal{Y} \otimes \mathcal{Y})) - \mathsf{H}_{\min}(\Phi \otimes \Phi) > 2\chi(\Psi). \quad (8.477)$$

Taking $\mathcal{W} = \mathcal{Z} \otimes \mathcal{X}$, the theorem is therefore proved. \square

One consequence of this theorem is that an analogous statement to the Holevo–Schumacher–Westmoreland theorem (Theorem 8.27), but without a regularization, does not hold in general. That is, because

$$\mathsf{C}(\Phi) \geq \frac{\chi(\Phi \otimes \Phi)}{2}, \quad (8.478)$$

it is the case that $\mathsf{C}(\Phi) > \chi(\Phi)$ for some choices of a channel Φ .

8.3.2 Super-activation of quantum channel capacity

The purpose of the present subsection is to demonstrate the phenomenon of *super-activation*, in which the tensor product of two zero-capacity channels have positive quantum capacity. As a byproduct, one obtains an example of a channel Ψ satisfying $\mathsf{I}_c(\Psi \otimes \Psi) > 2\mathsf{I}_c(\Psi)$.

Two classes of zero-capacity channels

It is possible to prove that certain classes of channels have zero quantum capacity. Self-complementary channels and channels whose Choi operators are PPT fall into this category. The following proposition establishes that channels whose Choi operators are PPT must have zero capacity.

Proposition 8.61 *Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} , such that $J(\Phi) \in \text{PPT}(\mathcal{Y} : \mathcal{X})$. It holds that $Q(\Phi) = 0$.*

Proof The first step of the proof is to establish that, for every choice of a complex Euclidean space \mathcal{W} and a state $\rho \in D(\mathcal{X} \otimes \mathcal{W})$, one has

$$(\Phi \otimes \mathbb{1}_{L(\mathcal{W})})(\rho) \in \text{PPT}(\mathcal{Y} : \mathcal{W}). \quad (8.479)$$

Toward this goal, observe that for any choice of a complex Euclidean space \mathcal{W} and a positive semidefinite operator $P \in \text{Pos}(\mathcal{X} \otimes \mathcal{W})$, there must exist a completely positive map $\Psi_P \in \text{CP}(\mathcal{X}, \mathcal{W})$ satisfying

$$P = (\mathbb{1}_{L(\mathcal{X})} \otimes \Psi_P)(\text{vec}(\mathbb{1}_{\mathcal{X}}) \text{vec}(\mathbb{1}_{\mathcal{X}})^*). \quad (8.480)$$

The map Ψ_P is, in fact, uniquely defined by this requirement; one may obtain its Choi representation by swapping the tensor factors of P . It follows that, for any complex Euclidean space \mathcal{W} and any state $\rho \in D(\mathcal{X} \otimes \mathcal{W})$, one must have

$$\begin{aligned} & (\mathbb{T} \otimes \mathbb{1}_{L(\mathcal{W})})((\Phi \otimes \mathbb{1}_{L(\mathcal{W})})(\rho)) \\ &= (\mathbb{1}_{L(\mathcal{Y})} \otimes \Psi_\rho)((\mathbb{T} \otimes \mathbb{1}_{L(\mathcal{X})})(J(\Phi))) \in \text{Pos}(\mathcal{Y} : \mathcal{W}) \end{aligned} \quad (8.481)$$

by virtue of the fact that Ψ_ρ is completely positive and $J(\Phi) \in \text{PPT}(\mathcal{Y} : \mathcal{X})$, which establishes (8.479).

As $J(\Phi) \in \text{PPT}(\mathcal{Y} : \mathcal{X})$, it follows that

$$J(\Phi^{\otimes n}) \in \text{PPT}(\mathcal{Y}^{\otimes n} : \mathcal{X}^{\otimes n}) \quad (8.482)$$

for every positive integer n . For every choice of positive integers n and m , for $\mathcal{Z} = \mathbb{C}^\Gamma$ for $\Gamma = \{0, 1\}$, and for any channel $\Xi \in C(\mathcal{Y}^{\otimes n}, \mathcal{Z}^{\otimes m})$, it therefore holds that

$$(\Xi \Phi^{\otimes n} \otimes \mathbb{1}_{L(\mathcal{Z})}^{\otimes m})(\rho) \in \text{PPT}(\mathcal{Z}^{\otimes m} : \mathcal{Z}^{\otimes m}) \quad (8.483)$$

for every state $\rho \in D(\mathcal{X}^{\otimes n} \otimes \mathcal{Z}^{\otimes m})$. By Proposition 6.42, one therefore has

$$F\left(2^{-m} \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m}) \text{vec}(\mathbb{1}_{\mathcal{Z}}^{\otimes m})^*, (\Xi \Phi^{\otimes n} \otimes \mathbb{1}_{L(\mathcal{Z})}^{\otimes m})(\rho)\right) \leq 2^{-m/2}. \quad (8.484)$$

For every choice of a positive real number $\alpha > 0$, it must therefore be the case that α fails to be an achievable rate for entanglement generation through Φ . Consequently, Φ has zero capacity for entanglement generation, which implies $Q(\Phi) = 0$ by Theorem 8.46. \square

The second category of channels mentioned above having zero quantum capacity are *self-complementary* channels. These are channels $\Phi \in C(\mathcal{X}, \mathcal{Y})$ such that there exists an isometry $A \in U(\mathcal{X}, \mathcal{Y} \otimes \mathcal{Y})$ such that

$$\Phi(X) = (\mathbb{1}_{L(\mathcal{Y})} \otimes \text{Tr})(AXA^*) = (\text{Tr} \otimes \mathbb{1}_{L(\mathcal{Y})})(AXA^*) \quad (8.485)$$

for every $X \in L(\mathcal{X})$. By Proposition 8.17, the coherent information of every state $\sigma \in D(\mathcal{X})$ through a self-complementary channel Φ must be zero:

$$I_c(\sigma; \Phi) = H(\Phi(\sigma)) - H(\Phi(\sigma)) = 0. \quad (8.486)$$

As every tensor power of a self-complementary channel is necessarily self-complementary, the quantum capacity theorem (Theorem 8.55) implies that self-complementary channels have zero quantum capacity. The following proposition states a more general variant of this observation.

Proposition 8.62 *Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ and $\Psi \in C(\mathcal{X}, \mathcal{Z})$ be complementary channels, for complex Euclidean spaces \mathcal{X}, \mathcal{Y} , and \mathcal{Z} , and suppose that there exists a channel $\Xi \in C(\mathcal{Z}, \mathcal{Y})$ such that $\Phi = \Xi\Psi$. It holds that Φ has zero quantum capacity: $Q(\Phi) = 0$.*

Proof Let n be a positive integer and let $\sigma \in D(\mathcal{X}^{\otimes n})$ be a state. One has

$$I_c(\sigma; \Phi^{\otimes n}) = I_c(\sigma; \Xi^{\otimes n} \Psi^{\otimes n}) \leq I_c(\sigma; \Psi^{\otimes n}) \quad (8.487)$$

by Proposition 8.15. Because Ψ is complementary to Φ , it holds that $\Psi^{\otimes n}$ is complementary to $\Phi^{\otimes n}$, and therefore

$$\begin{aligned} I_c(\sigma; \Phi^{\otimes n}) &= H(\Phi^{\otimes n}(\sigma)) - H(\Psi^{\otimes n}(\sigma)) \\ &= -I_c(\sigma; \Psi^{\otimes n}) \leq -I_c(\sigma; \Phi^{\otimes n}), \end{aligned} \quad (8.488)$$

which implies

$$I_c(\sigma; \Phi^{\otimes n}) \leq 0. \quad (8.489)$$

As this is so for every choice of n and every state $\sigma \in D(\mathcal{X}^{\otimes n})$, it follows that $Q(\Phi) = 0$ by Theorem 8.55. \square

Remark Channels of the form $\Phi \in C(\mathcal{X}, \mathcal{Y})$ for which there exists a channel $\Psi \in C(\mathcal{X}, \mathcal{Z})$ complementary to Φ , as well as a channel $\Xi \in C(\mathcal{Z}, \mathcal{Y})$ for which $\Phi = \Xi\Psi$, are known as *anti-degradable channels*.

50% erasure channels

A 50%-erasure channel is a simple type of self-complementary channel that plays a special role in the example of super-activation to be presented below. For any choice of a complex Euclidean space \mathcal{X} , the 50%-erasure channel defined with respect to \mathcal{X} is the channel $\Xi \in \mathcal{C}(\mathcal{X}, \mathbb{C} \oplus \mathcal{X})$ defined for each $X \in \mathcal{L}(\mathcal{X})$ as

$$\Xi(X) = \frac{1}{2} \begin{pmatrix} \text{Tr}(X) & 0 \\ 0 & X \end{pmatrix}. \quad (8.490)$$

Intuitively speaking, a 50%-erasure channel acts as the identity channel with probability 1/2, and otherwise its input is erased. Under the assumption that $\mathcal{X} = \mathbb{C}^\Sigma$, for Σ being a given alphabet, one may associate the complex Euclidean space $\mathbb{C} \oplus \mathcal{X}$ with $\mathbb{C}^{\{\#\} \cup \Sigma}$, for $\#$ being a special *blank symbol* that is not contained in Σ . With this interpretation, the event that the input is erased may be associated with the blank symbol $\#$ being produced, so that

$$\Xi(X) = \frac{1}{2}X + \frac{1}{2}\text{Tr}(X)E_{\#,\#} \quad (8.491)$$

for every $X \in \mathcal{L}(\mathcal{X})$.

For every choice of \mathcal{X} , the 50%-erasure channel $\Xi \in \mathcal{C}(\mathcal{X}, \mathbb{C} \oplus \mathcal{X})$ is self-complementary: one has

$$\Xi(X) = (\text{Tr} \otimes \mathbb{1})(AXA^*) = (\mathbb{1} \otimes \text{Tr})(AXA^*) \quad (8.492)$$

for $A \in \mathcal{U}(\mathcal{X}, (\mathbb{C} \oplus \mathcal{X}) \otimes (\mathbb{C} \oplus \mathcal{X}))$ being the isometry defined as

$$Ax = \frac{1}{\sqrt{2}}(0 \oplus x) \otimes (1 \oplus 0) + \frac{1}{\sqrt{2}}(1 \oplus 0) \otimes (0 \oplus x) \quad (8.493)$$

for every $x \in \mathcal{X}$. It follows that $\mathcal{Q}(\Xi) = 0$.

A theorem of Smith and Yard

The following theorem allows one to prove lower bounds on the maximum coherent information of a channel tensored with a 50%-erasure channel on a sufficiently large space. For a suitable choice of a zero-capacity channel tensored with a 50%-erasure channel, the theorem leads to a demonstration of the super-activation phenomenon.

Theorem 8.63 (Smith–Yard) *Let \mathcal{X} , \mathcal{Y} , and \mathcal{Z} be complex Euclidean spaces, let $A \in \mathcal{U}(\mathcal{X}, \mathcal{Y} \otimes \mathcal{Z})$ be an isometry, and let $\Phi \in \mathcal{C}(\mathcal{X}, \mathcal{Y})$ and $\Psi \in \mathcal{C}(\mathcal{X}, \mathcal{Z})$ be complementary channels defined as*

$$\Phi(X) = \text{Tr}_{\mathcal{Z}}(AXA^*) \quad \text{and} \quad \Psi(X) = \text{Tr}_{\mathcal{Y}}(AXA^*) \quad (8.494)$$

for every $X \in \mathcal{L}(\mathcal{X})$. Also let Σ be an alphabet, let $\eta : \Sigma \rightarrow \text{Pos}(\mathcal{X})$ be an ensemble of states, let \mathcal{W} be a complex Euclidean space satisfying

$$\dim(\mathcal{W}) \geq \sum_{a \in \Sigma} \text{rank}(\eta(a)), \quad (8.495)$$

and let $\Xi \in \mathcal{C}(\mathcal{W}, \mathbb{C} \oplus \mathcal{W})$ denote the 50%-erasure channel on \mathcal{W} . There exists a density operator $\rho \in \mathcal{D}(\mathcal{X} \otimes \mathcal{W})$ such that

$$\mathcal{I}_{\mathbb{C}}(\rho; \Phi \otimes \Xi) = \frac{1}{2}\chi(\Phi(\eta)) - \frac{1}{2}\chi(\Psi(\eta)). \quad (8.496)$$

Proof By the assumption

$$\dim(\mathcal{W}) \geq \sum_{a \in \Sigma} \text{rank}(\eta(a)), \quad (8.497)$$

one may choose a collection of vectors $\{u_a : a \in \Sigma\} \subset \mathcal{X} \otimes \mathcal{W}$ for which it holds that

$$\text{Tr}_{\mathcal{W}}(u_a u_a^*) = \eta(a) \quad (8.498)$$

for each $a \in \Sigma$, and for which

$$\{\text{Tr}_{\mathcal{X}}(u_a u_a^*) : a \in \Sigma\} \quad (8.499)$$

is an orthogonal set of operators. Let $\mathcal{V} = \mathbb{C}^\Sigma$, define a unit vector

$$u = \sum_{a \in \Sigma} e_a \otimes u_a \in \mathcal{V} \otimes \mathcal{X} \otimes \mathcal{W}, \quad (8.500)$$

and let $\rho = \text{Tr}_{\mathcal{V}}(uu^*)$. One may observe that, by virtue of the fact that (8.499) is an orthogonal set, it holds that

$$\text{Tr}_{\mathcal{W}}(uu^*) = \sum_{a \in \Sigma} E_{a,a} \otimes \eta(a). \quad (8.501)$$

For the unit vector $v \in \mathcal{V} \otimes \mathcal{Y} \otimes \mathcal{Z} \otimes \mathcal{W}$ defined as $v = (\mathbb{1}_{\mathcal{V}} \otimes A \otimes \mathbb{1}_{\mathcal{W}})u$, it therefore holds that

$$\text{Tr}_{\mathcal{W}}(vv^*) = \sum_{a \in \Sigma} E_{a,a} \otimes A\eta(a)A^*. \quad (8.502)$$

The 50%-erasure channel Ξ has the property that

$$\mathcal{H}((\Phi \otimes \Xi)(\rho)) = \frac{1}{2}\mathcal{H}((\Phi \otimes \mathbb{1}_{\mathcal{L}(\mathcal{W})})(\rho)) + \frac{1}{2}\mathcal{H}(\Phi(\text{Tr}_{\mathcal{W}}(\rho))) + 1, \quad (8.503)$$

and likewise for the channel Ψ in place of Φ . As Ψ is complementary to Φ

and Ξ is self-complementary, it follows that

$$\begin{aligned} I_c(\rho; \Phi \otimes \Xi) &= H((\Phi \otimes \Xi)(\rho)) - H((\Psi \otimes \Xi)(\rho)) \\ &= \frac{1}{2} H((\Phi \otimes \mathbb{1}_{L(\mathcal{V})})(\rho)) - \frac{1}{2} H((\Psi \otimes \mathbb{1}_{L(\mathcal{V})})(\rho)) \\ &\quad + \frac{1}{2} H(\Phi(\text{Tr}_{\mathcal{W}}(\rho))) - \frac{1}{2} H(\Psi(\text{Tr}_{\mathcal{W}}(\rho))). \end{aligned} \quad (8.504)$$

Now, let $V, Y, Z,$ and W be registers corresponding to the spaces $\mathcal{V}, \mathcal{Y}, \mathcal{Z},$ and $\mathcal{W},$ respectively, and consider the situation in which the compound register (V, Y, Z, W) is in the pure state vv^* . It holds that

$$\begin{aligned} H((\Phi \otimes \mathbb{1}_{L(\mathcal{W})})(\rho)) &= H(Y, W) = H(V, Z), \\ H((\Psi \otimes \mathbb{1}_{L(\mathcal{W})})(\rho)) &= H(Z, W) = H(V, Y), \\ H(\Phi(\text{Tr}_{\mathcal{W}}(\rho))) &= H(Y), \\ H(\Psi(\text{Tr}_{\mathcal{W}}(\rho))) &= H(Z), \end{aligned} \quad (8.505)$$

and therefore

$$I_c(\rho; \Phi \otimes \Xi) = \frac{1}{2} I(V : Y) - \frac{1}{2} I(V : Z) = \frac{1}{2} \chi(\Phi(\eta)) - \frac{1}{2} \chi(\Psi(\eta)), \quad (8.506)$$

as required. \square

An explicit example of super-activation

An example of the super-activation phenomenon, based on Theorem 8.63, will now be described. The first step is to define a zero-capacity channel Φ as follows. Let

$$\begin{aligned} A_1 &= \begin{pmatrix} 0 & 0 & \alpha & 0 \\ 0 & 0 & 0 & 0 \\ \gamma & 0 & 0 & 0 \\ 0 & \gamma & 0 & 0 \end{pmatrix}, & A_2 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \alpha \\ -\gamma & 0 & 0 & 0 \\ 0 & \gamma & 0 & 0 \end{pmatrix}, \\ A_3 &= \begin{pmatrix} \beta & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \beta & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & A_4 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ \beta & 0 & 0 & 0 \\ 0 & 0 & 0 & \beta \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ A_5 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \beta & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\beta \end{pmatrix}, & A_6 &= \begin{pmatrix} 0 & \beta & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \beta & 0 \end{pmatrix}, \end{aligned} \quad (8.507)$$

where

$$\alpha = \sqrt{\sqrt{2} - 1}, \quad \beta = \sqrt{1 - \frac{1}{\sqrt{2}}}, \quad \text{and} \quad \gamma = \sqrt{\frac{1}{\sqrt{2}} - \frac{1}{2}}, \quad (8.508)$$

and define $\Phi \in C(\mathbb{C}^4)$ as

$$\Phi(X) = \sum_{k=1}^6 A_k X A_k^* \quad (8.509)$$

for every $X \in L(\mathbb{C}^4)$.

The fact that Φ is a zero-capacity channel follows from the fact that the Choi representation of Φ is a PPT operator. One way to verify this claim is to check that

$$(T \otimes \mathbb{1}_{L(\mathbb{C}^4)})(J(\Phi)) = J(\Theta) \quad (8.510)$$

for $\Theta \in C(\mathbb{C}^4)$ being the channel defined as

$$\Theta(X) = \sum_{k=1}^6 B_k X B_k^* \quad (8.511)$$

for every $X \in L(\mathbb{C}^4)$, where

$$\begin{aligned} B_1 &= \begin{pmatrix} 0 & 0 & \alpha & 0 \\ 0 & 0 & 0 & 0 \\ \gamma & 0 & 0 & 0 \\ 0 & \gamma & 0 & 0 \end{pmatrix}, & B_2 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \alpha \\ \gamma & 0 & 0 & 0 \\ 0 & -\gamma & 0 & 0 \end{pmatrix}, \\ B_3 &= \begin{pmatrix} \beta & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \beta & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & B_4 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ \beta & 0 & 0 & 0 \\ 0 & 0 & 0 & -\beta \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ B_5 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \beta & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \beta \end{pmatrix}, & B_6 &= \begin{pmatrix} 0 & \beta & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \beta & 0 \end{pmatrix}. \end{aligned} \quad (8.512)$$

It therefore follows from Proposition 8.61 that Φ has zero quantum capacity.

A channel complementary to Φ is given by $\Psi \in C(\mathbb{C}^4, \mathbb{C}^6)$ defined as

$$\Psi(X) = \sum_{k=1}^4 C_k X C_k^* \quad (8.513)$$

for every $X \in L(\mathbb{C}^4)$, where

$$C_1 = \begin{pmatrix} 0 & 0 & \alpha & 0 \\ 0 & 0 & 0 & 0 \\ \beta & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & \beta & 0 & 0 \end{pmatrix}, \quad C_2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \alpha \\ 0 & 0 & 0 & 0 \\ \beta & 0 & 0 & 0 \\ 0 & \beta & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (8.514)$$

$$C_3 = \begin{pmatrix} \gamma & 0 & 0 & 0 \\ -\gamma & 0 & 0 & 0 \\ 0 & 0 & \beta & 0 \\ 0 & 0 & 0 & \beta \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad C_4 = \begin{pmatrix} 0 & \gamma & 0 & 0 \\ 0 & \gamma & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\beta \\ 0 & 0 & \beta & 0 \end{pmatrix}.$$

Finally, define density operators

$$\sigma_0 = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 0 \\ 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad \sigma_1 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & 0 \\ 0 & 0 & 0 & \frac{1}{2} \end{pmatrix}, \quad (8.515)$$

and define an ensemble $\eta : \{0, 1\} \rightarrow \text{Pos}(\mathbb{C}^4)$ as

$$\eta(0) = \frac{1}{2}\sigma_0 \quad \text{and} \quad \eta(1) = \frac{1}{2}\sigma_1. \quad (8.516)$$

It holds that

$$\Phi(\sigma_0) = \begin{pmatrix} \frac{2-\sqrt{2}}{2} & 0 & 0 & 0 \\ 0 & \frac{2-\sqrt{2}}{2} & 0 & 0 \\ 0 & 0 & \frac{\sqrt{2}-1}{2} & 0 \\ 0 & 0 & 0 & \frac{\sqrt{2}-1}{2} \end{pmatrix} \quad (8.517)$$

and

$$\Phi(\sigma_1) = \begin{pmatrix} \frac{\sqrt{2}-1}{2} & 0 & 0 & 0 \\ 0 & \frac{\sqrt{2}-1}{2} & 0 & 0 \\ 0 & 0 & \frac{2-\sqrt{2}}{2} & 0 \\ 0 & 0 & 0 & \frac{2-\sqrt{2}}{2} \end{pmatrix}, \quad (8.518)$$

while

$$\Psi(\sigma_0) = \Psi(\sigma_1) = \begin{pmatrix} \frac{\sqrt{2}-1}{2} & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{\sqrt{2}-1}{2} & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{2-\sqrt{2}}{4} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{2-\sqrt{2}}{4} & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{2-\sqrt{2}}{4} & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{2-\sqrt{2}}{4} \end{pmatrix}. \quad (8.519)$$

One therefore has that

$$\chi(\Phi(\eta)) = H\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right) - H\left(\frac{2-\sqrt{2}}{2}, \frac{2-\sqrt{2}}{2}, \frac{\sqrt{2}-1}{2}, \frac{\sqrt{2}-1}{2}\right) > \frac{1}{50}, \quad (8.520)$$

while $\chi(\Psi(\eta)) = 0$. By Theorem 8.63, there must exist a density operator $\rho \in D(\mathbb{C}^4 \otimes \mathbb{C}^4)$ such that

$$I_c(\rho; \Phi \otimes \Xi) > \frac{1}{100}, \quad (8.521)$$

for $\Xi \in C(\mathbb{C}^4, \mathbb{C} \oplus \mathbb{C}^4)$ being a 50%-erasure channel. One therefore has that $Q(\Phi) = Q(\Xi) = 0$, while $Q(\Phi \otimes \Xi) > 0$.

The need for a regularization in the quantum capacity theorem

The super-activation example described above illustrates that the maximum coherent information is not additive; one has

$$I_c(\Phi \otimes \Xi) > I_c(\Phi) + I_c(\Xi) \quad (8.522)$$

for the channels Φ and Ξ specified in that example. As these channels are different, it does not follow immediately that a strict inequality of the form

$$I_c(\Psi^{\otimes n}) > nI_c(\Psi) \quad (8.523)$$

holds for any choice of a channel Ψ and a positive integer n . It is possible, however, to conclude that such an inequality does hold (for $n = 2$) using a direct sum construction along similar lines to the one used in the context of the Holevo capacity and minimum output entropy. The following three propositions that concern direct sums of channels will be used to reach this conclusion.

Proposition 8.64 *Let $\mathcal{X}_0, \mathcal{X}_1, \mathcal{Y}_0, \mathcal{Y}_1, \mathcal{Z}_0$, and \mathcal{Z}_1 be complex Euclidean spaces, and let $\Phi_0 \in C(\mathcal{X}_0, \mathcal{Y}_0)$, $\Phi_1 \in C(\mathcal{X}_1, \mathcal{Y}_1)$, $\Psi_0 \in C(\mathcal{X}_0, \mathcal{Z}_0)$, and $\Psi_1 \in C(\mathcal{X}_1, \mathcal{Z}_1)$ be channels such that Ψ_0 is complementary to Φ_0 and Ψ_1 is complementary to Φ_1 . The channel $\Psi_0 \oplus \Psi_1$ is complementary to $\Phi_0 \oplus \Phi_1$.*

Proof Let $A_0 \in U(\mathcal{X}_0, \mathcal{Y}_0 \otimes \mathcal{Z}_0)$ and $A_1 \in U(\mathcal{X}_1, \mathcal{Y}_1 \otimes \mathcal{Z}_1)$ be isometries such that the following equations hold for all $X_0 \in L(\mathcal{X}_0)$ and $X_1 \in L(\mathcal{X}_1)$:

$$\begin{aligned}\Phi_0(X_0) &= \text{Tr}_{\mathcal{Z}_0}(A_0 X_0 A_0^*), & \Psi_0(X_0) &= \text{Tr}_{\mathcal{Y}_0}(A_0 X_0 A_0^*), \\ \Phi_1(X_1) &= \text{Tr}_{\mathcal{Z}_1}(A_1 X_1 A_1^*), & \Psi_1(X_1) &= \text{Tr}_{\mathcal{Y}_1}(A_1 X_1 A_1^*).\end{aligned}\quad (8.524)$$

Let $W \in U((\mathcal{Y}_0 \otimes \mathcal{Z}_0) \oplus (\mathcal{Y}_1 \otimes \mathcal{Z}_1), (\mathcal{Y}_0 \oplus \mathcal{Y}_1) \otimes (\mathcal{Z}_0 \oplus \mathcal{Z}_1))$ be the isometry defined by the equation

$$\begin{aligned}W((y_0 \otimes z_0) \oplus (y_1 \otimes z_1)) \\ = (y_0 \oplus 0) \otimes (z_0 \oplus 0) + (0 \oplus y_1) \otimes (0 \oplus z_1)\end{aligned}\quad (8.525)$$

for every $y_0 \in \mathcal{Y}_0$, $y_1 \in \mathcal{Y}_1$, $z_0 \in \mathcal{Z}_0$, and $z_1 \in \mathcal{Z}_1$. The equations

$$\begin{aligned}(\Phi_0 \oplus \Phi_1)(X) &= \text{Tr}_{\mathcal{Z}_0 \oplus \mathcal{Z}_1} \left(W \begin{pmatrix} A_0 & 0 \\ 0 & A_1 \end{pmatrix} X \begin{pmatrix} A_0^* & 0 \\ 0 & A_1^* \end{pmatrix} W^* \right) \\ (\Psi_0 \oplus \Psi_1)(X) &= \text{Tr}_{\mathcal{Y}_0 \oplus \mathcal{Y}_1} \left(W \begin{pmatrix} A_0 & 0 \\ 0 & A_1 \end{pmatrix} X \begin{pmatrix} A_0^* & 0 \\ 0 & A_1^* \end{pmatrix} W^* \right)\end{aligned}\quad (8.526)$$

hold for all $X \in L(\mathcal{X}_0 \oplus \mathcal{X}_1)$, which implies that $\Psi_0 \oplus \Psi_1$ is complementary to $\Phi_0 \oplus \Phi_1$, as required. \square

Proposition 8.65 *Let $\Phi_0 \in C(\mathcal{X}_0, \mathcal{Y}_0)$ and $\Phi_1 \in C(\mathcal{X}_1, \mathcal{Y}_1)$ be channels, for $\mathcal{X}_0, \mathcal{X}_1, \mathcal{Y}_0$, and \mathcal{Y}_1 being complex Euclidean spaces, and let $\sigma \in D(\mathcal{X}_0 \oplus \mathcal{X}_1)$ be an arbitrary state, written as*

$$\sigma = \begin{pmatrix} \lambda \sigma_0 & X \\ X^* & (1 - \lambda) \sigma_1 \end{pmatrix}\quad (8.527)$$

for $\lambda \in [0, 1]$, $\sigma_0 \in D(\mathcal{X}_0)$, $\sigma_1 \in D(\mathcal{X}_1)$, and $X \in L(\mathcal{X}_1, \mathcal{X}_0)$. It holds that

$$I_c(\sigma; \Phi_0 \oplus \Phi_1) = \lambda I_c(\sigma_0; \Phi_0) + (1 - \lambda) I_c(\sigma_1; \Phi_1).\quad (8.528)$$

Proof Observe first that

$$\begin{aligned}H((\Phi_0 \oplus \Phi_1)(\sigma)) &= H \begin{pmatrix} \lambda \Phi_0(\sigma_0) & 0 \\ 0 & (1 - \lambda) \Phi_1(\sigma_1) \end{pmatrix} \\ &= \lambda H(\Phi_0(\sigma_0)) + (1 - \lambda) H(\Phi_1(\sigma_1)) + H(\lambda, 1 - \lambda).\end{aligned}\quad (8.529)$$

Assuming that \mathcal{Z}_0 and \mathcal{Z}_1 are complex Euclidean spaces and $\Psi_0 \in C(\mathcal{X}_0, \mathcal{Z}_0)$ and $\Psi_1 \in C(\mathcal{X}_1, \mathcal{Z}_1)$ are channels complementary to Φ_0 and Φ_1 , respectively, one has that

$$\begin{aligned}H((\Psi_0 \oplus \Psi_1)(\sigma)) \\ = \lambda H(\Psi_0(\sigma_0)) + (1 - \lambda) H(\Psi_1(\sigma_1)) + H(\lambda, 1 - \lambda)\end{aligned}\quad (8.530)$$

by a similar calculation to (8.529). As $\Psi_0 \oplus \Psi_1$ is complementary to $\Phi_0 \oplus \Phi_1$, as established in Proposition 8.64, it follows that

$$\begin{aligned}I_c(\sigma; \Phi_0 \oplus \Phi_1) &= H((\Phi_0 \oplus \Phi_1)(\sigma)) - H((\Psi_0 \oplus \Psi_1)(\sigma)) \\ &= \lambda(H(\Phi_0(\sigma_0)) - H(\Psi_0(\sigma_0))) \\ &\quad + (1 - \lambda)(H(\Phi_1(\sigma_1)) - H(\Psi_1(\sigma_1))) \\ &= \lambda I_c(\sigma_0; \Phi_0) + (1 - \lambda) I_c(\sigma_1; \Phi_1)\end{aligned}\quad (8.531)$$

as required. \square

Proposition 8.66 *Let $\mathcal{X}_0, \mathcal{X}_1, \mathcal{Y}_0$, and \mathcal{Y}_1 be complex Euclidean spaces and let $\Phi_0 \in C(\mathcal{X}_0, \mathcal{Y}_0)$ and $\Phi_1 \in C(\mathcal{X}_1, \mathcal{Y}_1)$ be channels. It holds that*

$$I_c((\Phi_0 \oplus \Phi_1) \otimes (\Phi_0 \oplus \Phi_1)) \geq I_c(\Phi_0 \otimes \Phi_1).\quad (8.532)$$

Proof Define an isometry $W \in U(\mathcal{X}_0 \otimes \mathcal{X}_1, (\mathcal{X}_0 \oplus \mathcal{X}_1) \otimes (\mathcal{X}_0 \oplus \mathcal{X}_1))$ by the equation

$$W(x_0 \otimes x_1) = (x_0 \oplus 0) \otimes (0 \oplus x_1)\quad (8.533)$$

holding for all $x_0 \in \mathcal{X}_0$ and $x_1 \in \mathcal{X}_1$, and along similar lines, define an isometry $V \in U(\mathcal{Y}_0 \otimes \mathcal{Y}_1, (\mathcal{Y}_0 \oplus \mathcal{Y}_1) \otimes (\mathcal{Y}_0 \oplus \mathcal{Y}_1))$ by the equation

$$V(y_0 \otimes y_1) = (y_0 \oplus 0) \otimes (0 \oplus y_1)\quad (8.534)$$

for all $y_0 \in \mathcal{Y}_0$ and $y_1 \in \mathcal{Y}_1$. One has that

$$\begin{aligned}((\Phi_0 \oplus \Phi_1) \otimes (\Phi_0 \oplus \Phi_1))(W(X_0 \otimes X_1)W^*) \\ = \begin{pmatrix} \Phi_0(X_0) & 0 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 0 & \Phi_1(X_1) \end{pmatrix} \\ = V(\Phi_0(X_0) \otimes \Phi_1(X_1))V^*\end{aligned}\quad (8.535)$$

for all $X_0 \in L(\mathcal{X}_0)$ and $X_1 \in L(\mathcal{X}_1)$.

For every choice of a density operator $\sigma \in D(\mathcal{X}_0 \otimes \mathcal{X}_1)$, it follows that

$$I_c(W\sigma W^*; (\Phi_0 \oplus \Phi_1) \otimes (\Phi_0 \oplus \Phi_1)) = I_c(\sigma; \Phi_0 \otimes \Phi_1),\quad (8.536)$$

which implies the proposition. \square

Finally, consider the channel $\Psi = \Phi \oplus \Xi$, for Φ and Ξ as in the example of super-activation described above. By Proposition 8.65, one may conclude that $I_c(\Phi \oplus \Xi) = 0$, while Proposition 8.66 implies

$$I_c((\Phi \oplus \Xi) \otimes (\Phi \oplus \Xi)) \geq I_c(\Phi \otimes \Xi) > 0.\quad (8.537)$$

It therefore holds that the channel $\Psi = \Phi \oplus \Xi$ satisfies the strict inequality (8.523) for $n = 2$.

As a consequence of this fact, one has that the quantum capacity and maximum coherent information differ for some channels. In this sense, the regularization in the quantum capacity theorem (Theorem 8.55) is similar to the one in the Holevo–Schumacher–Westmoreland theorem (Theorem 8.27) in that it cannot generally be removed.

8.4 Exercises

Exercise 8.1 Let $\Phi_0 \in C(\mathcal{X}_0, \mathcal{Y}_0)$ and $\Phi_1 \in C(\mathcal{X}_1, \mathcal{Y}_1)$ be channels, for an arbitrary choice of complex Euclidean spaces $\mathcal{X}_0, \mathcal{X}_1, \mathcal{Y}_0$, and \mathcal{Y}_1 .

(a) Prove that

$$I_c(\Phi_0 \oplus \Phi_1) = \max\{I_c(\Phi_0), I_c(\Phi_1)\}. \quad (8.538)$$

(b) Prove that

$$\chi(\Phi_0 \oplus \Phi_1) = \max_{\lambda \in [0,1]} (\lambda\chi(\Phi_0) + (1-\lambda)\chi(\Phi_1) + H(\lambda, 1-\lambda)). \quad (8.539)$$

Exercise 8.2 Let $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$, and \mathcal{W} be complex Euclidean spaces, let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ and $\Psi \in C(\mathcal{Z}, \mathcal{W})$ be channels, and assume that Φ is an entanglement breaking channel (q.v. Exercise 6.1). Prove that the following identities hold:

$$(a) \quad H_{\min}(\Phi \otimes \Psi) = H_{\min}(\Phi) + H_{\min}(\Psi).$$

$$(b) \quad \chi(\Phi \otimes \Psi) = \chi(\Phi) + \chi(\Psi).$$

$$(c) \quad I_c(\Phi \otimes \Psi) = I_c(\Psi).$$

Exercise 8.3 Let $\Phi \in C(\mathcal{X}, \mathcal{Y})$ be a channel, for complex Euclidean spaces \mathcal{X} and \mathcal{Y} . It is said that Φ is *degradable* if there exists a complex Euclidean space \mathcal{Z} and a channel $\Psi \in C(\mathcal{Y}, \mathcal{Z})$ such that $\Psi\Phi$ is complementary to Φ .

(a) Prove that, for any choice of a degradable channel $\Phi \in C(\mathcal{X}, \mathcal{Y})$, states $\sigma_0, \sigma_1 \in D(\mathcal{X})$, and a real number $\lambda \in [0, 1]$, the following inequality holds:

$$I_c(\lambda\sigma_0 + (1-\lambda)\sigma_1; \Phi) \geq \lambda I_c(\sigma_0; \Phi) + (1-\lambda)I_c(\sigma_1; \Phi). \quad (8.540)$$

(Equivalently, the function $\sigma \mapsto I_c(\sigma; \Phi)$ defined on $D(\mathcal{X})$ is concave.)

(b) Prove that, for any choice of complex Euclidean spaces $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$, and \mathcal{W} and degradable channels $\Phi \in C(\mathcal{X}, \mathcal{Y})$ and $\Psi \in C(\mathcal{Z}, \mathcal{W})$, it holds that

$$I_c(\Phi \otimes \Psi) = I_c(\Phi) + I_c(\Psi). \quad (8.541)$$

Exercise 8.4 Let \mathcal{X} be a complex Euclidean space, let $\lambda \in [0, 1]$, and define a channel $\Xi \in C(\mathcal{X}, C \oplus \mathcal{X})$ as

$$\Xi(X) = \begin{pmatrix} \lambda \text{Tr}(X) & 0 \\ 0 & (1-\lambda)X \end{pmatrix} \quad (8.542)$$

for all $X \in L(\mathcal{X})$.

(a) Give a closed-form expression for the coherent information $I_c(\sigma; \Xi)$ of an arbitrary state $\sigma \in D(\mathcal{X})$ through Ξ .

(b) Give a closed-form expression for the entanglement-assisted classical capacity $C_E(\Xi)$ of Ξ .

(c) Give a closed-form expression for the quantum capacity $Q(\Xi)$ of Ξ .

The closed-form expressions for parts (b) and (c) should be functions of λ and $n = \dim(\mathcal{X})$ alone.

Exercise 8.5 Let n be a positive integer, let $\mathcal{X} = \mathbb{C}^{\mathbb{Z}_n}$, and let

$$\{W_{a,b} : a, b \in \mathbb{Z}_n\} \quad (8.543)$$

denote the set of discrete Weyl operators acting on \mathcal{X} (q.v. Section 4.1.2 of Chapter 4). Also let $p \in \mathcal{P}(\mathbb{Z}_n)$ be a probability vector, and define a channel $\Phi \in C(\mathcal{X})$ as

$$\Phi(X) = \sum_{a \in \mathbb{Z}_n} p(a) W_{0,a} X W_{0,a}^* \quad (8.544)$$

for all $X \in L(\mathcal{X})$. Prove that

$$I_c(\Phi) = \log(n) - H(p). \quad (8.545)$$

Exercise 8.6 For every positive integer n and every real number $\varepsilon \in [0, 1]$, define a channel $\Phi_{n,\varepsilon} \in C(\mathbb{C}^n)$ as

$$\Phi_{n,\varepsilon} = \varepsilon \mathbb{1}_n + (1-\varepsilon)\Omega_n, \quad (8.546)$$

where $\mathbb{1}_n \in C(\mathbb{C}^n)$ and $\Omega_n \in C(\mathbb{C}^n)$ denote the identity and completely depolarizing channels defined with respect to the space \mathbb{C}^n .

(a) Prove that, for every choice of a positive real number K , there exists a choice of n and ε for which

$$C_E(\Phi_{n,\varepsilon}) \geq K\chi(\Phi_{n,\varepsilon}) > 0. \quad (8.547)$$

(b) Prove that the fact established by a correct answer to part (a) remains true when $\chi(\Phi_{n,\varepsilon})$ is replaced by $C(\Phi_{n,\varepsilon})$.

8.5 Bibliographic remarks

The study of quantum channel capacities is, perhaps obviously, motivated in large part by Shannon's channel coding theorem (Shannon, 1948), and the goal of obtaining analogous statements for quantum channels. It was, however, realized early in the study of quantum information theory that there would not be a single capacity of a quantum channel, but rather several inequivalent but nevertheless fundamentally interesting capacities. The 1998 survey of Bennett and Shor (1998) provides a summary of what was known about channel capacities at a relatively early point in their study.

Holevo (1998) and Schumacher and Westmoreland (1997) independently proved the Holevo–Schumacher–Westmoreland theorem (Theorem 8.27), in both cases building on Hausladen, Jozsa, Schumacher, Westmoreland, and Wootters (1996). The definition of what is now called the Holevo capacity (or the *Holevo information* of a channel) originates with the work of Holevo and Schumacher and Westmoreland. Lemma 8.25 was proved by Hayashi and Nagaoka (2003), who used it in the analysis of generalizations of the Holevo–Schumacher–Westmoreland theorem.

The entanglement-assisted classical capacity theorem (Theorem 8.41) was proved by Bennett, Shor, Smolin, and Thapliyal (1999a). The proof of this theorem presented in this chapter is due to Holevo (2002). Lemma 8.38 is due to Adami and Cerf (1997).

Tasks that involve quantum information transmission through quantum channels, along with fundamental definitions connected with such tasks, were investigated by Schumacher (1996), Schumacher and Nielsen (1996), Adami and Cerf (1997), and Barnum, Nielsen, and Schumacher (1998), among others. The entanglement generation capacity of a channel was defined by Devetak (2005), and Theorems 8.45 and 8.46 follow from results proved by Barnum, Knill, and Nielsen (2000).

The coherent information of a state through a channel was defined by Schumacher and Nielsen (1996). Lloyd (1997) recognized the fundamental connection between the maximum coherent information of a channel and its quantum capacity, and provided a heuristic argument in support of the quantum capacity theorem (Theorem 8.55). The first rigorous proof of the quantum capacity theorem to be published was due to Devetak (2005). Shor reported a different proof of this theorem prior to Devetak's proof, although it was not published. A proof appearing in a subsequent paper of Hayden, Shor, and Winter (2008b) resembles Shor's original proof.

The proof of the quantum capacity theorem presented in this chapter is due to Hayden, M. Horodecki, Winter, and Yard (2008a), incorporating some

simplifying ideas due to Klesse (2008), who independently proved the same theorem based on similar techniques. The phenomenon of decoupling (as represented by Lemma 8.49) provides a key step in this proof; this basic technique was used by Devetak (2005), and was identified explicitly by M. Horodecki, Oppenheim, and Winter (2007) and Abeyesinghe, Devetak, Hayden, and Winter (2009). Further information on decoupling can be found in the PhD thesis of Dupuis (2009).

Shor (2004) proved that the non-additivity of Holevo capacity follows from the non-additivity of minimum output entropy. In the same paper, Shor also proved the converse implication, which naturally had greater relevance prior to Hastings proof that the minimum output entropy is non-additive, along with the equivalence of these two non-additivity statements with two other statements concerning the entanglement of formation. The direct sum construction of channels and its implications to the additivity of channel capacities was investigated by Fukuda and Wolf (2007).

The fact that the coherent information is not additive in general was first proved by DiVincenzo, Shor, and Smolin (1998). Various properties of quantum erasure channels were established by Bennett, DiVincenzo, and Smolin (1997). Theorem 8.63, along with the realization that it gives an example of the super-activation phenomenon, is due to Smith and Yard (2008). The channel Φ described in the chapter giving rise to an example of super-activation, which appears in Smith and Yard's paper as well, was identified by K. Horodecki, Pankowski, M. Horodecki, and P. Horodecki (2008), as it relates to a different capacity known as the *private capacity* of a channel.