



# On Fibonacci and Lucas Numbers of the Form $cx^2$

Refik Keskin and Zafer Yosma  
Department of Mathematics  
Faculty of Science and Arts  
Sakarya University  
Sakarya  
Turkey

[rkeskin@sakarya.edu.tr](mailto:rkeskin@sakarya.edu.tr)  
[d085016004@sakarya.edu.tr](mailto:d085016004@sakarya.edu.tr)

## Abstract

In this paper, by using some congruences concerning with Fibonacci and Lucas numbers, we completely solve the Diophantine equations  $L_n = 2L_mx^2$ ,  $F_n = 2F_mx^2$ ,  $L_n = 6L_mx^2$ ,  $F_n = 3F_mx^2$ , and  $F_n = 6F_mx^2$ .

## 1 Introduction

Fibonacci and Lucas sequences are defined as follows;  $F_0 = 0$ ,  $F_1 = 1$ ,  $F_n = F_{n-1} + F_{n-2}$  for  $n \geq 2$  and  $L_0 = 2$ ,  $L_1 = 1$ ,  $L_n = L_{n-1} + L_{n-2}$  for  $n \geq 2$ , respectively.  $F_n$  is called the  $n$ -th Fibonacci number and  $L_n$  is called the  $n$ -th Lucas number. Fibonacci and Lucas numbers for negative subscripts are given by  $F_{-n} = (-1)^{n+1} F_n$  for  $n \geq 1$  and  $L_{-n} = (-1)^n L_n$  for  $n \geq 1$ . It can be seen that  $L_n = F_{n-1} + F_{n+1}$  and  $L_{n-1} + L_{n+1} = 5F_n$  for every  $n \in \mathbb{Z}$ . For more information about Fibonacci and Lucas sequences, one can consult [9], [18].

Let  $\alpha$  and  $\beta$  denote the roots of the equation  $x^2 - x - 1 = 0$ . Then  $\alpha = (1 + \sqrt{5})/2$  and  $\beta = (1 - \sqrt{5})/2$ . It can be seen that  $\alpha\beta = -1$  and  $\alpha + \beta = 1$ . Moreover it is well known and easy to show that

$$F_n = (\alpha^n - \beta^n) / \sqrt{5}$$

and

$$L_n = \alpha^n + \beta^n$$

for every  $n \in \mathbb{Z}$ .

In the following section, we will give some congruences concerning with Fibonacci and Lucas numbers. By using these congruences, we may prove many properties known before.

## 2 Preliminaries

The problem of characterizing the square Fibonacci numbers was first introduced in the book by Ogilvy [12, p. 100]. In 1963, both Moser and Carlitz [10], and Rollet [17] proposed this problem. In 1964, the square conjecture was proved by Cohn [4] and independently by Wyler [19]. Later the problem of characterizing the square Lucas numbers was solved by Cohn [6] and by Alfred [1]. Moreover in 1965, Cohn solved the Diophantine equations  $F_n = 2x^2$  and  $L_n = 2x^2$  in [6].

We give the following theorem from [5].

**Theorem 1.** *If  $F_n = x^2$ , then  $n = 1, 2, 12$ . If  $F_n = 2x^2$ , then  $n = 3, 6$ . If  $L_n = x^2$ , then  $n = 1, 3$  and if  $L_n = 2x^2$ , then  $n = 6$ .*

The proofs of the following two theorems are given in [8].

**Theorem 2.** *Let  $n \in \mathbb{N} \cup \{0\}$  and  $k, m \in \mathbb{Z}$ . Then*

$$F_{2mn+k} \equiv (-1)^{mn} F_k \pmod{F_m} \quad (1)$$

and

$$L_{2mn+k} \equiv (-1)^{mn} L_k \pmod{F_m}. \quad (2)$$

**Theorem 3.** *Let  $n \in \mathbb{N} \cup \{0\}$  and  $k, m \in \mathbb{Z}$ . Then*

$$L_{2mn+k} \equiv (-1)^{(m+1)n} L_k \pmod{L_m} \quad (3)$$

and

$$F_{2mn+k} \equiv (-1)^{(m+1)n} F_k \pmod{L_m}. \quad (4)$$

From the identity (2), it follows that  $8 \nmid L_n$  for any natural number  $n$ .

Now we give two lemmas and a corollary, which will be needed later. The proofs of the lemmas can be achieved by induction. For the proof of the corollary, one can consult [2] or [11].

**Lemma 4.**  $L_{2^k} \equiv 3 \pmod{4}$  for the all positive integers  $k$  with  $k \geq 1$ .

**Lemma 5.** If  $r \geq 3$ , then  $L_{2^r} \equiv 2 \pmod{3}$ .

**Corollary 6.** If  $k \geq 1$ , then there is no integer  $x$  such that  $x^2 \equiv -1 \pmod{L_{2^k}}$ .

The following lemma can be proved by induction.

**Lemma 7.** If  $r \geq 2$ , then  $L_{2^r} \equiv 7 \pmod{8}$ .

The proofs of the following theorems can be found in [3], [18] or [8].

**Theorem 8.** *Let  $m, n \in \mathbb{N}$  and  $m \geq 2$ . Then  $L_m|L_n$  if and only if  $m|n$  and  $\frac{n}{m}$  is an odd integer.*

**Theorem 9.** *Let  $m, n \in \mathbb{N}$  and  $m \geq 3$ . Then  $F_m|F_n$  if and only if  $m|n$ .*

**Theorem 10.** *Let  $m, n \in \mathbb{N}$  and  $m \geq 2$ . Then  $L_m|F_n$  if and only if  $m|n$  and  $\frac{n}{m}$  is an even integer.*

Also we give some identities about Fibonacci and Lucas numbers which will be needed in the sequel:

$$L_{2n} = L_n^2 - 2(-1)^n \quad (5)$$

$$L_{3n} = L_n(L_n^2 - 3(-1)^n) \quad (6)$$

$$F_{2n} = F_n L_n \quad (7)$$

$$F_{3n} = F_n(5F_n^2 + 3(-1)^n) \quad (8)$$

$$L_n^2 - 5F_n^2 = 4(-1)^n \quad (9)$$

$$2|F_n \Leftrightarrow 2|L_n \Leftrightarrow 3|n \quad (10)$$

$$(F_n, L_n) = 1 \text{ or } (F_n, L_n) = 2 \quad (11)$$

Let  $\left(\frac{a}{p}\right)$  represent the Legendre symbol. Then we have

$$\left(\frac{2}{p}\right) = 1 \text{ if and only if } p \equiv \pm 1 \pmod{8} \quad (12)$$

and

$$\left(\frac{-2}{p}\right) = 1 \text{ if and only if } p \equiv 1, 3 \pmod{8}. \quad (13)$$

For the proof of (12) and (13), one can consult [2] or [11].

### 3 Main Theorems

Many authors investigated Fibonacci and Lucas numbers of the form  $cx^2$ . In [5], Cohn solved  $F_n = cx^2$  and  $L_n = cx^2$  for  $c = 1, 2$ . In [14], Robbins considered Fibonacci numbers of the form  $px^2$ . Robbins solved the equation  $F_n = px^2$  for all  $p$  such that  $p \equiv 3 \pmod{4}$  or  $p < 10000$ . Later, in [15] Robbins considered Fibonacci numbers of the form  $cx^2$ . The author obtained all solutions of  $F_n = cx^2$  for composite values of  $c \leq 1000$ . After that, in [16], the same author solved  $L_n = px^2$ , where  $p$  is an odd prime and  $p < 1000$ . Moreover,

in [20], Zhou dealt with Lucas numbers of the form  $L_n = px^2$ , where  $p$  is a prime number, and he gave solutions for  $1000 < p < 60000$ . In this section, we consider the equations  $L_n = 2L_mx^2$ ,  $F_n = 2F_mx^2$ ,  $L_n = 6L_mx^2$ ,  $F_n = 3F_mx^2$ , and  $F_n = 6F_mx^2$ .

In [13], Ribenboim considers square-classes of Fibonacci numbers.  $F_m, F_n$  are in the same square-class if there exist non-zero integers  $x, y$  such that  $F_mx^2 = F_ny^2$ ; or equivalently, when  $F_mF_n$  is a square. In a similar way, he considers square-classes of Lucas numbers. A square-class will be called trivial if it consists of only one number. Ribenboim showed that the square-class of  $L_m$  is trivial when  $m \neq 0, 1, 3$ , and  $6$ . He also showed that the square-class of  $F_m$  is trivial when  $m \neq 1, 2, 3, 6, 12$ . Now, we can give following two theorems, which can be obtained from Proposition 1 and Proposition 2 given in [13].

From now on, we will assume that  $n$  and  $m$  are positive integers.

**Theorem 11.** *Let  $m > 3$  be an integer and  $F_n = F_mx^2$  for some  $x \in \mathbb{Z}$ . Then  $n = m$ .*

**Theorem 12.** *Let  $m \geq 2$  be an integer and  $L_n = L_mx^2$  for some  $x \in \mathbb{Z}$ . Then  $n = m$ .*

The proofs of the following two theorems can be obtained from Theorem 6 and Theorem 12 given in [7], but we will give a different proof.

**Theorem 13.** *There is no integer  $x$  such that  $L_n = 2L_mx^2$  for  $m > 1$ .*

*Proof.* Assume that  $L_n = 2L_mx^2$ . Then  $L_m|L_n$  and therefore  $n = mk$  for some odd natural number  $k$  by Theorem 8. Firstly assume that  $m$  is an odd integer. Since  $2|L_n$ , we get  $3|n$  by (10). Thus we see that  $3 \nmid m$ . For if  $3|m$ , then  $L_3|L_m$ , i.e.,  $4|L_m$  by Theorem 8. This implies that  $8|L_n$ , which is impossible. Since  $3 \nmid m$ , it follows that  $3|k$ . That is,  $k = 3t$  for some odd positive integer  $t$ . Thus  $n = mk = 3mt$  and  $mt$  is an odd integer. Therefore, since  $3|n$ , it follows that  $L_3|L_n$ , i.e.,  $4|2L_mx^2$  by Theorem 8. Since  $3 \nmid m$ ,  $L_m$  is an odd integer. Therefore  $2|x^2$ , i.e.,  $x$  is an even integer. This implies that  $8|L_n$ , which is impossible.

Now assume that  $m$  is an even integer. If  $x$  is an even integer, then we see that  $8|L_n$ , which is impossible. Therefore  $x$  is an odd integer. Assume that  $3|m$ . Then  $L_m$  is an even integer. Therefore  $L_3|L_n$  by Theorem 8. It follows that  $n = 3b$  for some odd integer  $b$  by Theorem 8. That is,  $n$  is an odd integer. But this is impossible. Because since  $m$  is an even integer,  $n$  is also an even integer. Assume that  $3 \nmid m$ . Then since  $n = mk$  and  $3|n$ , we get  $3|k$ , i.e.,  $k = 3t$  for some odd integer  $t$ . Since  $t$  is an odd integer,  $t = 4q \pm 1$  for some nonnegative integer  $q$ . Thus  $n = mk = 3m(4q \pm 1) = 2 \cdot 6mq \pm 3m$ . Then

$$L_n = L_{2 \cdot 6mq \pm 3m} \equiv L_{\pm 3m} \pmod{F_6}$$

and therefore

$$2L_mx^2 \equiv L_{3m} \pmod{8}$$

by (2). Since  $x^2 \equiv 1 \pmod{8}$  and  $m$  is even integer, we get

$$2L_m \equiv L_m(L_m^2 - 3) \pmod{8}$$

by (6). Moreover, since  $3 \nmid m$ ,  $L_m$  is odd integer. Therefore we get

$$2 \equiv L_m^2 - 3 \pmod{8}.$$

Thus

$$2 \equiv -2 \pmod{8},$$

which is impossible. This completes the proof.  $\square$

In [5], it is shown that, for  $m = 1, 2$ , the equation  $F_n = 2F_m x^2 = 2x^2$  has solution only for  $n = 3, 6$ . More generally, we can give the following theorem.

**Theorem 14.** *If  $F_n = 2F_m x^2$  and  $m \geq 3$ , then  $m = 3$ ,  $x^2 = 36$ , and  $n = 12$  or  $m = 6$ ,  $x^2 = 9$ , and  $n = 12$ .*

*Proof.* If  $m = 3$ , then  $F_n = 2F_3 x^2 = (2x)^2$ . Thus it can be seen that  $n = 12$ ,  $x^2 = 36$  by Theorem 1. Assume that  $m > 3$  and  $F_n = 2F_m x^2$ . Then  $F_m | F_n$  and therefore  $n = mk$  for some natural number  $k$  by Theorem 9.

Firstly, assume that  $k$  is an even integer. Then  $k = 2t$  for some integer  $t$ . Therefore  $n = mk = 2mt$ . Thus

$$F_n = F_{2mt} = F_{mt} L_{mt} = 2F_m x^2$$

by (7). This shows that  $(F_{mt}/F_m) L_{mt} = 2x^2$ . It can be easily seen that if  $(F_{mt}/F_m, L_{mt}) = d$ , then  $d = 1$  or  $d = 2$  by (11). Thus we have the following equations:

$$\frac{F_{mt}}{F_m} = u^2, \quad L_{mt} = 2v^2, \quad (14)$$

$$\frac{F_{mt}}{F_m} = 2u^2, \quad L_{mt} = v^2, \quad (15)$$

$$\frac{F_{mt}}{F_m} = 2u^2, \quad L_{mt} = (2v)^2, \quad (16)$$

and

$$\frac{F_{mt}}{F_m} = (2u)^2, \quad L_{mt} = 2v^2. \quad (17)$$

Assume that (14) is satisfied. Then  $mt = m$ , i.e.,  $t = 1$  by Theorem 11. Therefore  $L_m = 2v^2$  and this implies that  $m = 6$  by Theorem 1. Thus we get  $m = 6$ ,  $x^2 = 9$ , and  $n = 12$ . By using Theorem 1 and Theorem 11, it can be seen that the other three cases are impossible.

Secondly, assume that  $k$  is an odd integer. Suppose that  $m$  is an even integer, i.e.,  $m = 2r$  for some natural number  $r$ . Then we can write  $n = mk = 2kr$ . Thus

$$F_n = F_{2kr} = F_{kr} L_{kr} = 2F_r L_r x^2$$

by (7). This shows that  $(F_{kr}/F_r) (L_{kr}/L_r) = 2x^2$ . A similar argument shows that the equation  $(F_{kr}/F_r) (L_{kr}/L_r) = 2x^2$  has no solution. Now assume that  $m$  is an odd integer. Firstly, suppose that  $3 \nmid k$ . Since  $k$  is an odd integer, we can write  $k = 6q \pm 1$  for some nonnegative integer  $q$ . Therefore  $n = mk = m(6q \pm 1) = 2 \cdot 3mq \pm m$ . Thus we get

$$F_n = F_{2 \cdot 3mq \pm m} \equiv F_{\pm m} \pmod{L_3},$$

i.e.,

$$F_n \equiv F_m \pmod{4}$$

by (4). Since  $F_n$  is even integer,  $F_m$  is also an even integer. Thus  $3|m$ , and therefore  $m = 3a$  for some integer  $a$  by (10). On the other hand, since  $F_m$  is even integer,  $4|F_n$ , and thus  $6|n$  by Theorem 9. Since  $n = mk = 3ak$ , we get  $6|3ak$ , i.e.,  $2|ak$ . Moreover, since  $k$  is odd integer, it is seen that  $2|a$ . This implies that  $2|m$ , which is impossible. Because  $m$  is an odd integer. Assume that  $3|k$ . Then  $k = 3s$  for some odd integer  $s$ . Therefore  $n = mk = 3ms$ . Thus since  $ms$  is odd integer, we get

$$F_n = F_{3ms} = F_{ms}(5F_{ms}^2 - 3) = 2F_mx^2$$

by (8). This shows that  $(F_{ms}/F_m)(5F_{ms}^2 - 3) = 2x^2$ . It can be easily seen that if  $d = (F_{ms}/F_m, 5F_{ms}^2 - 3)$ , then  $d = 1$  or  $d = 3$ . Assume that  $d = 3$ . Then  $3|F_{ms}$ , and thus  $4|ms$  by Theorem 9. But this is impossible, since  $ms$  is odd integer. Therefore  $d = 1$ . Then we get

$$\frac{F_{ms}}{F_m} = u^2, \quad 5F_{ms}^2 - 3 = 2v^2 \quad (18)$$

or

$$\frac{F_{ms}}{F_m} = 2u^2, \quad 5F_{ms}^2 - 3 = v^2 \quad (19)$$

for some integers  $u$  and  $v$ . Assume that (18) is satisfied. Then  $ms = m$ , i.e.,  $s = 1$  by Theorem 11. Therefore  $5F_m^2 - 3 = 2v^2$  and this shows that  $2v^2 = 5F_m^2 - 3 = L_m^2 + 1 = L_{2m} - 1$  by (9) and (5). This implies that  $L_{2m} = 2v^2 + 1$ . Since  $L_{2m} = 2v^2 + 1$ , we get  $3 \nmid m$ . Thus we can write  $m = 6q \pm 1 = 3 \cdot 2^{r+1}b \pm 1$ , where  $q = 2^r b$  for some odd integer  $b$  with  $r \geq 0$ . This shows that

$$L_{2m} = L_{2 \cdot 2^{r+1}3b \pm 2} \equiv -L_{\pm 2} \pmod{L_{2^{r+1}}}$$

and therefore

$$2v^2 + 1 \equiv -3 \pmod{L_{2^{r+1}}},$$

i.e.,

$$2v^2 \equiv -4 \pmod{L_{2^{r+1}}}$$

by (3). Since  $L_{2^{r+1}}$  is an odd integer, we get

$$v^2 \equiv -2 \pmod{L_{2^{r+1}}}.$$

This shows that  $\left(\frac{-2}{p}\right) = 1$  for every prime divisor of  $L_{2^{r+1}}$ . Then it follows that

$$p \equiv 1, 3 \pmod{8}$$

by (13) and therefore

$$L_{2^{r+1}} \equiv 1, 3 \pmod{8}.$$

This shows that  $r = 0$  by Lemma 7. Consequently,  $q$  is an odd integer. Therefore it can be easily seen that  $m = 12c + 5$  or  $m = 12c + 7$  for some integer  $c$ . Thus we get

$$L_m \equiv 3 \pmod{8}$$

or

$$L_m \equiv 5 \pmod{8}$$

by (2). On the other hand, since

$$2v^2 = L_m^2 + 1,$$

we get

$$2v^2 \equiv 1 \pmod{L_m},$$

and therefore

$$(2v)^2 \equiv 2 \pmod{L_m}.$$

This shows that  $\left(\frac{2}{p}\right) = 1$  for every prime divisor  $p$  of  $L_m$ . Then it follows that

$$p \equiv \pm 1 \pmod{8}$$

by (12) and therefore

$$L_m \equiv \pm 1 \pmod{8}.$$

But this contradicts with the fact that  $L_m \equiv 3, 5 \pmod{8}$ . Assume that (19) is satisfied. Then we get  $v^2 = 5F_{ms}^2 - 3 = L_{ms}^2 + 1$  by (9). This implies that  $L_{ms} = 0$ , which is impossible. This completes the proof.  $\square$

**Theorem 15.** *If  $L_n = 6L_mx^2$  and  $m \geq 1$ , then  $m = 2$ ,  $x^2 = 1$ , and  $n = 6$ .*

*Proof.* Assume that  $L_n = 6L_mx^2$  for some integer  $x$ . Then  $3|L_n$  and therefore  $n = 2k_0$  for some odd integer  $k_0$  by Theorem 8. Moreover, since  $2|L_n$ , we get  $3|n$  by (10). This shows that  $3|k_0$  and then  $k_0 = 3t$  for some odd integer  $t$ . Thus  $n = 6t = 6(2u + 1) = 12u + 6$ . Therefore

$$L_n = L_{12u+6} \equiv L_6 \pmod{8}$$

by (2). That is,

$$L_n \equiv 2 \pmod{8}.$$

Since  $8 \nmid L_n$ , it can be seen that  $x$  is an odd integer. Therefore

$$x^2 \equiv 1 \pmod{8},$$

which implies that

$$6L_mx^2 \equiv 6L_m \pmod{8}.$$

This shows that

$$6L_m \equiv 2 \pmod{8},$$

which implies that  $m \neq 1$ . Now assume that  $m > 2$ . Since  $L_m | L_n$ , there exists an odd integer  $k$  such that  $n = mk$  by Theorem 8. On the other hand, since  $2 | n$ , it is seen that  $2 | m$ . Therefore  $m = 2r$  for some odd integer  $r$ . If  $r = 6q + 3$ , then  $m = 2r = 12q + 6$  and therefore

$$L_m \equiv L_6 \pmod{8}$$

by (2). That is,

$$L_m \equiv 2 \pmod{8},$$

which is impossible since

$$6L_m \equiv 2 \pmod{8}.$$

Therefore  $3 \nmid r$ . Since  $n = mk$ ,  $m = 2r$  and  $3 \nmid r$ , it follows that  $3 | k$  and thus  $k = 3s$  for some odd integer  $s$ . Then

$$L_n = L_{mk} = L_{3ms} = L_{ms}(L_{ms}^2 - 3) = 6L_mx^2$$

by (6). It can be seen that  $(L_{ms}, L_{ms}^2 - 3) = 3$ . Thus  $\left(L_{ms}, \frac{L_{ms}^2 - 3}{3}\right) = 1$ . Then we get

$$\frac{L_{ms}}{L_m} \left(\frac{L_{ms}^2 - 3}{3}\right) = 2x^2.$$

This shows that

$$\frac{L_{ms}}{L_m} = 2u^2 \text{ and } \frac{L_{ms}^2 - 3}{3} = v^2 \tag{20}$$

or

$$\frac{L_{ms}}{L_m} = u^2 \text{ and } \frac{L_{ms}^2 - 3}{3} = 2v^2 \tag{21}$$

for some integers  $u$  and  $v$ . Assume that (20) is satisfied. Then  $3 \left(\frac{L_{ms}}{3}\right)^2 - 1 = v^2$  and therefore

$$v^2 \equiv -1 \pmod{3},$$

which is a contradiction. Now assume that (21) is satisfied. Then  $L_{ms} = L_mu^2$ , which implies that  $ms = m$  by Theorem 12. That is,  $s = 1$ . Thus  $L_m^2 - 3 = 6v^2$ . Since  $L_m^2 = L_{2m} + 2$  by (5), we see that  $L_{2m} - 1 = 6v^2$ . Moreover, since  $m = 2r$ , it follows that  $L_{4r} - 1 = 6v^2$ . On the other hand, we can write  $4r$  as  $4r = 4(4u \pm 1) = 16u \pm 4 = 2 \cdot 2^k a \pm 4$  for some odd integer  $a$  with  $k \geq 3$ . This shows that

$$L_{4r} = L_{2 \cdot 2^k a \pm 4} \equiv -L_{\pm 4} \pmod{L_{2^k}}$$

by (3) and therefore

$$1 + 6v^2 \equiv -7 \pmod{L_{2^k}}.$$

Then we get

$$6v^2 \equiv -8 \pmod{L_{2^k}}.$$

That is,

$$3v^2 \equiv -4 \pmod{L_{2^k}}.$$



Thus

$$(3v)^2 \equiv -12 \pmod{L_{2^k}}.$$

This shows that  $\left(\frac{-12}{p}\right) = 1$  for every prime divisor  $p$  of  $L_{2^k}$ . Then it follows that

$$p \equiv 1 \pmod{3}$$

and therefore

$$L_{2^k} \equiv 1 \pmod{3}.$$

But this contradicts with Lemma 5. This completes the proof.  $\square$

In [8], the authors showed that  $L_n = 3L_mx^2$  has no solution if  $m > 1$ . Now we give a similar result for Fibonacci numbers.

**Theorem 16.** *Let  $m \geq 3$  be an integer and  $F_n = 3F_mx^2$  for some integer  $x$ . Then  $m = 4$ ,  $x^2 = 16$ , and  $n = 12$ .*

*Proof.* Assume that  $m \geq 3$  and  $F_n = 3F_mx^2$  for some integer  $x$ . Then  $F_m | F_n$  and therefore  $n = mk$  for some integer  $k$  by Theorem 9.

Firstly, assume that  $k$  is an even integer. Then  $k = 2s$  for some  $s \in \mathbb{N}$ . Therefore  $n = mk = 2ms$ . Thus

$$F_n = F_{2ms} = F_{ms}L_{ms} = 3F_mx^2$$

by (7). This shows that

$$(F_{ms}/F_m)L_{ms} = 3x^2.$$

By using Theorem 1, Theorem 12, and Theorem 15, it can be shown that the equation  $(F_{ms}/F_m)L_{ms} = 3x^2$  has no solution.

Now assume that  $k$  is an odd integer. Since  $F_n = 3F_mx^2$ , we get  $4|n$  by Theorem 9. Moreover, since  $n = mk$  and  $k$  is odd, we get  $4|m$ . Assume that  $x$  is an even integer. Then  $4|F_n$ . Thus  $L_3|F_n$  and  $3|n$  by Theorem 10. Therefore since  $4|n$  and  $3|n$ , we get  $12|n$ . That is,  $n = 12t$  for some  $t \in \mathbb{N}$ . On the other hand since  $4|m$ , we get  $m = 4r$  for some  $r \in \mathbb{N}$ . Therefore  $12t = n = mk = 4rk$ . Then it follows that  $3t = rk$ . Thus

$$F_n = F_{12t} = F_{6t}L_{6t} = 3F_{2r}L_{2r}x^2$$

by (7). Since  $(6t/2r) = k$  and  $k$  is odd, we can write

$$\frac{F_{6t}}{F_{2r}} \cdot \frac{L_{6t}}{L_{2r}} = 3x^2.$$

Assume that  $3|r$ . Then, it can be seen that  $\left(\frac{F_{6t}}{F_{2r}}, \frac{L_{6t}}{L_{2r}}\right) = 1$ . Therefore

$$\frac{F_{6t}}{F_{2r}} = u^2, \quad \frac{L_{6t}}{L_{2r}} = 3v^2 \tag{22}$$

or

$$\frac{F_{6t}}{F_{2r}} = 3u^2, \quad \frac{L_{6t}}{L_{2r}} = v^2 \quad (23)$$

for some integers  $u$  and  $v$ . A similar argument shows that (22) and (23) are impossible. Now assume that  $3 \nmid r$ . Then since  $3t = rk$ , it follows that  $3 \mid k$ . Thus  $k = 3s$  for some  $s \in \mathbb{N}$ . Then  $3t = rk = 3rs$  and therefore  $t = rs$ . Also since  $3 \nmid r$ , it can be seen that  $\left(\frac{F_{6t}}{F_{2r}}, \frac{L_{6t}}{L_{2r}}\right) = 2$ .

Therefore

$$\frac{F_{6t}}{F_{2r}} = 2u^2, \quad \frac{L_{6t}}{L_{2r}} = 6v^2 \quad (24)$$

or

$$\frac{F_{6t}}{F_{2r}} = 6u^2, \quad \frac{L_{6t}}{L_{2r}} = 2v^2 \quad (25)$$

for some integers  $u$  and  $v$ . Assume that (24) is satisfied. Then  $2r = 2$  by Theorem 15. This shows that  $r = 1$  and  $t = s$ . Thus  $L_{6t} = 6L_2v^2 = L_6v^2$  and this implies that  $6t = 6$ , i.e.,  $t = 1$  by Theorem 12. Therefore  $k = 3s = 3t = 3$  and  $m = 4r = 4$ . Therefore  $n = 12$  and  $x^2 = 16$ .

Now assume that (25) is satisfied. Then it follows that

$$L_{6t} = 2L_{2r}v^2,$$

which is impossible by Theorem 13 and Theorem 15.

Now assume that  $x$  is an odd integer. Then

$$F_n \equiv 3F_m \pmod{8}.$$

Since  $4 \mid m$ , it follows that  $m = 12q$  or  $m = 12q \pm 4$  for some integer  $q$ . If  $m = 12q \pm 4$ , then

$$F_m \equiv F_{12q \pm 4} \equiv F_{\pm 4} \equiv \pm 3 \pmod{8}$$

by (1). Therefore

$$F_n \equiv \pm 1 \pmod{8},$$

which is impossible since  $4 \mid n$ . Because if  $4 \mid n$ , then  $n = 12r \pm 4$  or  $n = 12r$  for some integer  $r$ , and therefore  $F_n \equiv \pm 3, 0 \pmod{8}$  by (1). If  $m = 12q$ , then  $n = mk = 12qk$ . This shows that  $6qk/6q$  is an odd integer. Then from the identity

$$F_n = F_{12qk} = F_{6qk}L_{6qk} = 3F_mx^2 = 3F_{6q}L_{6q}x^2,$$

it follows that

$$\frac{F_{6qk}}{F_{6q}} \cdot \frac{L_{6qk}}{L_{6q}} = 3x^2.$$

Since  $\left(\frac{F_{6qk}}{F_{6q}}, \frac{L_{6qk}}{L_{6q}}\right) = 1$ , we get

$$\frac{F_{6qk}}{F_{6q}} = u^2, \quad \frac{L_{6qk}}{L_{6q}} = 3v^2 \quad (26)$$

or

$$\frac{F_{6qk}}{F_{6q}} = 3u^2, \quad \frac{L_{6qk}}{L_{6q}} = v^2 \quad (27)$$

for some integers  $u$  and  $v$ . Similarly, it can be seen that (26) and (27) are impossible. This completes the proof.  $\square$

Lastly, we can give the following theorem without proof since its proof is similar to that of Theorem 16.

**Theorem 17.** *There is no integer  $x$  such that  $F_n = 6F_mx^2$ .*

## References

- [1] B. U. Alfred, On square Lucas numbers, *Fibonacci Quart.* **2** (1964), 11–12.
- [2] D. M. Burton, *Elementary Number Theory*, McGraw-Hill, 1998.
- [3] L. Carlitz, A note on Fibonacci numbers, *Fibonacci Quart.* **1** (1964), 15–28.
- [4] J. H. E. Cohn, On square Fibonacci numbers, etc., *J. London Math. Soc.* **39** (1964), 537–540.
- [5] J. H. E. Cohn, Square Fibonacci numbers, etc., *Fibonacci Quart.* **2.2** (1964), 109–113.
- [6] J. H. E. Cohn, Lucas and Fibonacci numbers and some Diophantine equations, *Proc. Glasgow Math. Assoc.* **7** (1965), 24–28.
- [7] J. H. E. Cohn, Squares in some recurrent sequences, *Pacific J. Math.* **41** (1972), 631–646.
- [8] R. Keskin and B. Demirtürk, Fibonacci and Lucas congruences and their applications, *Acta Math. Sin. (Engl. Ser.)* **27** (2011), 725–736.
- [9] T. Koshy, *Fibonacci and Lucas Numbers With Applications*, John Wiley and Sons, Proc., New York-Toronto, 2001.
- [10] L. Moser and L. Carlitz, Advanced problem H-2, *Fibonacci Quart.* **1** (1963), 46.
- [11] I. Niven, H. S. Zuckerman, and H. L. Montgomery, *An Introduction to The Theory of Numbers*, John Wiley & Sons, 1991.

- [12] S. C. Ogilvy, *Tomorrow's Math: Unsolved Problems for the Amateur*, Oxford University Press, 1962.
- [13] P. Ribenboim, Square classes of Fibonacci and Lucas numbers, *Portugal. Math.* **46** (1989), 159–175.
- [14] N. Robbins, On Fibonacci numbers of the form  $px^2$ , where  $p$  is prime, *Fibonacci Quart.* **21** (1983), 266–271.
- [15] N. Robbins, Fibonacci numbers of the form  $cx^2$ , where  $1 \leq c \leq 1000$ , *Fibonacci Quart.* **28** (1990), 306–315.
- [16] N. Robbins, Lucas numbers of the form  $px^2$ , where  $p$  is prime, *Internat. J. Math. Math. Sci.* **14** (1991), 697–703.
- [17] A. P. Rollet, Advanced problem 5080, *Amer. Math. Monthly* **70** (1963), 216.
- [18] S. Vajda, *Fibonacci and Lucas Numbers and The Golden Section*, Ellis Horwood, 1989.
- [19] O. Wyler, In the Fibonacci series  $F_1 = 1$ ,  $F_2 = 1$ ,  $F_{n+1} = F_n + F_{n-1}$  the first, second and twelfth terms are squares, *Amer. Math. Monthly* **71** (1964), 221–222.
- [20] C. Zhou, A general conclusion on Lucas numbers of the form  $px^2$ , where  $p$  is prime, *Fibonacci Quart.* **37** (1999), 39–45.

---

2010 *Mathematics Subject Classification*: Primary 11B37; Secondary 11B39.  
*Keywords*: Fibonacci numbers, Lucas numbers, congruences.

---

(Concerned with sequence [A000045](#).)

---

Received April 29 2011; revised version received August 18 2011; September 25 2011. Published in *Journal of Integer Sequences*, October 16 2011.

---

Return to [Journal of Integer Sequences home page](#).