



On Pairwise Intersections of the Fibonacci, Sierpiński, and Riesel Sequences

Dan Ismailescu

Department of Mathematics

Hofstra University

103 Hofstra University

Hempstead, NY 11549

USA

dan.p.ismailescu@hofstra.edu

Peter Seho Park

Korea International School

373-6 Baekhyeon-dong, Budang-gu

Seongnam-si, Gyonggi-do

Korea

peterseho@yahoo.ca

Abstract

A *Sierpiński* number is an odd integer k with the property that $k \cdot 2^n + 1$ is composite for all positive integer values of n . A *Riesel* number is defined similarly; the only difference is that $k \cdot 2^n - 1$ is composite for all positive integer values of n .

In this paper we find Sierpiński and Riesel numbers among the terms of the well-known Fibonacci sequence. These numbers are smaller than all previously constructed examples. We also find a 23-digit number which is simultaneously a Sierpiński and a Riesel number. This improves on the current record established by Filaseta, Finch and Kozek in 2008. Finally, we prove that there are infinitely many values of n such that the Fibonacci numbers F_n and F_{n+1} are both Sierpiński numbers.

1 Constructing Sierpiński and Riesel numbers

In 1960, Sierpiński [11] proved that there are infinitely many odd numbers k such that $k \cdot 2^n + 1$ is composite for all positive integers n . A few years earlier, Riesel [9] showed that there are infinitely many odd integers k such that $k \cdot 2^n - 1$ is composite for all positive integers n .

The standard way to construct Sierpiński or Riesel numbers is due to Erdős [3]: one seeks a finite collection of triples $\{(a_i, b_i, p_i)\}_{i=1}^{i=t}$ that satisfy the following properties:

- (i) The set $\{(a_i, b_i)\}_{i=1}^{i=t}$ forms a *covering system*, that is, for every $1 \leq i \leq t$, a_i and b_i are integers, $b_i > 1$ and, for all integers n there exists $i \in \{1, \dots, t\}$ such that $n \equiv a_i \pmod{b_i}$.
- (ii) The numbers p_1, \dots, p_t are distinct primes that satisfy $p_i \mid 2^{b_i} - 1$ for all $i \in \{1, \dots, t\}$.

To obtain Sierpiński numbers, one imposes the system of congruences $k \cdot 2^{a_i} \equiv -1 \pmod{p_i}$ for $i \in \{1, \dots, t\}$. Since all primes p_i are odd, each of the above congruences has a solution for k in the form of an arithmetic progression modulo p_i . Then, as p_1, \dots, p_t are distinct primes, by the Chinese remainder theorem all odd positive k that satisfy the system of congruences form an arithmetic sequence modulo $2p_1 \cdots p_t$ (the fact that k must be odd property translates into an additional congruence $k \equiv 1 \pmod{2}$). All such k have the property that $k \cdot 2^n + 1$ is always divisible by at least one p_i for $i \in \{1, \dots, t\}$. Thus, if $k > \max\{p_i \mid i = 1, \dots, t\}$, then $k \cdot 2^n + 1$ is composite for all n , and therefore, k is a Sierpiński number.

In 1962 Selfridge showed that $k = 78557$ is a Sierpiński number by using the triples $\{(0, 2, 3), (1, 3, 7), (1, 4, 5), (11, 12, 13), (15, 36, 19), (27, 36, 37), (3, 36, 73)\}$. It is widely believed this is the smallest Sierpiński number; there are six smaller candidates which have not been eliminated as potential Sierpiński numbers, the smallest of which is 10223. Seventeen or Bust [6] is a distributed computing project which is testing these remaining numbers.

Similarly, to construct a Riesel number, we impose the system of congruences $k \cdot 2^{a_i} \equiv 1 \pmod{p_i}$ for all $i \in \{1, \dots, t\}$. Additionally, consider the congruence $k \equiv 1 \pmod{2}$ to ensure k is odd. Then, as above, by the Chinese remainder theorem, all odd positive k that satisfy the system of congruences form an arithmetic progression with common difference $2p_1 \cdots p_t$. All such k have the property that $k \cdot 2^n - 1$ is always divisible by at least one p_i for $i \in \{1, \dots, t\}$. Thus, if $k > \max\{p_i \mid i = 1, \dots, t\}$, then $k \cdot 2^n - 1$ is composite for all n , and therefore, k has the desired property. Riesel [9] proved that $k = 509203$ is a Riesel number by using the triples $\{(0, 2, 3), (2, 3, 7), (1, 4, 5), (7, 8, 17), (7, 12, 13), (3, 24, 241)\}$. As in the Sierpiński scenario, it is conjectured that 509203 is the smallest Riesel number. There are fifty five smaller numbers which are still being tested as possible candidates — for more information see [7].

2 Sierpiński and Riesel numbers in the Fibonacci sequence

Let F_n be the Fibonacci sequence defined by the recurrence relation $F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$, with $F_0 = 0$ and $F_1 = 1$. The Lucas sequence is defined analogously: $L_0 = 2$, $L_1 = 1$ and $L_n = L_{n-1} + L_{n-2}$ for $n \geq 2$.

Luca and Mejía-Huguet proved in [8] that there are infinitely many Fibonacci numbers which are Sierpiński numbers and infinitely many Fibonacci numbers which are Riesel numbers. Baczkowski, Fasoranti and Finch [1] proved similar results for the Lucas sequence. We use the same technique to improve the results for the Fibonacci numbers.

We construct $\{(a_i, b_i, p_i)\}_{i=1}^{i=t}$ that satisfies conditions (i) and (ii) as described in the previous section. Let k be a Sierpiński or Riesel number in the arithmetic sequence constructed from the above system. Then, $k \equiv \epsilon 2^{-a_i} \pmod{p_i}$ for all $i = 1, \dots, t$, where $\epsilon = -1$ for the Sierpiński case, and $\epsilon = 1$ for the Riesel case. For a positive integer m , let $h(m)$ be the m -th Pisano period, the period with which the Fibonacci sequence repeats modulo m . The existence of $h(m)$ is easy to establish, see, e. g., [12]. For every integer x , let

$$\mathcal{A}(x, m) = \{0 \leq y \leq h(m) - 1 \mid F_y \equiv x \pmod{m}\} + h(m)\mathbb{Z}. \quad (1)$$

In other words, $\mathcal{A}(x, m)$ denotes the set of all y modulo $h(m)$ such that $F_y \equiv x \pmod{m}$. For k to be a Fibonacci number, the congruence $k \equiv F_{y_i} \pmod{p_i}$ must have a solution y_i for each $i = 1, \dots, t$. Here, the set of possible solutions y_i is $\mathcal{A}(\epsilon 2^{-a_i}, p_i)$, where $\epsilon = -1$ for the Sierpiński case, and $\epsilon = 1$ for the Riesel case. So, in order for k to be a Fibonacci number, it is necessary and sufficient that

$$\bigcap_{i=1}^t \mathcal{A}(\epsilon 2^{-a_i}, p_i) \neq \emptyset.$$

Finally, since Sierpiński and Riesel numbers are odd, it is necessary that if $k = F_n$ (for k that is either a Sierpiński or Riesel number), then n cannot be divisible by 3.

Luca and Mejía-Huguet [8] proved the following:

Theorem 1. *For every $n \equiv 1807873 \pmod{3543120}$, F_n is a Riesel number.*

We managed to find two arithmetic sequences with the same property but smaller first terms. The first such sequence has a smaller common difference as well.

Theorem 2.

- (a) For every $n \equiv 820438 \pmod{1517040}$, F_n is a Riesel number.
- (b) For every $n \equiv 385297 \pmod{3543120}$, F_n is a Riesel number.

Proof. (a) Let $t = 7$ and

$$\{(a_i, b_i, p_i)\}_{i=1}^{i=7} = \{(1, 2, 3), (2, 4, 5), (4, 8, 17), (8, 16, 257), (0, 48, 7), (32, 48, 13), (16, 48, 97)\}.$$

It is easy to check that conditions (i) and (ii) are satisfied. For these choices of the primes p_i we have $(h(p_1), \dots, h(p_7)) = (8, 20, 36, 516, 16, 28, 196)$. It follows that

$$\begin{aligned} \mathcal{A}(2^{-a_1}, p_1) &= \mathcal{A}(2, 3) = \{3, 5, \mathbf{6}\} \pmod{8} \\ \mathcal{A}(2^{-a_2}, p_2) &= \mathcal{A}(4, 5) = \{9, 11, 12, \mathbf{18}\} \pmod{20} \\ \mathcal{A}(2^{-a_3}, p_3) &= \mathcal{A}(16, 17) = \{17, 19, 20, \mathbf{34}\} \pmod{36} \\ \mathcal{A}(2^{-a_4}, p_4) &= \mathcal{A}(256, 257) = \{257, 259, 260, \mathbf{514}\} \pmod{516} \\ \mathcal{A}(2^{-a_5}, p_5) &= \mathcal{A}(1, 7) = \{1, 2, \mathbf{6}, 15\} \pmod{16} \\ \mathcal{A}(2^{-a_6}, p_6) &= \mathcal{A}(3, 13) = \{4, \mathbf{10}\} \pmod{28} \\ \mathcal{A}(2^{-a_7}, p_7) &= \mathcal{A}(35, 97) = \{116, \mathbf{178}\} \pmod{196} \end{aligned}$$

It can be checked that if $n \equiv 820438 \pmod{1517040}$, then n belongs to all $\mathcal{A}(2^{-a_i}, p_i)$ for $i = 1, \dots, 7$. The numbers in boldface in the above sets represent the remainders of n modulo $h(p_i)$. Furthermore, since $820438 \equiv 1 \pmod{3}$ and $1517040 \equiv 0 \pmod{3}$, no n in the above arithmetic sequence is divisible by 3. Thus, for all such n , F_n is a Riesel number.

(b) Let $t = 7$ and

$$\{(a_i, b_i, p_i)\}_{i=1}^{i=7} = \{(0, 2, 3), (0, 3, 7), (3, 4, 5), (5, 12, 13), (13, 36, 19), (1, 36, 37), (25, 36, 73)\}.$$

As in part (a), it is easy to verify that conditions (i) and (ii) are satisfied. For these choices of the primes p_i we have $(h(p_1), \dots, h(p_7)) = (8, 16, 20, 28, 18, 76, 148)$. Straightforward computations show that

$$\begin{aligned} \mathcal{A}(2^{-a_1}, p_1) &= \mathcal{A}(1, 3) = \{\mathbf{1}, 2, 7\} \pmod{8} \\ \mathcal{A}(2^{-a_2}, p_2) &= \mathcal{A}(1, 7) = \{\mathbf{1}, 2, 6, 15\} \pmod{16} \\ \mathcal{A}(2^{-a_3}, p_3) &= \mathcal{A}(2, 5) = \{3, 14, 16, \mathbf{17}\} \pmod{20} \\ \mathcal{A}(2^{-a_4}, p_4) &= \mathcal{A}(11, 13) = \{11, \mathbf{17}\} \pmod{28} \\ \mathcal{A}(2^{-a_5}, p_5) &= \mathcal{A}(13, 19) = \{\mathbf{7}, 11\} \pmod{18} \\ \mathcal{A}(2^{-a_6}, p_6) &= \mathcal{A}(19, 37) = \{23, 48, \mathbf{53}, 66\} \pmod{76} \\ \mathcal{A}(2^{-a_7}, p_7) &= \mathcal{A}(4, 73) = \{\mathbf{53}, 95\} \pmod{148} \end{aligned}$$

It is easy to check that if $n \equiv 385297 \pmod{3543120}$, then n belongs to all $\mathcal{A}(2^{-a_i}, p_i)$ for $i = 1, \dots, 7$. The numbers in boldface in the above sets represent the remainders of n modulo $h(p_i)$. Furthermore, since $385297 \equiv 1 \pmod{3}$ and $3543120 \equiv 0 \pmod{3}$, no n in the above arithmetic sequence is divisible by 3. Thus, for all such n , F_n is a Riesel number. \square

In the same paper, Luca and Mejía-Huguet proved the following:

Theorem 3. *For every $n \equiv 20808199653121 \pmod{206353240410240}$, F_n is a Sierpiński number.*

Notice that the first term of the above arithmetic sequence is a 14-digit number while the common difference is a 15-digit number. We manage to improve this result by finding an arithmetic sequence with the same property, but with only a 6-digit initial term and a 7-digit common difference.

Theorem 4. *For every $n \equiv 696602 \pmod{1517040}$, F_n is a Sierpiński number.*

Proof. Let $t = 7$ and

$$\{(a_i, b_i, p_i)\}_{i=1}^{i=7} = \{(1, 2, 3), (0, 3, 7), (2, 4, 5), (4, 8, 17), (8, 12, 13), (16, 48, 97), (40, 48, 257)\}.$$

It is easy to verify that conditions (i) and (ii) are satisfied. For this selection of the primes p_i it is then straightforward to find the corresponding Pisano periods: $(h(p_1), \dots, h(p_7)) = (8, 16, 20, 36, 28, 196, 516)$. It follows that

$$\begin{aligned} \mathcal{A}(-2^{-a_1}, p_1) &= \mathcal{A}(1, 3) = \{1, \mathbf{2}, 7\} \pmod{8} \\ \mathcal{A}(-2^{-a_2}, p_2) &= \mathcal{A}(6, 7) = \{7, 9, \mathbf{10}, 14\} \pmod{16} \\ \mathcal{A}(-2^{-a_3}, p_3) &= \mathcal{A}(1, 5) = \{1, \mathbf{2}, 8, 19\} \pmod{20} \\ \mathcal{A}(-2^{-a_4}, p_4) &= \mathcal{A}(1, 17) = \{1, \mathbf{2}, 16, 35\} \pmod{36} \\ \mathcal{A}(-2^{-a_5}, p_5) &= \mathcal{A}(10, 13) = \{\mathbf{18}, 24\} \pmod{28} \\ \mathcal{A}(-2^{-a_6}, p_6) &= \mathcal{A}(62, 97) = \{\mathbf{18}, 80\} \pmod{196} \\ \mathcal{A}(-2^{-a_7}, p_7) &= \mathcal{A}(1, 257) = \{1, \mathbf{2}, 256, 515\} \pmod{516} \end{aligned}$$

If $n \equiv 696602 \pmod{1517040}$, then n belongs to all $\mathcal{A}(-2^{-a_i}, p_i)$ for $i = 1, \dots, 7$. As before, the boldface numbers in the above sets represent the remainders of n modulo $h(p_i)$. Furthermore, since $696602 \equiv 2 \pmod{3}$ and $1517040 \equiv 0 \pmod{3}$, no n in the above arithmetic progression is divisible by 3. Thus, for all such n , F_n is a Sierpiński number. \square

3 Consecutive Fibonacci-Sierpiński numbers

Despite their relative scarcity, we were able to find infinitely many values of n such that *both* F_n and F_{n+1} are Fibonacci-Sierpiński numbers.

Theorem 5. *For every $n \equiv 1510614062400961 \pmod{3021228124801920}$, both F_n and F_{n+1} are Fibonacci-Sierpiński numbers.*

Proof. Take $t = 7$ and

$$\{(a_i, b_i, p_i)\}_{i=1}^{i=7} = \{(1, 2, 3), (2, 4, 5), (4, 8, 17), (8, 16, 257), (16, 32, 65537), \\ (0, 64, 641), (32, 64, 6700417)\}.$$

We see that conditions (i) and (ii) are fulfilled. The Pisano periods sequence in this case is $(h(p_1), \dots, h(p_7)) = (8, 20, 36, 516, 14564, 640, 13400836)$. Then,

$$\begin{aligned} \mathcal{A}(-2^{-a_1}, p_1) &= \mathcal{A}(1, 3) = \{\mathbf{1}, \mathbf{2}, 7\} \pmod{8} \\ \mathcal{A}(-2^{-a_2}, p_2) &= \mathcal{A}(1, 5) = \{\mathbf{1}, \mathbf{2}, 8, 19\} \pmod{20} \\ \mathcal{A}(-2^{-a_3}, p_3) &= \mathcal{A}(1, 17) = \{\mathbf{1}, \mathbf{2}, 16, 35\} \pmod{36} \\ \mathcal{A}(-2^{-a_4}, p_4) &= \mathcal{A}(1, 257) = \{\mathbf{1}, \mathbf{2}, 256, 515\} \pmod{516} \\ \mathcal{A}(-2^{-a_5}, p_5) &= \mathcal{A}(1, 65537) = \{\mathbf{1}, \mathbf{2}, 7280, 14563\} \pmod{14564} \\ \mathcal{A}(-2^{-a_6}, p_6) &= \mathcal{A}(640, 641) = \{319, \mathbf{321}, \mathbf{322}, 638\} \pmod{640} \\ \mathcal{A}(-2^{-a_7}, p_7) &= \mathcal{A}(1, 6700417) = \{\mathbf{1}, \mathbf{2}, 6700416, 13400835\} \pmod{13400836} \end{aligned}$$

One sees that if $n \equiv 1510614062400961 \pmod{3021228124801920}$ then both n and $n + 1$ belongs to all sets $\mathcal{A}(-2^{-a_i}, p_i)$ for $i = 1, \dots, 7$. The boldface numbers in the sets above are the remainders of n and $n + 1$ modulo $h(p_i)$. Checking that $1510614062400961 \equiv 1 \pmod{3}$, $1510614062400962 \equiv 2 \pmod{3}$, and $3021228124801920 \equiv 0 \pmod{3}$, it follows that that no number in either of the two arithmetic sequences above is divisible by 3. Thus, for all such n , both F_n and F_{n+1} are Sierpiński numbers, thus proving the assertion in the theorem. \square

It would be interesting to decide if there exist consecutive Fibonacci numbers both of which are Riesel numbers. We were unsuccessful in constructing such an example.

4 Riesel-Sierpiński numbers

Cohen and Selfridge [2] were the first to prove that there exist infinitely many integers that are simultaneously Sierpiński numbers and Riesel numbers. Their discovery came in connection to the question whether there exist integers that cannot be written as the sum or difference of two prime powers. Apparently unaware of this result, Brier, Gallot and Vantieghem constructed other Riesel-Sierpiński numbers, but none smaller than Cohen and Selfridge's record, which is 26-digits long. For an interesting account of their efforts the reader is referred to [10, 5].

It was not until 2008 when Filaseta, Finch and Kozek [4] found a Riesel-Sierpiński number with only 24 digits: 143665583045350793098657. We were able to slightly refine their approach and construct a Riesel-Sierpiński number with only 23 digits. We present the details below.

To find a Riesel-Sierpiński number, one constructs a set of triples $\{(a_i, b_i, p_i)\}_{i=1}^{i=t}$ to obtain an arithmetic sequence of Sierpiński numbers and a set of triples $\{(a'_i, b'_i, p'_i)\}_{i=1}^{i=t'}$ which

generates an arithmetic sequence of Riesel numbers, with both sets of triples satisfying (i), (ii). If additionally we impose the condition that

$$\{k \mid 2^{a_i} k \equiv -1 \pmod{p_i} \forall i \in \{1, \dots, t\}\} \cap \{k' \mid 2^{a'_i} k' \equiv 1 \pmod{p'_i} \forall i \in \{1, \dots, t'\}\} \neq \emptyset,$$

then the Sierpiński numbers k and the Riesel numbers k' are each in an arithmetic sequence modulo $p_1 \cdots p_t$ and modulo $p'_1 \cdots p'_{t'}$, respectively. Taking into account the additional congruences $k \equiv 1 \pmod{2}$ and $k' \equiv 1 \pmod{2}$, the Chinese remainder theorem guarantees that $\{p_1, \dots, p_t\} \cap \{p'_1, \dots, p'_{t'}\} = \emptyset$ is sufficient to prove that there are infinitely many integers belonging to both arithmetic sequences.

However, this condition would require that $p = 3$ and the corresponding $b = 2$ can only be used once: either for the Sierpiński part or for the Riesel part.

Fortunately, $p = 3$ and $b = 2$ can be used for both Sierpiński and Riesel cases. As shown by Cohen and Selfridge in [2], if $\{p_1, \dots, p_t\} \cap \{p'_1, \dots, p'_{t'}\} = \{3\}$, and $(a_i, b_i, p_i)_{i=1}^{i=t}$ and $(a'_i, b'_i, p'_i)_{i=1}^{i=t'}$ have distinct a values (0 or 1) corresponding to $b = 2$ and $p = 3$, then there exist integers belonging to both arithmetic sequences.

Theorem 6. *Every $n \equiv 10439679896374780276373 \pmod{66483084961588510124010691590}$ is a Sierpiński-Riesel number.*

Proof. Let $t = 7$ and

$$\{(a_i, b_i, p_i)\}_{i=1}^{i=7} = \{(0, 2, 3), (3, 4, 5), (5, 8, 17), (9, 16, 257), (9, 48, 13), (17, 48, 97), (1, 48, 241)\}.$$

Let us notice that in this step we use the same set of primes as Filaseta, Finch and Kozek. It is straightforward to verify that the triples $(a_i, b_i, p_i)_{i=1}^{i=7}$ satisfy conditions (i) and (ii). With these choices the congruences $k \cdot 2^{a_i} \equiv -1 \pmod{p_i}$ imply that $k \equiv 2 \pmod{3}$, $k \equiv 3 \pmod{5}$, $k \equiv 9 \pmod{17}$, $k \equiv 129 \pmod{257}$, $k \equiv 5 \pmod{13}$, $k \equiv 31 \pmod{97}$ and $k \equiv 120 \pmod{241}$. Using the Chinese remainder theorem, we obtain that

$$k \equiv 18354878963 \pmod{19916152035}. \quad (2)$$

Every odd number satisfying (2) is a Sierpiński number. Let us now work the Riesel part. Let $t = 12$ and

$$\{(a'_i, b'_i, p'_i)\}_{i=1}^{i=12} = \{(1, 2, 3), (0, 3, 7), (8, 9, 73), (2, 18, 19), (32, 36, 37), (14, 36, 109), (4, 5, 31), (6, 10, 11), (10, 15, 151), (28, 30, 331), (12, 20, 41), (22, 60, 61)\}.$$

Filaseta, Finch and Kozek considered an almost identical set of primes: the only difference is that they used the prime 1321 instead of 41. It is not hard to check that the triples $(a'_i, b'_i, p'_i)_{i=1}^{i=12}$ satisfy conditions (i) and (ii). With these choices, the congruences $k \cdot 2^{a'_i} \equiv 1 \pmod{p'_i}$ imply that $k \equiv 2 \pmod{3}$, $k \equiv 1 \pmod{7}$, $k \equiv 2 \pmod{73}$, $k \equiv 5 \pmod{19}$, $k \equiv 16 \pmod{37}$, $k \equiv 93 \pmod{109}$, $k \equiv 2 \pmod{31}$, $k \equiv 5 \pmod{11}$, $k \equiv 32 \pmod{151}$, $k \equiv 4 \pmod{331}$, $k \equiv 10 \pmod{41}$, $k \equiv 49 \pmod{61}$.

Using the Chinese remainder theorem, we obtain that

$$k \equiv 4625814406597377449 \pmod{5007223647777439011}. \quad (3)$$

Every odd number satisfying (3) is a Riesel number. Now let us combine (2) and (3) using the Chinese remainder theorem for one last time. We get that

$$k \equiv 10439679896374780276373 \pmod{33241542480794255062005345795}$$

Taking into account that k must be odd, the statement in the theorem follows. \square

5 Conclusions

In this paper we find subsequences of the Fibonacci sequence whose terms have either the Sierpiński property or the Riesel property. We also construct a new arithmetic sequence, all of whose terms are simultaneously Sierpiński and Riesel numbers. While the existence of such numbers was known before, our main contribution is that our results are smaller than any of the previous ones.

The interesting question is whether there is a number which has *all three* properties simultaneously. Obtaining this would require the construction of two sets of triples (one for the Sierpiński case and one for the Riesel case) sharing no b value other than 2, and satisfying (i), (ii), and the condition that each set of triples generates a Fibonacci-Sierpiński or Fibonacci-Riesel arithmetic progression. If the two arithmetic progressions have nonempty intersection, then that intersection would be an arithmetic sequence of Fibonacci indices n for which F_n would be both Sierpiński and Riesel numbers. Unfortunately, so far we were unable to find such an example.

6 Acknowledgements

The authors would like to thank the anonymous referee whose thorough and competent suggestions greatly improved the exposition and the content of this paper.

References

- [1] D. Baczkowski, O. Fisoranti, and C. E. Finch, Lucas-Sierpiński and Lucas-Riesel numbers, *Fibonacci Quart.* **49** (2011), 334–339.
- [2] F. Cohen and J. L. Selfridge, Not every number is the sum or difference of two prime powers. *Math. Comp.* **29** (1975), 79–81.
- [3] P. Erdős, On integers of the form $2^k + p$ and some related problems. *Summa Brasil. Math.* **2** (1950), 113–123.

- [4] M. Filaseta, C. Finch, and M. Kozek, On powers associated with Sierpiński Numbers, Riesel numbers and Polignac’s conjecture, *J. Number Theory* **128** (2008), 1916–1940.
- [5] Y. Gallot, A search for some small Brier numbers, 2000, manuscript,
<http://yves.gallot.pagesperso-orange.fr/papers/smallbrier.pdf>.
- [6] L. Helm and D. Norris, Seventeen or bust: a distributed attack on the Sierpiński problem,
<http://www.seventeenorbust.com/>.
- [7] W. Keller, The Riesel problem: definition and status,
<http://www.prothsearch.net/rieselprob.html>.
- [8] F. Luca and V. J. Mejía-Huguet, Fibonacci-Riesel and Fibonacci-Sierpiński numbers,
Fibonacci Quart. **46/47** (2008/2009), 216–219.
- [9] H. Riesel, Några stora primtal, *Elementa* **39** (1956), 258–260.
- [10] C. Rivera, The prime puzzles & problem connection,
http://www.primepuzzles.net/problems/prob_029.htm.
- [11] W. Sierpiński, Sur un problème concernant les nombres $k \cdot 2^n + 1$, *Elem. Math.* **15** (1960), 73–74.
- [12] D. D. Wall, Fibonacci series modulo m , *Amer. Math. Monthly*, **67** (1960), 525–532.

2010 *Mathematics Subject Classification*: Primary 11B25; Secondary 11B50.

Keywords: Fibonacci number, Riesel number, Sierpiński number, covering congruence.

(Concerned with sequences [A000045](#), [A076336](#), and [A076337](#).)

Received June 1 2013; revised version received December 4 2013. Published in *Journal of Integer Sequences*, December 5 2013.

Return to [Journal of Integer Sequences home page](#).