



More Determinant Representations for Sequences

Alireza Moghaddamfar and Hadiseh Tajbakhsh

Department of Mathematics

K. N. Toosi University of Technology

P. O. Box 16315-1618

Tehran, Iran

and

Research Institute for Fundamental Sciences (RIFS)

Tabriz, Iran

moghadam@kntu.ac.ir

moghadam@ipm.ir

Abstract

In this paper, we will find some new families of infinite (integer) matrices whose entries satisfy a non-homogeneous recurrence relation and such that the sequence of their leading principal minors is a subsequence of the Fibonacci, Lucas, Jacobsthal, or Pell sequences.

1 Introduction

Throughout this paper, unless noted otherwise, we will use the following notation. Let $\alpha = (\alpha_i)_{i \geq 0}$ and $\beta = (\beta_i)_{i \geq 0}$ be two arbitrary sequences starting with a common first term $\alpha_0 = \beta_0$. We denote by $P_{\alpha, \beta}(n)$ the *generalized Pascal triangle* associated with the sequences α and β , which is introduced as follows. Actually, $P_{\alpha, \beta}(n) = [P_{i, j}]_{0 \leq i, j \leq n}$ is a square matrix of order $n + 1$ whose (i, j) -entry $P_{i, j}$ obeys the following rules:

$$P_{i, 0} = \alpha_i, \quad P_{0, j} = \beta_j \quad \text{for } i, j = 0, 1, 2, \dots, n, \quad \text{and} \quad P_{i, j} = P_{i, j-1} + P_{i-1, j} \quad \text{for } 1 \leq i, j \leq n.$$

We also denote by $T_{\alpha,\beta}(n) = [T_{i,j}]_{0 \leq i,j \leq n}$ the *Toeplitz matrix* of order $n + 1$ whose (i, j) -entry $T_{i,j}$ obeys the following rules:

$$T_{i,0} = \alpha_i, \quad T_{0,j} = \beta_j \quad \text{for } i, j = 0, 1, 2, \dots, n, \quad \text{and } T_{i,j} = T_{k,l} \quad \text{if } i - j = k - l.$$

The *unipotent lower triangular matrix* $L(n) = [L_{i,j}]_{0 \leq i,j \leq n}$ is again a square matrix of order $n + 1$ with entries:

$$L_{i,j} = \begin{cases} 0, & \text{if } 0 \leq i < j \leq n; \\ \binom{i}{j}, & \text{if } 0 \leq j \leq i \leq n. \end{cases}$$

We put $U(n) = L(n)^t$, where A^t signifies the transpose of matrix A . Moreover, a *lower Hessenberg matrix* $H(n) = [H_{i,j}]_{0 \leq i,j \leq n}$ is a square matrix of order $n + 1$, where $H_{i,j} = 0$ whenever $j > i + 1$ and $H_{i,i+1} \neq 0$ for some i , $0 \leq i \leq n - 1$.

Given a matrix A , we denote by $R_i(A)$ (resp., $C_j(A)$) the row i (resp., the column j) of A . We also denote by $A^{[1]}$ the submatrix obtained from A by deleting the first column of A .

Given a sequence $\varphi = (\varphi_i)_{i \geq 0}$, define the *binomial transform* of φ to be the sequence $\hat{\varphi} = (\hat{\varphi}_i)_{i \geq 0}$ with

$$\hat{\varphi}_i = \sum_{k=0}^i (-1)^{i+k} \binom{i}{k} \varphi_k.$$

The *Fibonacci sequence* ([A000045](#) in [3]) is defined by the recurrence relation:

$$F_0 = 0, \quad F_1 = 1, \quad F_n = F_{n-1} + F_{n-2} \quad \text{for } n \geq 2.$$

The *Lucas sequence* ([A000032](#) in [3]) is defined by the recurrence relation:

$$L_0 = 2, \quad L_1 = 1, \quad L_n = L_{n-1} + L_{n-2} \quad \text{for } n \geq 2.$$

The *Jacobsthal sequence* ([A001045](#) in [3]) is defined by the recurrence relation:

$$J_0 = 0, \quad J_1 = 1, \quad J_n = J_{n-1} + 2J_{n-2} \quad \text{for } n \geq 2.$$

The *Pell sequence* ([A000129](#) in [3]) is defined by the recurrence relation:

$$P_0 = 0, \quad P_1 = 1, \quad P_n = 2P_{n-1} + P_{n-2} \quad \text{for } n \geq 2.$$

Let $A = [A_{i,j}]_{i,j \geq 0}$ be an arbitrary infinite matrix. We denote the elementary row operation of type *three* by $O_{r,s}(\lambda)$, where $r \neq s$ and λ a scalar, that is

$$R_k(O_{r,s}(\lambda)A) = \begin{cases} R_r(A) + \lambda R_s(A), & \text{if } k = r; \\ R_k(A), & \text{if } k \neq r. \end{cases}$$

The n th *leading principal minor* of A , denoted by $d_n(A)$, is defined as follows:

$$d_n(A) = \det[A_{i,j}]_{0 \leq i,j \leq n}, \quad (n = 0, 1, 2, 3, \dots).$$

We put $D(A) = (d_n(A))_{n \geq 0}$. Two infinite matrices A and B are said to be *equimodular* if $D(A) = D(B)$. Given a sequence $\omega = (\omega_n)_{n \geq 0}$, a family $\{A_t \mid t \in I\}$ of equimodular matrices are said to be ω -*equimodular* if $D(A_t) = \omega$ for all $t \in I$. We will denote the family of ω -equimodular matrices by \mathcal{A}_ω . The infinite matrices in \mathcal{A}_ω are said to be *determinant representations of ω* . Note that for *any* sequence $\omega = (\omega_n)_{n \geq 0}$, there is a determinant representation of ω , in other words $\mathcal{A}_\omega \neq \emptyset$. Indeed, expanding along the last rows, it is easy to see that

$$\begin{pmatrix} \omega_0 & 1 & * & * & * & \cdots \\ -\omega_1 & 0 & 1 & * & * & \cdots \\ \omega_2 & 0 & 0 & 1 & * & \cdots \\ -\omega_3 & 0 & 0 & 0 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \in \mathcal{A}_\omega,$$

(see also Theorem 3.2 and the Remark after this theorem in [4]). Especially, there are many different determinant representations of ω , when ω is a (sub-)sequence of Fibonacci, Lucas, Jacobsthal and Pell sequences. Some examples of such matrices can be found in [1, 2].

In this paper, we are going to find some determinant representations of the sequences:

$$\mathcal{F} = (F_{n+1})_{n \geq 0}, \quad \mathcal{L} = (L_{n+1})_{n \geq 0}, \quad \mathcal{J} = (J_{n+1})_{n \geq 0} \quad \text{and} \quad \mathcal{P} = (P_{n+1})_{n \geq 0}.$$

It is worthwhile to point out that we will use *non-homogeneous recurrence relations* to construct these determinant representations.

In the sequel, we introduce a new family of (infinite) matrices $A(\infty) = [A_{i,j}]_{i,j \geq 0}$, whose entries obey a non-homogeneous recurrence relation. Actually, for two constants u and v , and arbitrary sequences $\lambda = (\lambda_i)_{i \geq 0}$ and $\mu = (\mu_i)_{i \geq 0}$ with $\mu_0 = 0$, the first column and row of matrix $A(\infty)$ are the sequences

$$(A_{i,0})_{i \geq 0} = (\lambda_0, \lambda_1, \lambda_2, \dots, A_{i,0} = \lambda_i, \dots),$$

and

$$(A_{0,j})_{j \geq 0} = (\lambda_0, \lambda_0 + u, \lambda_0 + 2u, \dots, A_{0,j} = \lambda_0 + ju, \dots),$$

respectively, while the remaining entries $A_{i,j}$ ($i, j \geq 1$) are obtained from the following non-homogeneous recurrence relation:

$$A_{i,j} = A_{i,j-1} + A_{i-1,j} - \lambda_{i-1} + \mu_i - \mu_{i-1} + (j-1)(v-u), \quad i, j \geq 1.$$

We denote by $A(n)$ the submatrix of $A(\infty)$ consisting of the entries in its first $n+1$ rows and columns. The matrix $A(3)$, for example, is then given by

$$A(3) = \begin{pmatrix} \lambda_0 & \lambda_0 + u & \lambda_0 + 2u & \lambda_0 + 3u \\ \lambda_1 & \lambda_1 + \mu_1 + u & \lambda_1 + 2\mu_1 + 2u + v & \lambda_1 + 3\mu_1 + 3u + 3v \\ \lambda_2 & \lambda_2 + \mu_2 + u & \lambda_2 + 2\mu_2 + \mu_1 + 2u + 2v & \lambda_2 + 3\mu_2 + 3\mu_1 + 3u + 7v \\ \lambda_3 & \lambda_3 + \mu_3 + u & \lambda_3 + 2\mu_3 + \mu_2 + \mu_1 + 2u + 3v & \lambda_3 + 3\mu_3 + 3\mu_2 + 4\mu_1 + 3u + 12v \end{pmatrix}.$$

Finally, the main result of this paper can be stated as follows:

Main Theorem. *The matrix $A(n)$, $n \geq 0$, defined as above, satisfies the following statements:*

(a) $A(n) = L(n) \cdot H(n) \cdot U(n)$, where

$$H(n) = \left(\begin{array}{c|cccc} \hat{\lambda}_0 & u & 0 & \cdots & 0 \\ \hat{\lambda}_1 & & & & \\ \hat{\lambda}_2 & & & & \\ \vdots & & & & \\ \hat{\lambda}_n & & & T_{(\hat{\mu}_1, \hat{\mu}_2, \hat{\mu}_3, \dots), (\hat{\mu}_1, v, 0, 0, \dots)}(n-1) & \end{array} \right).$$

In particular, we have $\det(A(n)) = \det(H(n))$.

(b) In the case when $u = v = 1$ and $\lambda_i = (2^i - 1)c + 1$, we have the following statements:

(b. 1) if $\mu_i = \left(2^i + \frac{(i-2)(i+1)}{2}\right)c - \frac{i(i-3)}{2}$, then $\det(A(n)) = F_{n+1}$.

(b. 2) if $\mu_i = \left(\frac{5 \cdot 3^i}{4} - 2^i - \frac{2i+1}{4}\right)c + \frac{5(3^i-1)}{4} + \frac{i}{2}$, then $\det(A(n)) = L_{n+1}$.

(b. 3) if $\mu_i = i^2c - i^2 + 2i$, then $\det(A(n)) = J_{n+1}$.

(b. 4) if $\mu_i = \left(2^{i+1} + \frac{(i+1)(i-4)}{2}\right)c + \frac{(5-i)i}{2}$, then $\det(A(n)) = P_{n+1}$.

As mentioned previously, we have obtained some determinant representations of the sequences:

$$\mathcal{F} = (F_{n+1})_{n \geq 0}, \quad \mathcal{L} = (L_{n+1})_{n \geq 0}, \quad \mathcal{J} = (J_{n+1})_{n \geq 0} \quad \text{and} \quad \mathcal{P} = (P_{n+1})_{n \geq 0},$$

which are presented in the following:

$$\left(\begin{array}{cccc} 1 & 2 & 3 & \cdots \\ c+1 & 2c+3 & 3c+6 & \cdots \\ 3c+1 & 7c+3 & 12c+8 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{array} \right) \in \mathcal{A}_{\mathcal{F}}, \quad \left(\begin{array}{cccc} 1 & 2 & 3 & \cdots \\ c+1 & 2c+5 & 3c+10 & \cdots \\ 3c+1 & 9c+13 & 16c+30 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{array} \right) \in \mathcal{A}_{\mathcal{L}},$$

$$\left(\begin{array}{cccc} 1 & 2 & 3 & \cdots \\ c+1 & 2c+3 & 3c+6 & \cdots \\ 3c+1 & 7c+2 & 12c+6 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{array} \right) \in \mathcal{A}_{\mathcal{J}} \quad \text{and} \quad \left(\begin{array}{cccc} 1 & 2 & 3 & \cdots \\ c+1 & 2c+4 & 3c+8 & \cdots \\ 3c+1 & 8c+5 & 14c+13 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{array} \right) \in \mathcal{A}_{\mathcal{P}}.$$

2 Main results

As the first result of this paper, we consider the following theorem.

Theorem 1. For two arbitrary sequences $(\lambda_i)_{i \geq 0}$ and $(\mu_i)_{i \geq 0}$, with $\mu_0 = 0$, and some integers u and v , let $A(\infty) = [A_{i,j}]_{i,j \geq 0}$ be an infinite dimensional matrix whose entries are given by

$$A_{i,j} = A_{i,j-1} + A_{i-1,j} - \lambda_{i-1} + \mu_i - \mu_{i-1} + (j-1)(v-u), \quad i, j \geq 1 \quad (1)$$

and the initial conditions $A_{i,0} = \lambda_i$ and $A_{0,i} = \lambda_0 + iu$, $i \geq 0$. If $A(n) = [A_{i,j}]_{0 \leq i,j \leq n}$, then we have

$$A(n) = L(n) \cdot H(n) \cdot U(n), \quad (2)$$

where

$$H(n) = \left(\begin{array}{c|cccc} \hat{\lambda}_0 & u & 0 & \cdots & 0 \\ \hat{\lambda}_1 & & & & \\ \hat{\lambda}_2 & & & & \\ \vdots & & & & \\ \hat{\lambda}_n & & & & \end{array} \right). \quad T_{(\hat{\mu}_1, \hat{\mu}_2, \hat{\mu}_3, \dots), (\hat{\mu}_1, v, 0, 0, \dots)}(n-1)$$

Proof. First of all, we recall that the entries of $L(n) = [L_{i,j}]_{0 \leq i,j \leq n}$ satisfy the following recurrence

$$L_{i,j} = L_{i-1,j-1} + L_{i-1,j}, \quad 1 \leq i, j \leq n. \quad (3)$$

Similarly, for the entries of $U(n) = [U_{i,j}]_{0 \leq i,j \leq n}$ we have

$$U_{i,j} = U_{i-1,j-1} + U_{i,j-1}, \quad 1 \leq i, j \leq n. \quad (4)$$

In what follows, for convenience, we will let $A = A(n)$, $L = L(n)$, $H = H(n)$ and $U = U(n)$. Now, for the proof of the desired factorization we compute the (i, j) -entry of $L \cdot H \cdot U$, that is

$$(L \cdot H \cdot U)_{i,j} = \sum_{r=0}^n \sum_{s=0}^n L_{i,r} H_{r,s} U_{s,j}. \quad (5)$$

In fact, we should establish

$$\begin{aligned} R_0(L \cdot H \cdot U) &= R_0(A) = (\lambda_0, \lambda_0 + u, \dots, \lambda_0 + nu), \\ C_0(L \cdot H \cdot U) &= C_0(A) = (\lambda_0, \lambda_1, \dots, \lambda_n), \end{aligned}$$

and finally, show that

$$(L \cdot H \cdot U)_{i,j} = (L \cdot H \cdot U)_{i-1,j-1} + (L \cdot H \cdot U)_{i-1,j} - \lambda_{i-1} + \mu_i - \mu_{i-1} + (j-1)(v-u), \quad (6)$$

for $1 \leq i, j \leq n$.

Let us do the required calculations. Assume first that $i = 0$. Then, we have

$$(L \cdot H \cdot U)_{0,j} = \sum_{r=0}^n \sum_{s=0}^n L_{0,r} H_{r,s} U_{s,j} = \sum_{s=0}^n H_{0,s} U_{s,j} = H_{0,0} U_{0,j} + H_{0,1} U_{1,j} = \lambda_0 + ju,$$

and so $R_0(L \cdot H \cdot U) = R_0(A) = (\lambda_0, \lambda_0 + u, \dots, \lambda_0 + nu)$.

Assume next that $j = 0$. In this case, we obtain

$$(L \cdot H \cdot U)_{i,0} = \sum_{r=0}^n \sum_{s=0}^n L_{i,r} H_{r,s} U_{s,0} = \sum_{r=0}^n L_{i,r} H_{r,0} = \sum_{r=0}^n \binom{i}{r} \hat{\lambda}_r = \lambda_i,$$

and hence we have $C_0(L \cdot H \cdot U) = C_0(A) = (\lambda_0, \lambda_1, \dots, \lambda_n)$.

Finally, we must establish (6). Let us for the moment assume that $1 \leq i, j \leq n$. In this case, we have

$$(L \cdot H \cdot U)_{i,j} = \sum_{r=0}^n \sum_{s=0}^n L_{i,r} H_{r,s} U_{s,j} = \sum_{r=0}^n L_{i,r} H_{r,0} U_{0,j} + \sum_{r=0}^n \sum_{s=1}^n L_{i,r} H_{r,s} U_{s,j}. \quad (7)$$

Let $\Omega(i, j) = \sum_{r=0}^n \sum_{s=1}^n L_{i,r} H_{r,s} U_{s,j}$. Then, using (4), we obtain

$$\begin{aligned} \Omega(i, j) &= \sum_{r=0}^n \sum_{s=1}^n L_{i,r} H_{r,s} (U_{s-1,j-1} + U_{s,j-1}) \\ &= \sum_{r=0}^n \sum_{s=1}^n L_{i,r} H_{r,s} U_{s-1,j-1} + \sum_{r=0}^n \sum_{s=1}^n L_{i,r} H_{r,s} U_{s,j-1} \\ &= \sum_{r=1}^n \sum_{s=1}^n L_{i,r} H_{r,s} U_{s-1,j-1} + (L \cdot H \cdot U)_{i,j-1} \\ &\quad + \sum_{s=1}^n L_{i,0} H_{0,s} U_{s-1,j-1} - \sum_{r=0}^n L_{i,r} H_{r,0} U_{0,j-1} \end{aligned} \quad (8)$$

For convenience, we write $\Theta(i, j) = \sum_{r=1}^n \sum_{s=1}^n L_{i,r} H_{r,s} U_{s-1,j-1}$. Now, we apply (3), to get

$$\begin{aligned}
\Theta(i, j) &= \sum_{r=1}^n \sum_{s=1}^n (L_{i-1,r-1} + L_{i-1,r}) H_{r,s} U_{s-1,j-1} \\
&= \sum_{r=1}^n \sum_{s=1}^n L_{i-1,r-1} H_{r,s} U_{s-1,j-1} + \sum_{r=1}^n \sum_{s=1}^n L_{i-1,r} H_{r,s} U_{s-1,j-1} \\
&= \sum_{r=2}^n \sum_{s=2}^n L_{i-1,r-1} H_{r,s} U_{s-1,j-1} + \sum_{r=1}^n L_{i-1,r-1} H_{r,1} U_{0,j-1} \\
&\quad + \sum_{s=2}^n L_{i-1,0} H_{1,s} U_{s-1,j-1} + \sum_{r=1}^n \sum_{s=1}^n L_{i-1,r} H_{r,s} U_{s-1,j-1} \\
&= \sum_{r=2}^n \sum_{s=2}^n L_{i-1,r-1} H_{r,s} U_{s-1,j-1} + \sum_{r=1}^n L_{i-1,r-1} H_{r,1} U_{0,j-1} \\
&\quad + \sum_{s=2}^n L_{i-1,0} H_{1,s} U_{s-1,j-1} + \sum_{r=1}^n \sum_{s=1}^n L_{i-1,r} H_{r,s} (U_{s,j} - U_{s,j-1}) \quad (\text{by (4)}) \\
&= \sum_{r=2}^n \sum_{s=2}^n L_{i-1,r-1} H_{r-1,s-1} U_{s-1,j-1} + \sum_{r=1}^n L_{i-1,r-1} H_{r,1} U_{0,j-1} \\
&\quad + \sum_{s=2}^n L_{i-1,0} H_{1,s} U_{s-1,j-1} + \sum_{r=1}^n \sum_{s=1}^n L_{i-1,r} H_{r,s} U_{s,j} \\
&\quad - \sum_{r=1}^n \sum_{s=1}^n L_{i-1,r} H_{r,s} U_{s,j-1} \quad (\text{by the structure of } H) \\
&= \sum_{r=1}^n \sum_{s=1}^n L_{i-1,r} H_{r,s} U_{s,j-1} + \sum_{r=1}^n L_{i-1,r-1} H_{r,1} U_{0,j-1} + \sum_{s=2}^n L_{i-1,0} H_{1,s} U_{s-1,j-1} \\
&\quad + \sum_{r=1}^n \sum_{s=0}^n L_{i-1,r} H_{r,s} U_{s,j} - \sum_{r=1}^n L_{i-1,r} H_{r,0} U_{0,j} - \sum_{r=1}^n \sum_{s=1}^n L_{i-1,r} H_{r,s} U_{s,j-1} \\
&\quad (\text{note that } L_{i-1,n-1} = U_{n-1,j-1} = 0) \\
&= \sum_{r=1}^n L_{i-1,r-1} H_{r,1} U_{0,j-1} + \sum_{s=2}^n L_{i-1,0} H_{1,s} U_{s-1,j-1} + \sum_{r=0}^n \sum_{s=0}^n L_{i-1,r} H_{r,s} U_{s,j} \\
&\quad - \sum_{s=0}^n L_{i-1,0} H_{0,s} U_{s,j} - \sum_{r=1}^n L_{i-1,r} H_{r,0} U_{0,j} \\
&= \sum_{r=1}^n L_{i-1,r-1} H_{r,1} U_{0,j-1} + \sum_{s=2}^n L_{i-1,0} H_{1,s} U_{s-1,j-1} + (L \cdot H \cdot U)_{i-1,j} \\
&\quad - \sum_{s=0}^n L_{i-1,0} H_{0,s} U_{s,j} - \sum_{r=1}^n L_{i-1,r} H_{r,0} U_{0,j} \quad (\text{by (5)}).
\end{aligned}$$

By substituting this in (8), we obtain

$$\begin{aligned}
\Omega(i, j) &= (L \cdot H \cdot U)_{i, j-1} + (L \cdot H \cdot U)_{i-1, j} \\
&\quad + \sum_{r=1}^n L_{i-1, r-1} H_{r, 1} U_{0, j-1} + \sum_{s=2}^n L_{i-1, 0} H_{1, s} U_{s-1, j-1} \\
&\quad - \sum_{s=0}^n L_{i-1, 0} H_{0, s} U_{s, j} - \sum_{r=1}^n L_{i-1, r} H_{r, 0} U_{0, j} \\
&\quad + \sum_{s=1}^n L_{i, 0} H_{0, s} U_{s-1, j-1} - \sum_{r=0}^n L_{i, r} H_{r, 0} U_{0, j-1}.
\end{aligned}$$

Finally, if the above expression is substituted in (7) and the sums are put together, then we obtain

$$(L \cdot H \cdot U)_{i, j} = (L \cdot H \cdot U)_{i-1, j} + (L \cdot H \cdot U)_{i, j-1} + \Psi(i, j),$$

where

$$\begin{aligned}
\Psi(i, j) &:= \sum_{r=0}^n L_{i, r} H_{r, 0} U_{0, j} + \sum_{r=1}^n L_{i-1, r-1} H_{r, 1} U_{0, j-1} + \sum_{s=2}^n L_{i-1, 0} H_{1, s} U_{s-1, j-1} \\
&\quad - \sum_{s=0}^n L_{i-1, 0} H_{0, s} U_{s, j} - \sum_{r=1}^n L_{i-1, r} H_{r, 0} U_{0, j} + \sum_{s=1}^n L_{i, 0} H_{0, s} U_{s-1, j-1} \\
&\quad - \sum_{r=0}^n L_{i, r} H_{r, 0} U_{0, j-1}.
\end{aligned}$$

However, by easy calculations one can show that

$$\begin{aligned}
\sum_{r=0}^n L_{i, r} H_{r, 0} U_{0, j} - \sum_{r=0}^n L_{i, r} H_{r, 0} U_{0, j-1} &= 0, \\
\sum_{r=1}^n L_{i-1, r-1} H_{r, 1} U_{0, j-1} &= \sum_{r=1}^n \binom{i-1}{r-1} \hat{\mu}_r = \sum_{r=1}^n \left(\binom{i}{r} - \binom{i-1}{r} \right) \hat{\mu}_r = \mu_i - \mu_{i-1}, \\
\sum_{r=1}^n L_{i-1, r} H_{r, 0} U_{0, j} &= \sum_{r=0}^n \hat{\lambda}_r - \lambda_0 = \lambda_{i-1} - \lambda_0, \\
\sum_{s=2}^n L_{i-1, 0} H_{1, s} U_{s-1, j-1} &= (j-1)v, \\
\sum_{s=0}^n L_{i-1, 0} H_{0, s} U_{s, j} &= \lambda_0 + ju, \\
\sum_{s=1}^n L_{i, 0} H_{0, s} U_{s-1, j-1} &= u,
\end{aligned}$$

and so

$$\Psi(i, j) = \mu_i - \mu_{i-1} - \lambda_{i-1} + (j-1)(v-u).$$

This completes the proof. \square

Before stating the next result, we need to introduce some additional definitions. Let $\lambda = (\lambda_i)_{i \geq 0}$ and $\mu = (\mu_i)_{i \geq 0}$ be two arbitrary sequences. The *convolution* of λ and μ is the sequence $\nu = (\nu_i)_{i \geq 0}$, where

$$\nu_i = \sum_{k=0}^i \lambda_k \mu_{i-k}.$$

The *convolution matrix* associated with sequences λ and μ is the infinite matrix $A(\infty)$ whose first column $C_0(A(\infty))$ is λ and whose j th column ($j = 1, 2, \dots$) is the convolution of sequences $C_{j-1}(A(\infty))$ and μ . We say that the convolution matrix of the sequences λ and λ is the convolution matrix of the sequence λ . There are many well-known integer matrices which can be written as convolution matrices of some sequences. For instance, $U(\infty)$ is the convolution matrix of the sequences $(1, 0, 0, \dots)$ and $(1, 1, 0, 0, \dots)$ and $P_{(1,1,\dots),(1,1,\dots)}(\infty)$ is the convolution matrix of the sequence $(1, 1, \dots)$.

We will need the following technical result [4, Theorem 3.1].

Proposition 2. *Let*

$$A(x) = \sum_{n=1}^{\infty} a_n x^{n-1}, \quad B(x) = \sum_{n=0}^{\infty} b_n x^n, \quad V(x) = \sum_{n=0}^{\infty} v_n x^n \quad \text{and} \quad W(x) = \sum_{n=0}^{\infty} w_n x^n$$

be the generating functions for the sequences $(a_n)_{n \geq 1}$, $(b_n)_{n \geq 0}$, $(v_n)_{n \geq 0}$, and $(w_n)_{n \geq 0}$, respectively. Consider an infinite dimensional matrix of the following form:

$$M(\infty) = \left(\begin{array}{c|ccc} b_0 & v_0 & v_0 w_0 & \cdots \\ b_1 & v_1 & v_0 w_1 + v_1 w_0 & \cdots \\ b_2 & v_2 & v_0 w_2 + v_1 w_1 + v_2 w_0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{array} \right)$$

where $C_0(M(\infty)) = (b_0, b_1, \dots)^t$ and $M(\infty)^{[1]}$ is the convolution matrix of the sequences $(v_i)_{i \geq 0}$ and $(w_j)_{j \geq 0}$. If

$$A(W(x)) = B(x)/V(x), \tag{9}$$

then for any non-negative integer n , there holds

$$\det(M(n)) = (-1)^n v_0^{n+1} w_1^{n(n+1)/2} a_{n+1},$$

where $M(n)$ is the $(n+1) \times (n+1)$ upper left corner matrix of $M(\infty)$.

We are now in a position to prove the following theorem which is the second result of this paper.

Theorem 3. *Let $A(n)$ be defined as in Theorem 1 and let c be a constant. In the case when $u = v = 1$ and $\lambda_i = (2^i - 1)c + 1$, we have the following statements:*

- (a) if $\mu_i = \left(2^i + \frac{(i-2)(i+1)}{2}\right)c - \frac{i(i-3)}{2}$, then $\det(A(n)) = F_{n+1}$.
- (b) if $\mu_i = \left(\frac{5 \cdot 3^i}{4} - 2^i - \frac{2i+1}{4}\right)c + \frac{5(3^i-1)}{4} + \frac{i}{2}$, then $\det(A(n)) = L_{n+1}$.
- (c) if $\mu_i = i^2c - i^2 + 2i$, then $\det(A(n)) = J_{n+1}$.
- (d) if $\mu_i = \left(2^{i+1} + \frac{(i+1)(i-4)}{2}\right)c + \frac{(5-i)i}{2}$, then $\det(A(n)) = P_{n+1}$.

Proof. Let $\mu = (\mu_i)_{i \geq 0}$ be a sequence with $\mu_0 = 0$ and let c be a constant. Let $\lambda = (\lambda_i)_{i \geq 0}$ be a sequence with $\lambda_i = (2^i - 1)c + 1$. We consider the infinite matrices $A(\infty) = [A_{i,j}]_{i,j \geq 0}$ whose entries satisfy

$$A_{i,j} = A_{i-1,j} + A_{i,j-1} - (2^i - 1)c - 1 + \mu_i - \mu_{i-1} \quad \text{for } i, j \geq 1, \quad (10)$$

with the initial conditions $A_{i,0} = (2^i - 1)c + 1$ and $A_{0,i} = 1 + i$, $i \geq 0$. By Theorem 2, we observe that

$$A(n) = L(n) \cdot H(n) \cdot U(n),$$

where

$$H(n) = \left(\begin{array}{c|cccc} 1 & 1 & 0 & \cdots & 0 \\ \hline c & & & & \\ c & & & & \\ \vdots & & & & \\ c & & & & \end{array} T_{(\hat{\mu}_1, \hat{\mu}_2, \hat{\mu}_3, \dots), (\hat{\mu}_1, v, 0, 0, \dots)}(n-1) \right).$$

Evidently $\det(A(n)) = \det(H(n))$, so it suffices to find $\det(H(n))$. From the structure of matrix $H(\infty)$, we have

$$C_0(H(\infty)) = (b_i)_{i \geq 0} = (1, c, c, \dots)^t,$$

whose generating function is

$$B(x) = \frac{1 + (c-1)x}{1-x}.$$

- (a) Let $\mu_i = \left(2^i + \frac{(i-2)(i+1)}{2}\right)c - \frac{i(i-3)}{2}$. In this case, we have the following infinite dimensional matrices:

$$A(\infty) = \begin{pmatrix} 1 & 2 & 3 & 4 & \cdots \\ c+1 & 2c+3 & 3c+6 & 4c+10 & \cdots \\ 3c+1 & 7c+3 & 12c+8 & 18c+17 & \cdots \\ 7c+1 & 17c+2 & 32c+8 & 53c+23 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

and

$$H(\infty) = \left(\begin{array}{c|cccc} 1 & 1 & 0 & \cdots & 0 \\ \hline c & & & & \\ c & & T_{(c+1, 2c-1, c, c, \dots), (c+1, 1, 0, 0, \dots)}(\infty) & & \\ \vdots & & & & \end{array} \right).$$

Note that the submatrix $H(\infty)^{[1]}$ is the convolution of sequences

$$(v_i)_{i \geq 0} = (1, c+1, 2c-1, c, c, \dots), \quad \text{and} \quad (w_i)_{i \geq 0} = (0, 1, 0, 0, 0, \dots),$$

whose generating functions are

$$V(x) = \frac{1 + cx + (c-2)x^2 - (c-1)x^3}{1-x} \quad \text{and} \quad W(x) = x,$$

respectively. Plugging these generating functions into (9) yields

$$A(W(x)) = A(x) = \frac{\frac{1+(c-1)x}{1-x}}{\frac{1+cx+(c-2)x^2-(c-1)x^3}{1-x}} = 1 - x + 2x^2 - \cdots + (-1)^n F_{n+1} x^n + \cdots,$$

and it follows by Proposition 2 that

$$\det(H(n)) = (-1)^n v_0^{n+1} w_1^{n(n+1)/2} a_{n+1} = (-1)^n a_{n+1} = F_{n+1},$$

as required.

- (b) Let $\mu_i = \left(\frac{5 \cdot 3^i}{4} - 2^i - \frac{2i+1}{4} \right) c + \frac{5(3^i-1)}{4} + \frac{i}{2}$. The infinite dimensional matrices created in this case are as follows:

$$A(\infty) = \begin{pmatrix} 1 & 2 & 3 & 4 & \cdots \\ c+1 & 2c+5 & 3c+10 & 4c+16 & \cdots \\ 3c+1 & 9c+13 & 16c+30 & 24c+53 & \cdots \\ 7c+1 & 31c+36 & 62c+88 & 101c+163 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

and

$$H(\infty) = \left(\begin{array}{c|cccc} 1 & 1 & 0 & \cdots & 0 \\ \hline c & & & & \\ c & & T_{(c+3, 4c+5, 9c+10, 19c+20, \dots), (c+3, 1, 0, 0, \dots)}(\infty) & & \\ \vdots & & & & \end{array} \right).$$

Again, one can easily see that the submatrix $H(\infty)^{[1]}$ is the convolution of sequences

$$(v_i)_{i \geq 0} = (1, c+3, 4c+5, 9c+10, 19c+20, \dots),$$

(with general form $v_0 = 1$, $v_1 = c + 3$ and $v_i = (5 \cdot 2^{i-2})(c + 1) - c$ for $i \geq 2$), and

$$(w_i)_{i \geq 0} = (0, 1, 0, 0, 0, \dots).$$

The generating functions for these sequences are

$$V(x) = \frac{(1 + (c-1)x)(-x^2 + x + 1)}{(1-x)(1-2x)}, \quad \text{and} \quad W(x) = x,$$

respectively. If $B(x)$, $V(x)$ and $W(x)$ are substituted in (9), then we obtain

$$A(W(x)) = A(x) = \frac{\frac{1+(c-1)x}{1-x}}{\frac{(1+(c-1)x)(-x^2+x+1)}{(1-x)(1-2x)}} = 1 - 3x + 4x^2 - 7x^3 + \dots + (-1)^n L_{n+1} x^n + \dots,$$

and by Proposition 2, it follows that

$$\det(H(n)) = (-1)^n v_0^{n+1} w_1^{n(n+1)/2} a_{n+1} = (-1)^n a_{n+1} = L_{n+1},$$

as required.

(c) Let $\mu_i = i^2 c - i^2 + 2i$. In this case, we have the following infinite dimensional matrices:

$$A(\infty) = \begin{pmatrix} 1 & 2 & 3 & 4 & \dots \\ c+1 & 2c+3 & 3c+6 & 4c+10 & \dots \\ 3c+1 & 7c+2 & 12c+6 & 18c+14 & \dots \\ 7c+1 & 16c-1 & 30c+1 & 50c+11 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

and

$$H(\infty) = \left(\begin{array}{c|cccc} 1 & 1 & 0 & \dots & 0 \\ \hline c & & & & \\ c & & & & \\ \vdots & & & & \end{array} T_{(c+1, 2c-2, 0, 0, \dots), (c+1, 1, 0, 0, \dots)}(\infty) \right).$$

Moreover, from the structure of $H(\infty)$, we see that the submatrix $H(\infty)^{[1]}$ is the convolution of sequences

$$(v_i)_{i \geq 0} = (1, c+1, 2c-2, 0, 0, \dots), \quad \text{and} \quad (w_i)_{i \geq 0} = (0, 1, 0, 0, \dots),$$

with generating functions $V(x) = 1 + (c+1)x + (2c-2)x^2$ and $W(x) = x$, respectively. Substituting the obtained generating functions in (9), we obtain

$$A(W(x)) = A(x) = \frac{\frac{1+(c-1)x}{1-x}}{1 + (c+1)x + (2c-2)x^2} = 1 - x + 3x^2 - \dots + (-1)^n J_{n+1} x^n + \dots.$$

Therefore, it follows from Proposition 2 that

$$\det(H(n)) = (-1)^n v_0^{n+1} w_1^{n(n+1)/2} a_{n+1} = (-1)^n a_{n+1} = J_{n+1},$$

as required.

- (d) Let $\mu_i = \left(2^{i+1} + \frac{(i+1)(i-4)}{2}\right) c + \frac{(5-i)i}{2}$. This time, we will deal with the following matrices:

$$A(\infty) = \begin{pmatrix} 1 & 2 & 3 & 4 & \cdots \\ c+1 & 2c+4 & 3c+8 & 4c+13 & \cdots \\ 3c+1 & 8c+5 & 14c+13 & 21c+26 & \cdots \\ 7c+1 & 21c+5 & 41c+17 & 68c+42 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

and

$$H(\infty) = \left(\begin{array}{c|ccc} 1 & 1 & 0 & \cdots & 0 \\ \hline c & & & & \\ c & & T_{(c+2, 3c-1, 2c, 2c, \dots), (c+2, 1, 0, 0, \dots)}(\infty) & & \\ \vdots & & & & \end{array} \right).$$

In addition, the submatrix $H(\infty)^{[1]}$ of $H(\infty)$ is the convolution of sequences:

$$(v_i)_{i \geq 0} = (1, c+2, 3c-1, 2c, 2c, \dots) \quad \text{and} \quad (w_i)_{i \geq 0} = (0, 1, 0, 0, \dots).$$

Note that the generating functions of these sequences are

$$V(x) = \frac{1 + (1+c)x + (2c-3)x^2 - (c-1)x^3}{1-x} \quad \text{and} \quad W(x) = x,$$

respectively. After having substituted these generating functions in (9), we obtain

$$A(W(x)) = A(x) = \frac{\frac{1+(c-1)x}{1-x}}{\frac{1+(1+c)x+(2c-3)x^2-(c-1)x^3}{1-x}} = 1 - 2x + 5x^2 - \cdots + (-1)^n P_{n+1} x^n + \cdots.$$

Now, by Proposition 2, we deduce that

$$\det(H(n)) = (-1)^n v_0^{n+1} w_1^{n(n+1)/2} a_{n+1} = (-1)^n a_{n+1} = P_{n+1},$$

as required.

This completes the proof. □

3 Some remarks

In this section, we will explain how the sequences $(\lambda_i)_{i \geq 0}$ and $(\mu_i)_{i \geq 0}$ in Theorem 3, are determined. Consider the following lower Hessenberg matrix

$$H(\infty) = [H_{i,j}]_{i,j \geq 0} = \begin{pmatrix} h_{0,0} & h_{0,1} & 0 & 0 & 0 & \cdots \\ h_{1,0} & h_{1,1} & h_{1,2} & 0 & 0 & \cdots \\ h_{2,0} & h_{2,1} & h_{1,1} & h_{1,2} & 0 & \cdots \\ h_{3,0} & h_{3,1} & h_{2,1} & h_{1,1} & h_{1,2} & \cdots \\ h_{4,0} & h_{4,1} & h_{3,1} & h_{2,1} & h_{1,1} & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Let $H(n) = [H_{i,j}]_{0 \leq i,j \leq n}$, and let d_n be the n th determinant of $H(n)$. In what follows, we show that the sequence of principal minors of $H(\infty)$, i.e., $D(H(\infty)) = (d_n)_{n \geq 0}$, satisfies a recurrence relation.

Proposition 4. *With the above notation, we have*

$$d_n = \begin{cases} h_{0,0}, & \text{if } n = 0, \\ (-1)^n h_{0,1} (h_{1,2})^{n-1} h_{n,0} + \sum_{k=0}^{n-1} h_{n-k,1} (-h_{1,2})^{n-k-1} d_k, & \text{if } n \geq 1. \end{cases}$$

Proof. Obviously, $d_0 = h_{0,0}$. Hence, from now on we assume $n > 1$. First, we apply the following row operations:

$$\begin{aligned} H_1(n) &= \left(\prod_{i=1}^n O_{i,0} \left(\frac{-h_{i,1}}{h_{0,1}} \right) \right) H(n), \\ H_2(n) &= \left(\prod_{i=1}^{n-1} O_{i+1,1} \left(\frac{-h_{i,1}}{h_{1,2}} \right) \right) H_1(n), \\ H_3(n) &= \left(\prod_{i=1}^{n-2} O_{i+2,2} \left(\frac{-h_{i,1}}{h_{1,2}} \right) \right) H_2(n), \\ &\vdots \\ H_n(n) &= \left(\prod_{i=1}^1 O_{i+(n-1),n-1} \left(\frac{-h_{i,1}}{h_{1,2}} \right) \right) H_{n-1}(n). \end{aligned}$$

It is obvious that, step by step, the columns are “emptied” until finally the following matrix

$$H_n(n) = \begin{pmatrix} \tilde{h}_{0,0} & h_{0,1} & 0 & 0 & 0 & \cdots & 0 \\ \tilde{h}_{1,0} & 0 & h_{1,2} & 0 & 0 & \cdots & 0 \\ \tilde{h}_{2,0} & 0 & 0 & h_{1,2} & 0 & \cdots & 0 \\ \tilde{h}_{3,0} & 0 & 0 & 0 & h_{1,2} & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \tilde{h}_{n-1,0} & 0 & 0 & 0 & 0 & \cdots & h_{1,2} \\ \tilde{h}_{n,0} & 0 & 0 & 0 & 0 & \cdots & 0 \end{pmatrix}_{(n+1) \times (n+1)},$$

is obtained, where

$$\tilde{h}_{i,0} = \begin{cases} h_{0,0}, & \text{if } i = 0; \\ h_{1,0} - \frac{h_{1,1}}{h_{0,1}}h_{0,0}, & \text{if } i = 1; \\ h_{i,0} - \frac{h_{i,1}}{h_{0,1}}h_{0,0} - \frac{1}{h_{1,2}} \sum_{k=1}^{i-1} h_{i-k,1} \tilde{h}_{k,0}, & \text{if } i \geq 2. \end{cases} \quad (11)$$

Evidently, $d_n = \det(H_n(n))$. Expanding the determinant along the last row of $\det(H_n(n))$, we obtain

$$d_n = (-1)^n \tilde{h}_{n,0} h_{0,1} (h_{1,2})^{n-1}, \quad (n \geq 1). \quad (12)$$

Finally, after some simplification, it follows that

$$\begin{aligned} d_n &= (-1)^n \tilde{h}_{n,0} h_{0,1} (h_{1,2})^{n-1} \\ &= (-1)^n h_{0,1} (h_{1,2})^{n-1} \left[h_{n,0} - \frac{h_{n,1}}{h_{0,1}} h_{0,0} - \frac{1}{h_{1,2}} \sum_{k=1}^{n-1} h_{n-k,1} \tilde{h}_{k,0} \right] \quad (\text{by (11)}) \\ &= (-1)^n h_{0,1} (h_{1,2})^{n-1} h_{n,0} + (-1)^{n+1} (h_{1,2})^{n-1} h_{n,1} h_{0,0} \\ &\quad + (-1)^{n+1} h_{0,1} (h_{1,2})^{n-2} \sum_{k=1}^{n-1} h_{n-k,1} \tilde{h}_{k,0} \\ &= (-1)^n h_{0,1} (h_{1,2})^{n-1} h_{n,0} + (-1)^{n+1} (h_{1,2})^{n-1} h_{n,1} h_{0,0} \\ &\quad + \sum_{k=1}^{n-1} h_{n-k,1} (-h_{1,2})^{n-k-1} d_k \quad (\text{by (12)}) \\ &= (-1)^n h_{0,1} (h_{1,2})^{n-1} h_{n,0} + \sum_{k=0}^{n-1} h_{n-k,1} (-h_{1,2})^{n-k-1} d_k. \end{aligned}$$

and the result follows. \square

In Proposition 4, if we take $h_{0,0} = h_{0,1} = 1$, $h_{1,2} = 1$, $h_{i,0} = c$ and $h_{i,1} = \hat{\mu}_i$ for $i \geq 1$, then we obtain

$$d_n = \begin{cases} 1, & \text{if } n = 0; \\ (-1)^n c + \sum_{k=0}^{n-1} \hat{\mu}_{n-k} (-1)^{n-k-1} d_k, & \text{if } n \geq 1. \end{cases}$$

Now, if $(d_n)_{n \geq 0} \in \{\mathcal{F}, \mathcal{L}, \mathcal{J}, \mathcal{P}\}$, then

$$\hat{\mu}_n = c + (-1)^{n-1} d_n + \sum_{k=1}^{n-1} (-1)^{k+1} \hat{\mu}_{n-k} d_k,$$

from which we determine the sequence $(\hat{\mu}_i)_{i \geq 1}$. Now, we form

$$H(\infty) = \left(\begin{array}{c|ccc} 1 & 1 & 0 & \dots \\ \hline c & & & \\ c & & & \\ \vdots & & & \\ c & & & \end{array} T_{(\hat{\mu}_1, \hat{\mu}_2, \hat{\mu}_3, \dots), (\hat{\mu}_1, 1, 0, 0, \dots)}(\infty) \right).$$

Finally, the sequences $(\lambda_i)_{i \geq 0}$ and $(\mu_i)_{i \geq 0}$ are determined by the equation $A(n) = L(n) \cdot H(n) \cdot U(n)$.

4 Acknowledgement

Our special thanks go to the Research Institute for Fundamental Sciences, Tabriz, Iran, for having sponsored this paper.

References

- [1] A. R. Moghaddamfar and S. M. H. Pooya, Generalized Pascal triangles and Toeplitz matrices, *Electron. J. Linear Algebra*, **18** (2009), 564–588.
- [2] A. R. Moghaddamfar and H. Tajbakhsh, Lucas numbers and determinants, *Integers*, **12** (2012), 21–51.
- [3] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*. Published electronically at <http://oeis.org>, 2013.
- [4] Y. Yang and M. Leonard, Evaluating determinants of convolution-like matrices via generating functions, *Int. J. Inf. Syst. Sci.*, **3** (2007), 569–580.

2010 *Mathematics Subject Classification*: Primary 11C20; Secondary 15B36, 10A35.

Keywords: determinant, generalized Pascal triangle, Toeplitz matrix, matrix factorization, convolution matrix, lower Hessenberg matrix.

(Concerned with sequences [A000032](#), [A000045](#), [A000129](#), and [A001045](#).)

Received December 3 2013; revised version received March 2 2014. Published in *Journal of Integer Sequences*, March 26 2014.

Return to [Journal of Integer Sequences home page](#).