



Counting Toroidal Binary Arrays, II

S. N. Ethier¹

Department of Mathematics
University of Utah
155 South 1400 East
Salt Lake City, UT 84112
USA

ethier@math.utah.edu

Jiyeon Lee²

Department of Statistics
Yeungnam University
214-1 Daedong, Kyeongsan
Kyeongbuk 712-749
South Korea

leejy@yu.ac.kr

Abstract

We derive formulas for (i) the number of distinct toroidal $n \times n$ binary arrays, allowing rotation of rows and/or columns as well as matrix transposition, and (ii) the number of distinct toroidal $n \times n$ binary arrays, allowing rotation and/or reflection of rows and/or columns as well as matrix transposition.

¹Partially supported by a grant from the Simons Foundation (209632).

²Supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future Planning (No. 2013R1A1A3A04007670).

1 Introduction

A previous paper [1] found the number of (distinct) toroidal $m \times n$ binary arrays, allowing rotation of rows and/or columns, to be

$$a(m, n) := \frac{1}{mn} \sum_{c|m} \sum_{d|n} \varphi(c)\varphi(d) 2^{mn/\text{lcm}(c,d)}, \quad (1)$$

where φ is Euler's phi function and lcm stands for least common multiple. This is [A184271](#) in the *On-Line Encyclopedia of Integer Sequences* [2]. The main diagonal is [A179043](#). It was also shown that, allowing rotation and/or reflection of rows and/or columns, the number becomes

$$b(m, n) := b_1(m, n) + b_2(m, n) + b_3(m, n) + b_4(m, n), \quad (2)$$

where

$$b_1(m, n) := \frac{1}{4mn} \sum_{c|m} \sum_{d|n} \varphi(c)\varphi(d) 2^{mn/\text{lcm}(c,d)},$$

$$b_2(m, n) := \frac{1}{4n} \sum_{d|n} \varphi(d) 2^{mn/d} + \begin{cases} (4n)^{-1} \sum' \varphi(d) (2^{(m+1)n/(2d)} - 2^{mn/d}), & \text{if } m \text{ is odd;} \\ (8n)^{-1} \sum' \varphi(d) (2^{mn/(2d)} + 2^{(m+2)n/(2d)} - 2 \cdot 2^{mn/d}), & \text{if } m \text{ is even,} \end{cases}$$

with $\sum' := \sum_{d|n: d \text{ is odd}}$

$$b_3(m, n) := b_2(n, m),$$

and

$$b_4(m, n) := \begin{cases} 2^{(mn-3)/2}, & \text{if } m \text{ and } n \text{ are odd;} \\ 3 \cdot 2^{mn/2-3}, & \text{if } m \text{ and } n \text{ have opposite parity;} \\ 7 \cdot 2^{mn/2-4}, & \text{if } m \text{ and } n \text{ are even.} \end{cases}$$

(The formula for $b_2(m, n)$ given in [1] is simplified here.) This is [A222188](#) in the *OEIS* [2]. The main diagonal is [A209251](#).

Our aim here is to derive the corresponding formulas when $m = n$ and we allow matrix transposition as well. More precisely, we show that the number of (distinct) toroidal $n \times n$ binary arrays, allowing rotation of rows and/or columns as well as matrix transposition, is

$$\alpha(n) = \frac{1}{2} a(n, n) + \frac{1}{2n} \sum_{d|n} \varphi(d) 2^{n(n+d-2\lfloor d/2 \rfloor)/(2d)}, \quad (3)$$

where $a(n, n)$ is from (1). When we allow rotation and/or reflection of rows and/or columns as well as matrix transposition, the number becomes

$$\beta(n) = \frac{1}{2} b(n, n) + \frac{1}{4n} \sum_{d|n} \varphi(d) 2^{n(n+d-2\lfloor d/2 \rfloor)/(2d)} + \begin{cases} 2^{(n^2-5)/4}, & \text{if } n \text{ is odd;} \\ 5 \cdot 2^{n^2/4-3}, & \text{if } n \text{ is even,} \end{cases} \quad (4)$$

where $b(n, n)$ is from (2). These are the sequences [A255015](#) and [A255016](#), respectively, recently added to the *OEIS* [2].

For an alternative description, we could define a group action on the set of $n \times n$ binary arrays, which has 2^{n^2} elements. If the group is generated by σ (row rotation) and τ (column rotation), then the number of orbits is given by $a(n, n)$; see [1]. If the group is generated by σ , τ , and ζ (matrix transposition), then the number of orbits is given by $\alpha(n)$; see Theorem 1 below. If the group is generated by σ , τ , ρ (row reflection), and θ (column reflection), then the number of orbits is given by $b(n, n)$; see [1]. If the group is generated by σ , τ , ρ , θ , and ζ , then the number of orbits is given by $\beta(n)$; see Theorem 2 below.

Both theorems are proved using Pólya's enumeration theorem (actually, the simplified unweighted version; see, e.g., van Lint and Wilson [3, Theorem 37.1, p. 524]).

To help clarify the distinction between the various group actions, we consider the case of 3×3 binary arrays as in [1]. When the group is generated by σ and τ (allowing rotation of rows and/or columns), there are 64 orbits, which were listed in [1]. When the group is generated by σ , τ , and ζ (allowing rotation of rows and/or columns as well as matrix transposition), there are 44 orbits, which are listed in Table 1 below. When the group is generated by σ , τ , ρ , and θ (allowing rotation and/or reflection of rows and/or columns), there are 36 orbits, which were listed in [1]. When the group is generated by σ , τ , ρ , θ , and ζ (allowing rotation and/or reflection of rows and/or columns as well as matrix transposition), there are 26 orbits, which are listed in Table 2 below.

Table 3 provides numerical values for $\alpha(n)$ and $\beta(n)$ for small n .

We take this opportunity to correct a small gap in the proof of Theorem 2 in [1]. The proof assumed implicitly that $m, n \geq 3$. The theorem is correct as stated for $m, n \geq 1$, so the proof is incomplete if m or n is 1 or 2. Following the proof of Theorem 2 below, we supply the missing steps.

2 Rotation of rows and columns, and matrix transposition

Let $X_n := \{0, 1\}^{\{0, 1, \dots, n-1\}^2}$ be the set of $n \times n$ matrices of 0s and 1s, which has 2^{n^2} elements. Let $\alpha(n)$ denote the number of orbits of the group action on X_n by the group of order $2n^2$ generated by σ (row rotation), τ (column rotation), and ζ (matrix transposition). (Exception: If $n = 1$, the group is of order 1.)

Informally, $\alpha(n)$ is the number of (distinct) toroidal $n \times n$ binary arrays, allowing rotation of rows and/or columns as well as matrix transposition.

Theorem 1. *With $a(n, n)$ defined using (1), $\alpha(n)$ is given by (3).*

Proof. Let us assume that $n \geq 2$. By Pólya's enumeration theorem,

$$\alpha(n) = \frac{1}{2n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (2^{A_{ij}} + 2^{E_{ij}}), \quad (5)$$

Table 1: A list of the 44 orbits of the group action in which the group generated by σ , τ , and ζ acts on the set of 3×3 binary arrays. (Rows and/or columns can be rotated and matrices can be transposed.) Each orbit is represented by its minimal element in 9-bit binary form. Subscripts indicate orbit size. Bars separate different numbers of 1s.

$$\begin{aligned}
& \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}_1 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}_9 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}_9 \mid \\
& \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 1 & 1 \end{pmatrix}_6 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}_9 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 0 \end{pmatrix}_9 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}_3 \\
& \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}_3 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}_9 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}_9 \\
& \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}_9 \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{pmatrix}_9 \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix}_{18} \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}_9 \\
& \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix}_9 \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}_9 \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \end{pmatrix}_9 \mid \\
& \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_6 \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_9 \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 0 & 1 \end{pmatrix}_9 \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix}_{18} \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}_3 \\
& \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}_3 \mid \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{18} \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}_9 \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_9 \mid \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_1
\end{aligned}$$

where A_{ij} (resp., E_{ij}) is the number of cycles in the permutation $\sigma^i \tau^j$ (resp., $\sigma^i \tau^j \zeta$); here σ rotates the rows (row 0 becomes row 1, row 1 becomes row 2, ..., row $n-1$ becomes row 0), τ rotates the columns, and ζ transposes the matrix. We know from [1] that

$$a(n, n) = \frac{1}{n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{A_{ij}}, \quad (6)$$

so it remains to find E_{ij} . The permutation ζ has n fixed points and $\binom{n}{2}$ transpositions, so $E_{00} = n(n+1)/2$.

Notice that σ and τ commute, whereas $\sigma\zeta = \zeta\tau$ and $\tau\zeta = \zeta\sigma$. Let $(i, j) \in \{0, 1, \dots, n-$

Table 2: A list of the 26 orbits of the group action in which the group generated by σ , τ , ρ , θ , and ζ acts on the set of 3×3 binary arrays. (Rows and/or columns can be rotated and/or reflected and matrices can be transposed.) Each orbit is represented by its minimal element in 9-bit binary form. Subscripts indicate orbit size. Bars separate different numbers of 1s.

$$\begin{aligned}
 & \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}_1 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix}_{18} \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}_{18} \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 1 & 1 \end{pmatrix}_6 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}_{36} \\
 & \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}_6 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}_{36} \mid \\
 & \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_6 \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{36} \\
 & \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}_6 \mid \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{18} \mid \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{18} \mid \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_9 \mid \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_1
 \end{aligned}$$

Table 3: The values of $\alpha(n)$ and $\beta(n)$ for $n = 1, 2, \dots, 12$.

| n | $\alpha(n)$ | $\beta(n)$ |
|-----|---|---|
| 1 | 2 | 2 |
| 2 | 6 | 6 |
| 3 | 44 | 26 |
| 4 | 2209 | 805 |
| 5 | 674384 | 172112 |
| 6 | 954623404 | 239123150 |
| 7 | 5744406453840 | 1436120190288 |
| 8 | 144115192471496836 | 36028817512382026 |
| 9 | 14925010120653819583840 | 3731252531904348833632 |
| 10 | 6338253001142965335834871200 | 1584563250300891724601560272 |
| 11 | 10985355337065423791175013899922368 | 2746338834266358751489231123956672 |
| 12 | 77433143050453552587418968170813573149024 | 19358285762613388352671214587818634041520 |

$1\}^2 - \{(0, 0)\}$ be arbitrary. Then

$$(\sigma^i \tau^j \zeta)^2 = (\sigma^i \tau^j \zeta)(\zeta \tau^i \sigma^j) = \sigma^{i+j} \tau^{i+j},$$

hence

$$\begin{aligned}(\sigma^i \tau^j \zeta)^{2d} &= \sigma^{(i+j)d} \tau^{(i+j)d} = ((\sigma\tau)^{i+j})^d, \\(\sigma^i \tau^j \zeta)^{2d+1} &= \sigma^{(i+j)d+i} \tau^{(i+j)d+j} \zeta.\end{aligned}$$

Clearly, $(\sigma^i \tau^j \zeta)^{2d+1}$ cannot be the identity permutation, so $\sigma^i \tau^j \zeta$ is of even order. Using the fact that, in the cyclic group $\{a, a^2, \dots, a^{n-1}, a^n = e\}$ of order n , a^k is of order $n/\gcd(k, n)$, we find that the permutation $\sigma^i \tau^j \zeta$ is of order $2d$, where $d := n/\gcd(i+j, n)$. Therefore, every cycle of this permutation must have length that divides $2d$.

We claim that all cycles have length d or $2d$. Accepting that for now, let us determine how many cycles have length d . A cycle that includes entry (k, l) has length d if (k, l) is a fixed point of $(\sigma^i \tau^j \zeta)^d$. For this to hold we must have d odd (otherwise there would be no fixed points because we have excluded the case $i = j = 0$ and $(i+j)d/2 = \text{lcm}(i+j, n)/2$ is not a multiple of n). Since

$$(\sigma^i \tau^j \zeta)^d = \sigma^{(i+j)(d-1)/2+i} \tau^{(i+j)(d-1)/2+j} \zeta,$$

we must also have

$$(k, l) = ([l + (i+j)(d-1)/2 + j], [k + (i+j)(d-1)/2 + i]), \quad (7)$$

where $d := n/\gcd(i+j, n)$ and, for simplicity, $[r] := (r \bmod n) \in \{0, 1, \dots, n-1\}$. For each $k \in \{0, 1, \dots, n-1\}$, there is a unique l (namely, $l := [k + (i+j)(d-1)/2 + i]$) such that (7) holds; indeed,

$$\begin{aligned}[l + (i+j)(d-1)/2 + j] &= [[k + (i+j)(d-1)/2 + i] + (i+j)(d-1)/2 + j] \\&= [k + (i+j)(d-1)/2 + i + (i+j)(d-1)/2 + j] \\&= [k + (i+j)d] \\&= [k + (i+j)(n/\gcd(i+j, n))] \\&= [k + \text{lcm}(i+j, n)] \\&= k.\end{aligned}$$

This shows that there are n fixed points of $(\sigma^i \tau^j \zeta)^d$. Each cycle of length d of $\sigma^i \tau^j \zeta$ will account for d such fixed points, hence there are n/d such cycles. All remaining cycles will have length $2d$, and so there are $n(n-1)/(2d)$ of these. The total number of cycles is therefore $n(n+1)/(2d)$.

The other possibility is that d is even and all cycles have the same length, $2d$, so there are $n^2/(2d)$ of them. Notice that d is a divisor of n , so the contribution to

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{E_{ij}}$$

from odd d is

$$\sum_{d|n: d \text{ is odd}} n\varphi(d)2^{n(n+1)/(2d)} \quad (8)$$

and from even d is

$$\sum_{d|n: d \text{ is even}} n\varphi(d)2^{n^2/(2d)}. \quad (9)$$

The reason for the coefficient $n\varphi(d)$ is that, if $d|n$, then the number of elements of the cyclic group $\{e, \sigma\tau, (\sigma\tau)^2, \dots, (\sigma\tau)^{n-1}\}$ that are of order d is $\varphi(d)$. And for a given $(i, j) \in \{0, 1, \dots, n-1\}^2$, there are n pairs $(k, l) \in \{0, 1, \dots, n-1\}^2$ such that $[k+l] = [i+j]$. Putting (8) and (9) together, we obtain

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{E_{ij}} = \sum_{d|n} n\varphi(d)2^{n(n+d-2\lfloor d/2 \rfloor)/(2d)}, \quad (10)$$

which, together with (5) and (6), yields (3).

It remains to prove our claim that, for $(i, j) \in \{0, 1, \dots, n-1\}^2 - \{(0, 0)\}$, the permutation $\sigma^i \tau^j \zeta$ cannot have any cycles whose length is a proper divisor of $d := n/\gcd(i+j, n)$. Let $c|d$ with $1 \leq c < d$. We must show that $(\sigma^i \tau^j \zeta)^c$ has no fixed points. We can argue as above with c in place of d . For (k, l) to be a fixed point of $(\sigma^i \tau^j \zeta)^c$ we must have $(i+j)c$ a multiple of n . But $d := n/\gcd(i+j, n)$ is the smallest integer c such that $(i+j)c$ is a multiple of n because $(i+j)n/\gcd(i+j, n) = \text{lcm}(i+j, n)$.

Finally, we excluded the case $n = 1$ at the beginning of the proof, but we notice that the formula (3) gives $\alpha(1) = 2$, which is correct. \square

3 Rotation and reflection of rows and columns, and matrix transposition

Let $X_n := \{0, 1\}^{\{0, 1, \dots, n-1\}^2}$ be the set of $n \times n$ matrices of 0s and 1s, which has 2^{n^2} elements. Let $\beta(n)$ denote the number of orbits of the group action on X_n by the group of order $8n^2$ generated by σ (row rotation), τ (column rotation), ρ (row reflection), θ (column reflection), and ζ (matrix transposition). (Exceptions: If $n = 2$, the group is of order 8; if $n = 1$, the group is of order 1.)

Informally, $\beta(n)$ is the number of (distinct) toroidal $n \times n$ binary arrays, allowing rotation and/or reflection of rows and/or columns as well as matrix transposition.

Theorem 2. *With $b(n, n)$ defined using (2), $\beta(n)$ is given by (4).*

Proof. Let us assume that $n \geq 3$. (We will treat the cases $n = 1$ and $n = 2$ later.) By Pólya's enumeration theorem,

$$\beta(n) = \frac{1}{8n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (2^{A_{ij}} + 2^{B_{ij}} + 2^{C_{ij}} + 2^{D_{ij}} + 2^{E_{ij}} + 2^{F_{ij}} + 2^{G_{ij}} + 2^{H_{ij}}),$$

where A_{ij} (resp., $B_{ij}, C_{ij}, D_{ij}, E_{ij}, F_{ij}, G_{ij}, H_{ij}$) is the number of cycles in the permutation $\sigma^i \tau^j$ (resp., $\sigma^i \tau^j \rho, \sigma^i \tau^j \theta, \sigma^i \tau^j \rho \theta, \sigma^i \tau^j \zeta, \sigma^i \tau^j \rho \zeta, \sigma^i \tau^j \theta \zeta, \sigma^i \tau^j \rho \theta \zeta$); here σ rotates the rows (row 0 becomes row 1, row 1 becomes row 2, \dots , row $n-1$ becomes row 0), τ rotates the columns, ρ reflects the rows (rows 0 and $n-1$ are interchanged, rows 1 and $n-2$ are interchanged, \dots , rows $\lfloor n/2 \rfloor - 1$ and $n - \lfloor n/2 \rfloor$ are interchanged), θ reflects the columns, and ζ transposes the matrix. The order of the group generated by $\sigma, \tau, \rho, \theta$, and ζ is $8n^2$, using the assumption that $n \geq 3$.

We have already evaluated

$$a(n, n) = \frac{1}{n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{A_{ij}},$$

$$\alpha(n) = \frac{1}{2n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (2^{A_{ij}} + 2^{E_{ij}}),$$

and

$$b(n, n) = \frac{1}{4n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (2^{A_{ij}} + 2^{B_{ij}} + 2^{C_{ij}} + 2^{D_{ij}}),$$

so

$$\beta(n) = \frac{1}{2} b(n, n) + \frac{1}{4} \left(\alpha(n) - \frac{1}{2} a(n, n) \right) + \frac{1}{8n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (2^{F_{ij}} + 2^{G_{ij}} + 2^{H_{ij}}). \quad (11)$$

Let us begin with

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{H_{ij}}.$$

Here we are concerned with the permutations $\sigma^i \tau^j \rho \theta \zeta$ for $(i, j) \in \{0, 1, \dots, n-1\}^2$. We will need some multiplication rules for the permutations $\sigma, \tau, \rho, \theta$, and ζ , specifically

$$\sigma\tau = \tau\sigma, \quad \sigma\theta = \theta\sigma, \quad \tau\rho = \rho\tau, \quad \rho\theta = \theta\rho, \quad \sigma\rho = \rho\sigma^{-1}, \quad \tau\theta = \theta\tau^{-1},$$

and

$$\sigma\zeta = \zeta\tau, \quad \tau\zeta = \zeta\sigma, \quad \rho\zeta = \zeta\theta, \quad \theta\zeta = \zeta\rho.$$

It follows that (with $\tau^{-i} := (\tau^{-1})^i$)

$$\sigma^i \tau^j \rho \theta \zeta = \sigma^i \tau^j \zeta \theta \rho = \zeta \tau^i \sigma^j \theta \rho = \zeta \theta \tau^{-i} \sigma^j \rho = \zeta \theta \rho \tau^{-i} \sigma^{-j},$$

and hence

$$(\sigma^i \tau^j \rho \theta \zeta)^2 = (\sigma^i \tau^j \rho \theta \zeta)(\zeta \theta \rho \tau^{-i} \sigma^{-j}) = \sigma^{i-j} \tau^{-i+j} = (\sigma \tau^{-1})^{i-j} = (\sigma^{-1} \tau)^{-i+j}. \quad (12)$$

In particular, if $i \in \{0, 1, \dots, n-1\}$, then the permutation $\sigma^i \tau^i \rho \theta \zeta$ is of order 2. Furthermore, under this permutation, the entry in position (k, l) moves to position $(n-1-[l+i], n-1-[k+i])$, where, as before, $[r] := (r \bmod n) \in \{0, 1, \dots, n-1\}$. Thus, (k, l) is a fixed point if and only if

$$(k, l) = (n-1-[l+i], n-1-[k+i]). \quad (13)$$

For each $k \in \{0, 1, \dots, n-1\}$ there is a unique $l \in \{0, 1, \dots, n-1\}$ (namely $l := n-1-[k+i]$) such that (13) holds; indeed,

$$\begin{aligned} n-1-[l+i] &= n-1-[n-1-[k+i]+i] = n-1-[n-1-(k+i)+i] \\ &= n-1-[n-1-k] = n-1-(n-1-k) = k. \end{aligned}$$

Thus, $\sigma^i \tau^i \rho \theta \zeta$ with $i \in \{0, 1, \dots, n-1\}$ is of order 2 and has exactly n fixed points, hence $\binom{n}{2}$ transpositions. This implies that $H_{ii} = n(n+1)/2$ for such i .

Now we let $(i, j) \in \{0, 1, \dots, n-1\}^2$ be arbitrary but with $i \neq j$. Let us generalize (12) to

$$\begin{aligned} (\sigma^i \tau^j \rho \theta \zeta)^{2d} &= \sigma^{(i-j)d} \tau^{(-i+j)d} = ((\sigma \tau^{-1})^{i-j})^d = ((\sigma^{-1} \tau)^{-i+j})^d, \\ (\sigma^i \tau^j \rho \theta \zeta)^{2d+1} &= \sigma^{(i-j)d+i} \tau^{(-i+j)d+j} \rho \theta \zeta. \end{aligned}$$

The proof proceeds much like the proof of Theorem 1. Specifically, $\sigma^i \tau^j \rho \theta \zeta$ is of order $2d$, where $d := n/\gcd(|i-j|, n)$. All cycles have length d or $2d$. In fact, if d is odd, there are n/d cycles of length d and $n(n-1)/(2d)$ cycles of length $2d$. If d is even, there are $n^2/(2d)$ cycles, all of length $2d$. And for a given $(i, j) \in \{0, 1, \dots, n-1\}^2$, there are n pairs $(k, l) \in \{0, 1, \dots, n-1\}^2$ such that $[k-l] = [|i-j|]$. We arrive at the conclusion that

$$\frac{1}{8n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{H_{ij}} = \frac{1}{8n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{E_{ij}} = \frac{1}{4} \left(\alpha(n) - \frac{1}{2} a(n, n) \right). \quad (14)$$

Next we evaluate

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{F_{ij}} = \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{G_{ij}}, \quad (15)$$

where the equality holds by symmetry. We consider the permutations $\sigma^i \tau^j \rho \zeta$ for $(i, j) \in \{0, 1, \dots, n-1\}^2$. From the multiplication rules, it follows that

$$\sigma^i \tau^j \rho \zeta = \zeta \theta \tau^{-i} \sigma^j$$

and hence

$$(\sigma^i \tau^j \rho \zeta)^2 = (\sigma^i \tau^j \rho \zeta)(\zeta \theta \tau^{-i} \sigma^j) = \sigma^i \tau^j \rho \theta \tau^{-i} \sigma^j = \sigma^{i-j} \tau^{i+j} \rho \theta = \theta \rho \sigma^{-i+j} \tau^{-i-j}, \quad (16)$$

which implies

$$(\sigma^i \tau^j \rho \zeta)^4 = (\sigma^{i-j} \tau^{i+j} \rho \theta)(\theta \rho \sigma^{-i+j} \tau^{-i-j}) = e.$$

So the permutation $\sigma^i \tau^j \rho \zeta$ is of order 4. The entry in position (k, l) moves to position $([l + j], n - 1 - [k + i])$ under this permutation. Thus, $(k, l) \in \{0, 1, \dots, n - 1\}^2$ is a fixed point of $\sigma^i \tau^j \rho \zeta$ if and only if

$$(k, l) = ([l + j], n - 1 - [k + i]).$$

There is a solution (k, l) if and only if there exists $l \in \{0, 1, \dots, n - 1\}$ such that, with $k := [l + j]$, we have $n - 1 - [k + i] = l$ or, equivalently,

$$[l + i + j] = n - 1 - l. \tag{17}$$

When $i + j \leq n - 1$, (17) is equivalent to

$$l + i + j = n - 1 - l \quad \text{or} \quad l + i + j - n = n - 1 - l$$

or to

$$l = (n - 1 - i - j)/2 \quad \text{or} \quad l = (2n - 1 - i - j)/2.$$

If n is odd and $i + j$ is odd, then there is one fixed point, $(k, l) = ([(2n - 1 - i + j)/2], [(2n - 1 - i - j)/2])$. If n is odd and $i + j$ is even, then there is one fixed point, $(k, l) = ([(n - 1 - i + j)/2], [(n - 1 - i - j)/2])$. If n is even and $i + j$ is odd, then there are two fixed points, namely

$$\begin{aligned} (k, l) &= ([(n - 1 - i + j)/2], [(n - 1 - i - j)/2]), \\ (k, l) &= ([(2n - 1 - i + j)/2], [(2n - 1 - i - j)/2]). \end{aligned}$$

Finally, if n is even and $i + j$ is even, then there is no fixed point.

When $i + j \geq n$, (17) is equivalent to

$$l + i + j - n = n - 1 - l \quad \text{or} \quad l + i + j - 2n = n - 1 - l$$

or to

$$l = (2n - 1 - i - j)/2 \quad \text{or} \quad l = (3n - 1 - i - j)/2.$$

If n is odd and $i + j$ is odd, then there is one fixed point, $(k, l) = ([(2n - 1 - i + j)/2], [(2n - 1 - i - j)/2])$. If n is odd and $i + j$ is even, then there is one fixed point, $(k, l) = ([(3n - 1 - i + j)/2], [(3n - 1 - i - j)/2])$. If n is even and $i + j$ is odd, then there are two fixed points, namely

$$\begin{aligned} (k, l) &= ([(2n - 1 - i + j)/2], [(2n - 1 - i - j)/2]), \\ (k, l) &= ([(n - 1 - i + j)/2], [(n - 1 - i - j)/2]). \end{aligned}$$

Finally, if n is even and $i + j$ is even, then there is no fixed point. Notice that the results are the same for $i + j \geq n$ as for $i + j \leq n - 1$.

Using (16), under the permutation $(\sigma^i \tau^j \rho \zeta)^2$, the entry in position (k, l) moves to position $(n-1-[k+i-j], n-1-[l+i+j])$. Thus, $(k, l) \in \{0, 1, \dots, n-1\}^2$ is a fixed point of $(\sigma^i \tau^j \rho \zeta)^2$ if and only if

$$(k, l) = (n-1-[k+i-j], n-1-[l+i+j]).$$

A necessary and sufficient condition on (k, l) is (17) together with $[k+i-j] = n-1-k$. Solutions have l as before. On the other hand, k must satisfy

$$k+i-j-n = n-1-k, \quad k+i-j = n-1-k, \quad \text{or} \quad k+i-j+n = n-1-k,$$

or equivalently,

$$k = [(n-1-i+j)/2] \quad \text{or} \quad k = [(2n-1-i+j)/2].$$

If n is odd, the only fixed points of $(\sigma^i \tau^j \rho \zeta)^2$ are those already shown to be fixed points of $\sigma^i \tau^j \rho \zeta$. If n is even and $i+j$ is odd, there are two fixed points of $(\sigma^i \tau^j \rho \zeta)^2$ that are not fixed points of $\sigma^i \tau^j \rho \zeta$, namely

$$(k, l) = ([(n-1-i+j)/2], [(2n-1-i-j)/2]), \\ (k, l) = ([(2n-1-i+j)/2], [(n-1-i-j)/2]).$$

Finally, there are no fixed points when n is even and $i+j$ is even.

Consequently, if n is odd, then the permutation $\sigma^i \tau^j \rho \zeta$, which is of order 4, has only one fixed point. Therefore, it has one cycle of length 1 and $(n^2-1)/4$ cycles of length 4. Thus,

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{F_{ij}} = n^2 2^{(n^2+3)/4}.$$

For even n , if $i+j$ is odd, then the permutation $\sigma^i \tau^j \rho \zeta$ has two cycles of length 1 and one cycle of length 2, and the remaining cycles are of length 4. If $i+j$ is even, then all cycles of the permutation $\sigma^i \tau^j \rho \zeta$ are of length 4, hence there are $n^2/4$ of them. Thus,

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{F_{ij}} = \frac{1}{2} n^2 2^{(n^2-4)/4+3} + \frac{1}{2} n^2 2^{n^2/4} = 5n^2 2^{n^2/4-1}.$$

These results, together with (3), (10), (11), (14), and (15), yield (4).

Finally, recall that we have assumed that $n \geq 3$. We notice that the formula (4) gives $\beta(1) = 2$ and $\beta(2) = 6$, which are correct, as we can see by direct enumeration. \square

In the derivation of (2) in [1], the proof requires $m, n \geq 3$ because the group $D_m \times D_n$ used in the application of Pólya's enumeration theorem (D_m being the dihedral group of order $2m$), is incorrect if m or n is 1 or 2. If $m = 2$, row rotation and row reflection are the same, so the latter is redundant. Thus, D_2 should be replaced by C_2 , the cyclic group

of order 2. The reason (2) is still valid is that $b_1(2, n) = b_2(2, n)$ and $b_3(2, n) = b_4(2, n)$, as is easily verified. If $m = 1$, again row reflection is redundant, so D_1 should be replaced by C_1 . Here (2) remains valid because $b_1(1, n) = b_2(1, n)$ and $b_3(1, n) = b_4(1, n)$. A similar remark applies to $n = 2$ and $n = 1$, except that here $b_1(m, 2) = b_3(m, 2)$, $b_2(m, 2) = b_4(m, 2)$, $b_1(m, 1) = b_3(m, 1)$, and $b_2(m, 1) = b_4(m, 1)$.

References

- [1] S. N. Ethier, Counting toroidal binary arrays, *J. Integer Sequences* **16** (2013), [Article 13.4.7](#).
- [2] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, <http://oeis.org/>, 2015.
- [3] J. H. van Lint and R. M. Wilson, *A Course in Combinatorics*, 2nd ed., Cambridge University Press, 2001.

2010 *Mathematics Subject Classification*: Primary 05A05.

Keywords: toroidal array, Euler's phi function, group action, orbit, Pólya's enumeration theorem.

(Concerned with sequences [A179043](#), [A184271](#), [A209251](#), [A222188](#), [A255015](#), and [A255016](#).)

Received February 19 2015; revised version received June 3 2015. Published in *Journal of Integer Sequences*, July 16 2015.

Return to [Journal of Integer Sequences home page](#).