



Largest Values of the Stern Sequence, Alternating Binary Expansions and Continuants

Roland Paulin
Max-Planck-Institut für Mathematik
Vivatsgasse 7
D-53111, Bonn
Germany
paulinroland@gmail.com

Abstract

We study the largest values of the r th row of Stern's diatomic array. In particular, we prove some conjectures of Lansing. Our main tool is the connection between the Stern sequence, alternating binary expansions, and continuants. This allows us to reduce the problem of ordering the elements of the Stern sequence to the problem of ordering continuants. We describe an operation that increases the value of a continuant, allowing us to reduce the problem of largest continuants to ordering continuants of very special shape. Finally, we order these special continuants using some identities and inequalities involving Fibonacci numbers.

1 Introduction

The Stern sequence [A002487](#) is defined as follows: $s(0) = 0$, $s(1) = 1$, $s(2n) = s(n)$ and $s(2n+1) = s(n) + s(n+1)$ for every $n \geq 0$. Stern's diatomic array consists of rows indexed by $0, 1, 2, \dots$, where if $r \geq 0$, then the r th row is $s(2^r), s(2^r + 1), \dots, s(2^{r+1})$. Stern's diatomic array can also be constructed in the following way. Start with the 0th row $1, 1$. If $r \geq 1$, then to construct the r th row, copy the previous row, and between every two consecutive numbers x, y , write their sum $x + y$. This array was first studied by Stern [7]. Lehmer [4]

summarized several properties of this array. Recently Defant [1] gave upper bounds for the elements of the Stern sequence.

For $r \geq 0$ and $m \geq 1$, let $L_m(r)$ denote the m th largest distinct value of the r th row of Stern's diatomic array. If there are fewer than m distinct values in the r th row, then we just take $L_m(r) = -\infty$. Lucas [5] has observed that the largest value in the r th row is $L_1(r) = F_{r+2}$, where F_n denotes the n th Fibonacci number. Lansing [3] determined the second and third largest values $L_2(r)$ and $L_3(r)$, and formulated the following conjectures about the $L_m(r)$'s. Her Conjecture 7 [3] says that if $m \geq 1$ and $r \geq 4m - 2$, then

$$L_m(r) = L_m(r - 1) + L_m(r - 2).$$

Her Conjecture 9 [3] says that if $m \geq 2$ and $r \geq 4m - 4$, then

$$L_m(r) = L_{m-1}(r) - F_{r-(4m-5)} = F_{r+2} - \sum_{j=2}^m F_{r-(4j-5)}.$$

The following theorem is our main result. It gives a formula for $L_m(r)$ for certain values of r and m , and it implies the two conjectures of Lansing stated above.

Theorem 1. *If $r \in \mathbb{Z}_{\geq 0}$, then*

$$\{L_1(r), \dots, L_{\lceil \frac{r}{2} \rceil}(r)\} = \{F_{r+2} - F_i F_j; i, j \in \mathbb{Z}_{\geq 0}, i + j = r - 1\}.$$

More explicitly, if $1 \leq m \leq \lceil \frac{r}{2} \rceil$, then

$$L_m(r) = F_{r+2} - F_{2m-2-b} F_{r-2m+1+b},$$

where

$$b = b(m, r) = \begin{cases} 0, & \text{if } m \leq \lfloor \frac{r+3}{4} \rfloor \text{ or } 2 \mid r; \\ 1, & \text{if } m > \lfloor \frac{r+3}{4} \rfloor \text{ and } 2 \nmid r. \end{cases}$$

We show now that Theorem 1 implies the two conjectures stated above.

Proof of Conjecture 7 [3]. Let $m \geq 1$ and $r \geq 4m - 2$. Then $m \leq \lfloor \frac{(r-1)+3}{4} \rfloor$, so $L_m(r) = F_{r+2} - F_{2m-2} F_{r-2m+1}$ and $L_m(r-1) = F_{r+1} - F_{2m-2} F_{r-2m}$ by Theorem 1. If $r \geq 4m - 1$, then $m \leq \lfloor \frac{(r-2)+3}{4} \rfloor$, while if $r = 4m - 2$, then $2 \mid r - 2$, so either way $L_m(r-2) = F_r - F_{2m-2} F_{r-2m-1}$. The conjecture now follows from

$$F_{r+2} - F_{2m-2} F_{r-2m+1} = (F_{r+1} - F_{2m-2} F_{r-2m}) + (F_r - F_{2m-2} F_{r-2m-1}).$$

□

Proof of Conjecture 9 [3]. It is enough to prove $L_{m-1}(r) - L_m(r) = F_{r-(4m-5)}$, because $L_1(r) = F_{r+2}$. Now $m - 1 \leq \lfloor \frac{r+3}{4} \rfloor$, and either $2 \mid r = 4m - 4$, or $m \leq \lfloor \frac{r+3}{4} \rfloor$. So $L_{m-1}(r) = F_{r+2} - F_{2m-4} F_{r-2m+3}$ and $L_m(r) = F_{r+2} - F_{2m-2} F_{r-2m+1}$ by Theorem 1. Hence

$$L_{m-1}(r) - L_m(r) = F_{2m-2} F_{r-2m+1} - F_{2m-4} F_{r-2m+3} = (-1)^{2m-4} F_2 F_{r-(4m-5)} = F_{r-(4m-5)}$$

by Lemma 3. □

Here is a brief description of the contents of the paper. In Section 2 we state some basic results about Fibonacci numbers, alternating binary expansions and continuants. In Section 3 we describe the connection between the Stern sequence, alternating binary expansions and continuants. Using this description we reduce the problem of comparing the elements of the Stern sequence to the problem of comparing continuants, which is the subject of Section 4. (For earlier work on comparing continuants, see Ramharter's paper [6].) Using some identities involving Fibonacci numbers, we finish the proof of Theorem 1 in Section 5. Finally, in Section 6 we discuss possible extensions of our results.

2 Preliminaries

In this section we introduce some notation, and state some basic results about Fibonacci numbers, alternating binary expansions and continuants. Let F_n denote the n th Fibonacci number for every $n \in \mathbb{Z}$. So $F_0 = 0$, $F_1 = 1$, and $F_n = F_{n-1} + F_{n-2}$ for every $n \in \mathbb{Z}$. Note that $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^n = \begin{pmatrix} F_{n+1} & F_n \\ F_n & F_{n-1} \end{pmatrix}$ for every $n \in \mathbb{Z}$. The following basic lemma describes the sign of the Fibonacci numbers.

Lemma 2. *If $n \in \mathbb{Z}$, then $F_{-n} = (-1)^{n+1}F_n$. Hence*

$$F_n \begin{cases} = 0, & \text{if } n = 0; \\ > 0, & \text{if } n > 0 \text{ or } 2 \nmid n; \\ < 0, & \text{if } n < 0 \text{ and } 2 \mid n. \end{cases}$$

Proof. The first part is easy to check by induction on n , and the second part follows from the first part. \square

The following lemma describes a useful identity, which allows us to compare $F_i F_j$'s with fixed $i + j$.

Lemma 3. *If $i, j, k, l \in \mathbb{Z}$ and $i + j = k + l$, then*

$$F_i F_j - F_k F_l = (-1)^k F_{i-k} F_{j-k}.$$

Hence if $i + j = k + l$, then $F_i F_j = F_k F_l$ if and only if $\{i, j\} = \{k, l\}$.

Proof. If $n, i, j \in \mathbb{Z}$, then

$$F_{n+i} F_{n+j} - F_n F_{n+i+j} = (-1)^n F_i F_j.$$

Dickson [2, p. 404] attributes this identity to A. Tagiuri. Substituting k , $i - k$ and $j - k$ into n , i and j , we obtain the stated identity. To prove the second part, note that by Lemma 2, if $n \in \mathbb{Z}$, then $F_n = 0$ if and only if $n = 0$. \square

The following lemma describes the ordering of $F_i F_j$'s with $i, j \geq 0$ and $i + j$ fixed. We need this result to prove the second half of Theorem 1.

Lemma 4. *Let $n \in \mathbb{Z}_{\geq 0}$, then the set*

$$\{F_i F_j; i, j \in \mathbb{Z}_{\geq 0}, i + j = n\}$$

has cardinality $\lceil \frac{n+1}{2} \rceil$, and if $m \in \{1, \dots, \lceil \frac{n+1}{2} \rceil\}$, then the m th smallest element of this set is $F_{2m-2-c} F_{n-(2m-2-c)}$, where

$$c = \begin{cases} 0, & \text{if } m \leq \lfloor \frac{n+4}{4} \rfloor \text{ or } 2 \nmid n; \\ 1, & \text{if } m > \lfloor \frac{n+4}{4} \rfloor \text{ and } 2 \mid n. \end{cases}$$

Proof. The last part of Lemma 3 implies that the cardinality is $\lceil \frac{n+1}{2} \rceil$. For $m \in \{1, \dots, \lceil \frac{n+1}{2} \rceil\}$ let $c(m)$ be defined as c in the statement above, and let $u(m) = 2m - 2 - c(m)$ and $v(m) = n - (2m - 2 - c(m))$. Then $u(m), v(m) \geq 0$ and $u(m) + v(m) = n$, so $F_{u(m)} F_{v(m)}$ is an element of the set. Let $1 \leq m < m' \leq \lceil \frac{n+1}{2} \rceil$. All we need to prove is that $F_{u(m)} F_{v(m)} < F_{u(m')} F_{v(m')}$. The difference is

$$F_{u(m')} F_{v(m')} - F_{u(m)} F_{v(m)} = (-1)^{u(m)} F_{u(m')-u(m)} F_{v(m')-u(m)}$$

by Lemma 3. Here

$$u(m') - u(m) = 2(m' - m) - c(m') + c(m) \geq 2 - 1 > 0,$$

so $F_{u(m')-u(m)} > 0$. Therefore we need to prove that $(-1)^{c(m)} F_{v(m')-u(m)} > 0$. First suppose that $2 \nmid n$. Then $c(m) = c(m') = 0$, so $v(m') - u(m) = n - 2(m' + m - 2)$ is odd, hence $(-1)^{c(m)} F_{v(m')-u(m)} > 0$ by Lemma 2. So let $2 \mid n$.

Suppose that $m > \lfloor \frac{n+4}{4} \rfloor$. Then $c(m) = c(m') = 1$, and $m > \lfloor \frac{n+4}{4} \rfloor = \lceil \frac{n+2}{4} \rceil$, so $m - 1 \geq \lceil \frac{n+2}{4} \rceil \geq \frac{n+2}{4} > \frac{n}{4}$, hence

$$2 \mid v(m') - u(m) = n - 2(m' + m - 3) \leq n - 2(2m + 1 - 3) = n - 4(m - 1) < 0.$$

Thus $(-1)^{c(m)} F_{v(m')-u(m)} > 0$ by Lemma 2.

Finally, let $m \leq \lfloor \frac{n+4}{4} \rfloor$. Then $c(m) = 0$. If $m' > \lfloor \frac{n+4}{4} \rfloor$, then $c(m') = 1$ and $2 \nmid v(m') - u(m)$, so $F_{v(m')-u(m)} > 0$. If $m' \leq \lfloor \frac{n+4}{4} \rfloor$, then $c(m') = 0$ and $4m' - 4 \leq n$, so

$$v(m') - u(m) = n - 2(m' + m - 2) \geq n - 2(2m' - 3) > n - (4m' - 4) \geq 0,$$

hence $F_{v(m')-u(m)} > 0$. □

Now we turn to the discussion of alternating binary expansions. For $d, l_0 \in \mathbb{Z}_{\geq 0}$, $l_1, \dots, l_d \in \mathbb{Z}_{\geq 1}$ we define

$$A(l_0, \dots, l_d) = \sum_{i=0}^d (-1)^{d-i} 2^{l_0 + \dots + l_i}.$$

We call this an alternating binary expansion. The following lemma gives a bound for $A(l_0, \dots, l_d)$.

Lemma 5. *If $d, l_0 \in \mathbb{Z}_{\geq 0}$ and $l_1, \dots, l_d \in \mathbb{Z}_{\geq 1}$, then*

$$2^{l_0 + \dots + l_d - 1} \leq A(l_0, \dots, l_d) \leq 2^{l_0 + \dots + l_d}.$$

If $d > 0$, then $A(l_0, \dots, l_d) < 2^{l_0 + \dots + l_d}$.

Proof. We prove by induction on d . For $d = 0$ this is trivial, so suppose that $d \geq 1$ and that the statement is true for smaller values of d . Let $k_i = l_0 + \dots + l_i$ for every $i \in \{0, 1, \dots, d\}$. Then $0 \leq k_0 < k_1 < \dots < k_d$, and $A(l_0, \dots, l_d) = 2^{k_d} - 2^{k_{d-1}} + 2^{k_{d-2}} - \dots + (-1)^d 2^{k_0} = 2^{k_d} - A(l_0, \dots, l_{d-1})$. Here $0 < 2^{k_{d-1}-1} \leq A(l_0, \dots, l_{d-1}) \leq 2^{k_{d-1}} \leq 2^{k_d-1}$ by the induction hypothesis, so $2^{k_d-1} \leq A(l_0, \dots, l_d) < 2^{k_d}$. \square

The following lemma describes the number of alternating binary expansions of a positive integer.

Lemma 6. *Every $n \in \mathbb{Z}_{\geq 1}$ has exactly two alternating binary expansions, and exactly one of these has the form $n = A(l_0, 1, l_2, \dots, l_d)$, where $l_0 \in \mathbb{Z}_{\geq 0}$ and $d, l_2, \dots, l_d \in \mathbb{Z}_{\geq 1}$. If n is a power of 2, then $d = 1$ and the other expansion is $n = A(l_0)$, while otherwise $d \geq 2$ and the other expansion is $A(l_0, l_2 + 1, l_3, \dots, l_d)$.*

Proof. It is easy to check that $A(l_0, 1) = A(l_0) = 2^{l_0}$ and $A(l_0, 1, l_2, \dots, l_d) = A(l_0, l_2 + 1, l_3, \dots, l_d)$ for $d \geq 2$. Hence it is enough to prove that every $n \in \mathbb{Z}_{\geq 1}$ has exactly one alternating binary expansion of the form $n = A(l_0, \dots, l_d)$ such that $d \geq 1$ and $l_1 = 1$. We prove by induction on n . Let k be the unique positive integer such that $2^{k-1} \leq n < 2^k$. Let $k_i = l_0 + \dots + l_i$ for $i \in \{0, \dots, d\}$. Then $2^{k_d} \leq n < 2^{k_d}$ by Lemma 5, so $k_d = k$. Moreover $n = 2^k - A(l_0, \dots, l_{d-1})$, so $n' = A(l_0, \dots, l_{d-1}) = 2^k - n$. If $d = 1$, then $l_0 = k - 1$ and $n = A(l_0, 1) = 2^{k-1}$. Now let $d \geq 2$. Then $A(l_0, \dots, l_{d-1}) < 2^{k_{d-1}} \leq 2^{k-1}$ by Lemma 5, so $2^{k-1} < n < 2^k$ and $n' < 2^{k-1} < n$. By the induction hypothesis, there is a unique expansion $n' = A(l_0, \dots, l_{d-1})$ with $l_1 = 1$, where furthermore $l_0 + \dots + l_{d-1} \leq k - 1$, because $n' < 2^{k-1}$. Then $l_d = k - (l_0 + \dots + l_{d-1}) \in \mathbb{Z}_{\geq 1}$ is also determined. \square

Now we will recall some basic facts about continuants. Continuants are polynomials in several variables, which come up often when working with continued fractions. We define the d th continuant $K_d(X_1, \dots, X_d) \in \mathbb{Z}[X_1, \dots, X_d]$ for every $d \in \mathbb{Z}_{\geq 0}$, by the following recursion: $K_0 = 1$, $K_1(X_1) = X_1$, and

$$K_d(X_1, \dots, X_d) = X_d K_{d-1}(X_1, \dots, X_{d-1}) + K_{d-2}(X_1, \dots, X_{d-2})$$

for every $d \geq 2$. We can safely write $K(X_1, \dots, X_d)$ instead of $K_d(X_1, \dots, X_d)$, because d is anyway determined by the number of variables. If $\underline{X} = (X_1, \dots, X_d)$, then we will also use the notation $K(\underline{X}) = K(X_1, \dots, X_d)$. The continuants are related to continued fractions by the following identity:

$$[a_0, a_1, \dots, a_d] = a_0 + \frac{1}{a_1 + \frac{1}{\ddots + \frac{1}{a_d}}}} = \frac{K(a_0, \dots, a_d)}{K(a_1, \dots, a_d)}$$

for every $d \geq 0$.

The following lemma gives maybe the most practical description of continuants, using 2×2 matrices.

Lemma 7. *If $d \geq 2$, then*

$$\begin{pmatrix} K_d(X_1, \dots, X_d) & K_{d-1}(X_1, \dots, X_{d-1}) \\ K_{d-1}(X_2, \dots, X_d) & K_{d-2}(X_2, \dots, X_{d-1}) \end{pmatrix} = \begin{pmatrix} X_1 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} X_d & 1 \\ 1 & 0 \end{pmatrix}.$$

So if $d \geq 0$, then $K_d(X_1, \dots, X_d) = M_{1,1}$, where $M = \begin{pmatrix} X_1 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} X_d & 1 \\ 1 & 0 \end{pmatrix}$.

Proof. This easily follows from the defining recursion by induction on d . □

The continuants have the following symmetry property.

Lemma 8. $K(X_1, \dots, X_d) = K(X_d, \dots, X_1)$ for every $d \geq 0$.

Proof. Let $M = \begin{pmatrix} X_1 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} X_d & 1 \\ 1 & 0 \end{pmatrix}$, then

$$M^T = \begin{pmatrix} X_d & 1 \\ 1 & 0 \end{pmatrix}^T \cdots \begin{pmatrix} X_1 & 1 \\ 1 & 0 \end{pmatrix}^T = \begin{pmatrix} X_d & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} X_1 & 1 \\ 1 & 0 \end{pmatrix},$$

hence

$$K(X_d, \dots, X_1) = (M^T)_{1,1} = M_{1,1} = K(X_1, \dots, X_d)$$

by Lemma 7. □

The following lemma states a few simple identities involving continuants.

Lemma 9. *If $d \geq 1$, then*

$$K(X_1, \dots, X_d) = K(X_1 - 1, X_2, \dots, X_d) + K(X_2, \dots, X_d).$$

If $d \geq 2$, then

$$K(1, X_2, \dots, X_d) = K(X_2, \dots, X_d) + K(X_3, \dots, X_d).$$

If $d \geq 1$, then

$$K(1, X_1, X_2, \dots, X_d) = K(X_1 + 1, X_2, \dots, X_d)$$

and

$$K(X_1, \dots, X_{d-1}, X_d, 1) = K(X_1, \dots, X_{d-1}, X_d + 1).$$

Proof. The identities are trivial for $d = 1$, so assume that $d \geq 2$. Using Lemma 8 and the defining recursion of the continuants, we obtain

$$K(X_1, \dots, X_d) = X_1 K(X_2, \dots, X_d) + K(X_3, \dots, X_d).$$

Substituting $X_1 - 1$ into X_1 , we get

$$K(X_1 - 1, X_2, \dots, X_d) = (X_1 - 1)K(X_2, \dots, X_d) + K(X_3, \dots, X_d).$$

These two equations immediately imply the first part of the lemma. Substituting 1 into X_1 in the first equation, we obtain the second part of the lemma.

The fourth identity follows from the third one by Lemma 8. Finally,

$$K(X_1 + 1, X_2, \dots, X_d) = K(X_1, \dots, X_d) + K(X_2, \dots, X_d) = K(1, X_1, \dots, X_d)$$

by the first two parts of the lemma. □

3 Connecting the Stern sequence, alternating binary expansions and continuants

The following proposition describes the connection between the Stern sequence, alternating binary expansions and continuants. This result plays a central role in this paper: it allows us to reduce the problem of ordering the elements of the Stern sequence to the problem of ordering continuants.

Proposition 10. *If $d, l_0 \in \mathbb{Z}_{\geq 0}$ and $l_1, \dots, l_d \in \mathbb{Z}_{\geq 1}$, then*

$$s(A(l_0, l_1, \dots, l_d)) = K(l_1, \dots, l_d).$$

Proof. We prove by induction on $l_0 + \dots + l_d$. So let $S \in \mathbb{Z}_{\geq 0}$, and suppose the statement is true if $l_0 + \dots + l_d < S$. Now let $l_0 + \dots + l_d = S$. If $l_0 > 0$, then $A(l_0, l_1, \dots, l_d) = 2^{l_0} A(0, l_1, \dots, l_d)$, so $s(A(l_0, l_1, \dots, l_d)) = s(A(0, l_1, \dots, l_d)) = K(l_1, \dots, l_d)$ by the induction hypothesis. So assume that $l_0 = 0$.

Let $k_i = l_0 + \dots + l_i$ for every $i \in \{0, \dots, d\}$, and let $n = A(l_0, l_1, \dots, l_d) = \sum_{i=0}^d (-1)^{d-i} 2^{k_i}$. Note that $k_0 = l_0 = 0$, so $k_1 = l_1$. Here $n > 0$ by Lemma 5, and n is odd, so $n = 2m + 1$ for some $m \in \mathbb{Z}_{\geq 0}$. If $d = 0$, then $n = 1$, and $s(1) = 1 = K()$. So let $d \geq 1$.

First suppose that $l_1 \geq 2$. If d is even, then $m = A(l_1 - 1, l_2, \dots, l_d)$ and $m + 1 = A(0, l_1 - 1, l_2, \dots, l_d)$, while if d is odd, then $m + 1 = A(l_1 - 1, l_2, \dots, l_d)$ and $m = A(0, l_1 - 1, l_2, \dots, l_d)$. (Note that $l_1 - 1 \geq 1$.) So either way we get

$$\begin{aligned} s(n) &= s(m) + s(m + 1) = s(A(0, l_1 - 1, l_2, \dots, l_d)) + s(A(l_1 - 1, l_2, \dots, l_d)) \\ &= K(l_1 - 1, l_2, \dots, l_d) + K(l_2, \dots, l_d) = K(l_1, \dots, l_d) \end{aligned}$$

using the induction hypothesis and Lemma 9.

Now suppose that $l_1 = 1$. If $d = 1$, then $n = A(0, 1) = 1$ and $s(n) = 1 = K(1)$. So let $d \geq 2$. If d is odd, then $m = A(l_2, \dots, l_d)$ and $m + 1 = A(0, l_2, \dots, l_d)$, while if d is even, then $m + 1 = A(l_2, \dots, l_d)$ and $m = A(0, l_2, \dots, l_d)$. So either way we get

$$\begin{aligned} s(n) &= s(m) + s(m + 1) = s(A(0, l_2, \dots, l_d)) + s(A(l_2, \dots, l_d)) \\ &= K(l_2, \dots, l_d) + K(l_3, \dots, l_d) = K(1, l_2, \dots, l_d) = K(l_1, \dots, l_d) \end{aligned}$$

using the induction hypothesis and Lemma 9. □

For $r \in \mathbb{Z}$ let us define

$$E_r = \{(l_1, \dots, l_d); d \in \mathbb{Z}_{\geq 1}, l_1, \dots, l_d \in \mathbb{Z}_{\geq 1}, l_1 = l_d = 1, l_1 + \dots + l_d = r + 1\}$$

and

$$E'_r = \{(l_1, \dots, l_d); d \in \mathbb{Z}_{\geq 1}, l_1, \dots, l_d \in \mathbb{Z}_{\geq 1}, l_1 = l_d = 1, l_1 + \dots + l_d \leq r + 1\}.$$

Using Lemmas 5, 6 and Proposition 10, we obtain the following corollary.

Corollary 11. *If $r \in \mathbb{Z}_{\geq 0}$, then the set of values of the r th row of Stern's diatomic array is*

$$\{s(n); 2^r \leq n \leq 2^{r+1}\} = \{K(\underline{l}); \underline{l} \in E'_r\}.$$

Proof. Suppose that $\underline{l} \in E'_r$. Let $l_0 = r + 1 - \sum_{i=1}^r l_i$ and $n = A(l_0, \dots, l_d)$. Then $2^r \leq n \leq 2^{r+1}$ and $s(n) = K(\underline{l})$ by Lemma 5 and Proposition 10. So the left hand side contains the right hand side. Conversely, let $n \in \{2^r, \dots, 2^{r+1}\}$. If $n = 2^r$ or $n = 2^{r+1}$, then $s(n) = 1 = K(1)$. So suppose that $2^r < n < 2^{r+1}$. According to Lemma 6, n has an alternating binary expansion $n = A(l_0, \dots, l_d)$ with $d \geq 1$ and $l_1 = 1$. Here $l_1 + \dots + l_d \leq l_0 + \dots + l_d = r + 1$ by Lemma 5, and $s(n) = K(l_1, \dots, l_d)$ by Proposition 10. So if $l_d = 1$, then $\underline{l} = (l_1, \dots, l_d) \in E'_r$ and $s(n) = K(\underline{l})$. If $l_d > 1$, then $\underline{l}' = (l_1, \dots, l_d - 1, 1) \in E'_r$ and $s(n) = K(\underline{l}) = K(\underline{l}')$ by Lemma 9. So the right hand side contains the left hand side. \square

This corollary implies that $L_m(r)$ is the m th largest distinct value in $\{K(\underline{l}); \underline{l} \in E'_r\}$. So we need to compare the continuants $K(\underline{l})$, where $\underline{l} \in E'_r$.

4 Comparing continuants

The following proposition describes a simple operation on (l_1, \dots, l_d) that increases $K(l_1, \dots, l_d)$. This operation is our main tool in comparing continuants.

Proposition 12. *Let $d, l_1, \dots, l_d \in \mathbb{Z}_{\geq 1}$, $j \in \{1, \dots, d\}$, and suppose that $l_j = u + v$ for some $u, v \in \mathbb{Z}_{\geq 1}$. Then*

$$K(l_1, \dots, l_d) \leq K(l_1, \dots, l_{j-1}, u, v, l_{j+1}, \dots, l_d),$$

where equality holds if and only if $j = 1$ and $u = 1$, or $j = d$ and $v = 1$.

Proof. Lemma 7 implies that

$$K(l_1, \dots, l_d) = (P \begin{pmatrix} u+v & 1 \\ 1 & 0 \end{pmatrix} Q)_{1,1}$$

and

$$K(l_1, \dots, l_{j-1}, u, v, l_{j+1}, \dots, l_d) = (P \begin{pmatrix} u & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} v & 1 \\ 1 & 0 \end{pmatrix} Q)_{1,1},$$

where $P = \begin{pmatrix} l_1 & 1 \\ 1 & 0 \end{pmatrix} \dots \begin{pmatrix} l_{j-1} & 1 \\ 1 & 0 \end{pmatrix}$ and $Q = \begin{pmatrix} l_{j+1} & 1 \\ 1 & 0 \end{pmatrix} \dots \begin{pmatrix} l_d & 1 \\ 1 & 0 \end{pmatrix}$. Using

$$\begin{pmatrix} u & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} v & 1 \\ 1 & 0 \end{pmatrix} - \begin{pmatrix} u+v & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} (u-1)(v-1) & u-1 \\ v-1 & 1 \end{pmatrix} = \begin{pmatrix} u-1 \\ 1 \end{pmatrix} (v-1, 1)$$

we obtain

$$\begin{aligned} \Delta &= K(l_1, \dots, l_{j-1}, u, v, l_{j+1}, \dots, l_d) - K(l_1, \dots, l_d) = (P \begin{pmatrix} u-1 \\ 1 \end{pmatrix} (v-1, 1) Q)_{1,1} \\ &= (P_{1,1}(u-1) + P_{1,2})(Q_{1,1}(v-1) + Q_{2,1}). \end{aligned}$$

Here $P_{1,1}, P_{1,2}, Q_{1,1}, Q_{2,1}, u-1, v-1 \geq 0$, so $\Delta \geq 0$. Note that $\det P, \det Q \in \{-1, 1\}$, since $\det \begin{pmatrix} X & 1 \\ 1 & 0 \end{pmatrix} = -1$. So $(P_{1,1}, P_{1,2}) \neq (0, 0)$ and $(Q_{1,1}, Q_{2,1}) \neq (0, 0)$. Hence $\Delta = 0$ if and only if $u = 1$ and $P_{1,2} = 0$, or $v = 1$ and $Q_{2,1} = 0$. It is easy to see that $P_{1,2} = 0$ if and only if $j = 1$, and similarly, $Q_{2,1} = 0$ if and only if $j = d$. \square

We introduce some notation. Let

$$h(l_1, \dots, l_d) = \sum_{i=1}^d (l_i - 1) = l_1 + \dots + l_d - d.$$

For $r \in \mathbb{Z}$ and $a \in \mathbb{Z}_{\geq 0}$ let

$$E_{r,a} = \{(l_1, \dots, l_d) \in E_r; h(l_1, \dots, l_d) = a\}$$

and

$$E'_{r,a} = \{(l_1, \dots, l_d) \in E'_r; h(l_1, \dots, l_d) = a\}.$$

Then $E'_r = \bigcup_{a \geq 0} E'_{r,a} = \bigcup_{t \leq r} \bigcup_{a \geq 0} E_{t,a}$.

For $s, p_0, p_1, \dots, p_s \in \mathbb{Z}_{\geq 0}$ let

$$w_{p_0, \dots, p_s}(X_1, \dots, X_s) = (\underbrace{1, \dots, 1}_{p_0}, X_1, \underbrace{1, \dots, 1}_{p_1}, X_2, \dots, \underbrace{1, \dots, 1}_{p_{s-1}}, X_s, \underbrace{1, \dots, 1}_{p_s})$$

and

$$\kappa_{p_0, \dots, p_s}(X_1, \dots, X_s) = K(w_{p_0, \dots, p_s}(X_1, \dots, X_s)).$$

For $s = 0$ we simply write

$$w_{p_0} = (\underbrace{1, \dots, 1}_{p_0})$$

and $\kappa_{p_0} = K(w_{p_0})$.

The idea is that starting from $(l_1, \dots, l_d) \in E'_r$, and using the operation of Proposition 12 several times, and also possibly increasing elements or adding new elements to (l_1, \dots, l_d) , we can increase $K(l_1, \dots, l_n)$ to $K(w_{r+1})$. If we stop a bit earlier, we get the largest continuants. The precise statement is described in the following proposition.

Proposition 13. *If $r \in \mathbb{Z}_{\geq 0}$ and $\underline{l} \in E'_r \setminus (E_{r,0} \cup E_{r,1})$, then there is an $\underline{m} \in E_{r-1,0} \cup E_{r,2}$ such that $K(\underline{l}) \leq K(\underline{m})$.*

Proof. We will use several times that if $u_1, \dots, u_d \in \mathbb{Z}_{\geq 1}$, then

$$K(u_1, \dots, u_d) \leq K(u_1, \dots, u_d, 1),$$

and if $u'_1, \dots, u'_d \in \mathbb{Z}_{\geq 1}$ and $u_i \leq u'_i$ for every i , then

$$K(u_1, \dots, u_d) \leq K(u'_1, \dots, u'_d).$$

We prove by induction on $h(\underline{l})$. If $h(\underline{l}) = 0$, then $\underline{l} = w_d$ for some $d \leq r$, so we can take $\underline{m} = w_r \in E_{r-1,0}$. If $h(\underline{l}) = 1$, then $\underline{l} = w_{p_0, p_1}(2)$ for some $p_0, p_1 \in \mathbb{Z}_{\geq 1}$ with $p_0 + p_1 \leq r - 2$, so we can take $\underline{m} = w_{p_0, r-2-p_0}(3) \in E_{r,2}$. Finally, let $h(\underline{l}) \geq 2$, and suppose that the statement is true for smaller values of h . Then there is a $j \in \{2, \dots, d-1\}$ such that $l_j \geq 2$. Let

$$\underline{l}' = (l_1, \dots, l_{j-1}, l_j - 1, 1, l_{j+1}, \dots, l_d),$$

then $\underline{l}' \in E'_r$, and $K(\underline{l}) \leq K(\underline{l}')$ by Proposition 12. Moreover $h(\underline{l}') = h(\underline{l}) - 1 \geq 1$, so $\underline{l}' \notin E_{r,0}$. If $\underline{l}' \notin E_{r,1}$, then by the induction hypothesis there is an $\underline{m} \in E_{r-1,0} \cup E_{r,2}$ such that $K(\underline{l}) \leq K(\underline{l}') \leq K(\underline{m})$. So suppose that $\underline{l}' \in E_{r,1}$. Then $\underline{l} \in E_{r,2}$, so we can take $\underline{m} = \underline{l}$. \square

5 Fibonacci identities

Based on Proposition 13, our next goal is to calculate $K(\underline{l})$ for $\underline{l} \in E_{r,0} \cup E_{r,1} \cup E_{r-1,0} \cup E_{r,2}$. If $r \geq 1$, then $E_{r,0} = \{w_{r+1}\}$, $E_{r-1,0} = \{w_r\}$,

$$E_{r,1} = \{w_{p_0,p_1}(2); p_0, p_1 \in \mathbb{Z}_{\geq 1}, p_0 + p_1 = r - 1\},$$

and $E_{r,2} = U_r \cup V_r$, where

$$U_r = \{w_{p_0,p_1}(3); p_0, p_1 \in \mathbb{Z}_{\geq 1}, p_0 + p_1 = r - 2\}$$

and

$$V_r = \{w_{p_0,p_1,p_2}(2); p_0, p_2 \in \mathbb{Z}_{\geq 1}, p_1 \in \mathbb{Z}_{\geq 0}, p_0 + p_1 + p_2 = r - 3\}.$$

In general $\kappa_{p_0, \dots, p_s}(X_1, \dots, X_s) = M_{1,1}$, where

$$M = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^{p_0} \begin{pmatrix} X_1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^{p_1} \begin{pmatrix} X_2 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} X_s & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^{p_s}.$$

Note that $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^p = \begin{pmatrix} F_{p+1} & F_p \\ F_p & F_{p-1} \end{pmatrix}$ for every $p \in \mathbb{Z}$. Calculating the matrix products, we get

$$\kappa_{p_0} = F_{p_0+1},$$

$$\kappa_{p_0,p_1}(2) = 2F_{p_0+1}F_{p_1+1} + F_{p_0+1}F_{p_1} + F_{p_0}F_{p_1+1},$$

$$\kappa_{p_0,p_1}(3) = 3F_{p_0+1}F_{p_1+1} + F_{p_0+1}F_{p_1} + F_{p_0}F_{p_1+1},$$

$$\begin{aligned} \kappa_{p_0,p_1,p_2}(2,2) &= F_{p_0+1}F_{p_1}F_{p_2} + F_{p_0}F_{p_1+1}F_{p_2} + 2F_{p_0+1}F_{p_1+1}F_{p_2} + F_{p_0+1}F_{p_1-1}F_{p_2+1} \\ &\quad + F_{p_0}F_{p_1}F_{p_2+1} + 4F_{p_0+1}F_{p_1}F_{p_2+1} + 2F_{p_0}F_{p_1+1}F_{p_2+1} + 4F_{p_0+1}F_{p_1+1}F_{p_2+1}. \end{aligned}$$

In the following lemma we express these values in more useful forms.

Lemma 14. *If $p_0, p_1, p_2 \in \mathbb{Z}_{\geq 0}$, then*

$$\kappa_{p_0,p_1}(2) = F_{p_0+p_1+3} - F_{p_0}F_{p_1} = F_{p_0+p_1+2} + F_{p_0+p_1} + F_{p_0-1}F_{p_1-1},$$

$$\kappa_{p_0,p_1}(3) = F_{p_0+p_1+3} + F_{p_0+p_1+1} - 2F_{p_0+p_1-2} - 2F_{p_0-2}F_{p_1-2},$$

$$\begin{aligned} \kappa_{p_0,p_1,p_2}(2,2) &= (F_{p_0+p_1+p_2+4} + F_{p_0+p_1+p_2+2} - F_{p_0+p_1+p_2-4}) \\ &\quad - (F_{p_1}(F_{p_0-1}F_{p_2-1} + 3F_{p_0-2}F_{p_2-1} + 3F_{p_0-1}F_{p_2-2}) + 2F_{p_0-2}F_{p_1+1}F_{p_2-2}). \end{aligned}$$

Proof. If $i, j \in \mathbb{Z}$, then

$$\begin{pmatrix} F_{i+j+1} & F_{i+j} \\ F_{i+j} & F_{i+j-1} \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^{i+j} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^i \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^j = \begin{pmatrix} F_{i+1} & F_i \\ F_i & F_{i-1} \end{pmatrix} \begin{pmatrix} F_{j+1} & F_j \\ F_j & F_{j-1} \end{pmatrix},$$

hence $F_{i+j} = F_{i+1}F_j + F_iF_{j-1}$. Applying this identity twice, we get that $F_{i+j+k} = F_{i+j+1}F_k + F_{i+j}F_{k-1} = (F_{i+1}F_{j+1} + F_iF_j)F_k + (F_{i+1}F_j + F_iF_{j-1})F_{k-1}$ for $i, j, k \in \mathbb{Z}$. Using these identities, one can express every term in the statement of the Lemma as a polynomial of $F_{p_0}, F_{p_0+1}, F_{p_1}, F_{p_1+1}, F_{p_2}, F_{p_2+1}$. Comparing the obtained polynomials, one can check the stated identities. The calculations could be done by hand, but they are tedious. Instead we have used Mathematica [8] to carry out these symbolic calculations, this is done in the attached file. \square

Corollary 15. *If $r \geq 6$, then*

$$\begin{aligned} \max\{K(\underline{l}); \underline{l} \in E_{r,1}\} &< F_{r+2} = \kappa_{r+1}, \\ \min\{K(\underline{l}); \underline{l} \in E_{r,1}\} &= \kappa_{1,r-2}(2) = F_{r+1} + F_{r-1}, \\ \max\{K(\underline{l}); \underline{l} \in U_r\} &= \kappa_{2,r-4}(3) = F_{r+1} + F_{r-1} - 2F_{r-4}, \\ \max\{K(\underline{l}); \underline{l} \in V_r\} &= \kappa_{2,0,r-5}(2, 2) = F_{r+1} + F_{r-1} - F_{r-7}. \end{aligned}$$

So if $r \geq 6$, then

$$\max\{K(\underline{l}); \underline{l} \in E_{r-1,0} \cup E_{r,2}\} \leq \min\{K(\underline{l}); \underline{l} \in E_{r,1}\},$$

with strict inequality for $r \neq 7$.

Proof. Let $r \geq 6$. Recalling the description of $E_{r,1}$, U_r and V_r , we see that

$$\begin{aligned} \{K(\underline{l}); \underline{l} \in E_{r,1}\} &= \{\kappa_{p_0,p_1}(2); p_0, p_1 \in \mathbb{Z}_{\geq 1}, p_0 + p_1 = r - 1\}, \\ \{K(\underline{l}); \underline{l} \in U_r\} &= \{\kappa_{p_0,p_1}(3); p_0, p_1 \in \mathbb{Z}_{\geq 1}, p_0 + p_1 = r - 2\}, \\ \{K(\underline{l}); \underline{l} \in V_r\} &= \{\kappa_{p_0,p_1,p_2}(2); p_0, p_2 \in \mathbb{Z}_{\geq 1}, p_1 \in \mathbb{Z}_{\geq 0}, p_0 + p_1 + p_2 = r - 3\}. \end{aligned}$$

Using Lemma 14 one can easily reduce the first part of the proposition to the following statements. If $p_0, p_1 \in \mathbb{Z}_{\geq 1}$ and $p_0 + p_1 = r - 1$, then $F_{p_0}F_{p_1} > 0$ and $F_{p_0-1}F_{p_1-1} \geq 0$. If $p_0, p_1 \in \mathbb{Z}_{\geq 1}$ and $p_0 + p_1 = r - 2$, then $2F_{p_0-2}F_{p_1-2} \geq 0$. If $p_0, p_2 \in \mathbb{Z}_{\geq 1}$, $p_1 \in \mathbb{Z}_{\geq 0}$ and $p_0 + p_1 + p_2 = r - 3$, then $F_{p_1}(F_{p_0-1}F_{p_2-1} + 3F_{p_0-2}F_{p_2-1} + 3F_{p_0-1}F_{p_2-2}) + 2F_{p_0-2}F_{p_1+1}F_{p_2-2} \geq 0$. These statements follow from the facts that $F_n > 0$ for $n \geq 1$, and $F_n \geq 0$ for $n \geq -1$.

Now we prove the last part. If $\underline{l} \in E_{r-1,0}$, then $K(\underline{l}) = \kappa_r = F_{r+1} < F_{r+1} + F_{r-1}$. Since $E_{r,2} = U_r \cup V_r$, the statement follows from $F_{r+1} + F_{r-1} - 2F_{r-4} < F_{r+1} + F_{r-1}$ and $F_{r+1} + F_{r-1} - F_{r-7} \leq F_{r+1} + F_{r-1}$. Note that here $F_{r-7} > 0$ if $r \geq 6$ and $r \neq 7$. \square

Now we are ready to prove our main result.

Proof of Theorem 1. The second part of the theorem follows from the first part by Lemma 4. We prove now the first part. For $r \in \{0, 1, 2, 3, 4, 5\}$ one could check the statement by hand. The attached Mathematica file contains a program that does this.

Suppose that $r \geq 6$, and let $m \geq 1$. By Corollary 11, $L_m(r)$ is the m th largest distinct value in $\{K(\underline{l}); \underline{l} \in E'_r\}$. If $\underline{l} \in E'_r \setminus (E_{r,0} \cup E_{r,1})$, then

$$K(\underline{l}) \leq \min(K(\underline{l}'); \underline{l}' \in E_{r,1})$$

by Proposition 13 and Corollary 15. Moreover

$$\{K(\underline{l}); \underline{l} \in E_{r,0}\} = \{F_{r+2}\}$$

and

$$\{K(\underline{l}); \underline{l} \in E_{r,1}\} = \{F_{r+2} - F_i F_j; i, j \in \mathbb{Z}_{\geq 1}, i + j = r - 1\}$$

by Lemma 14. So $\{L_1(r), \dots, L_{|H|}(r)\} = H$, where

$$H = \{F_{r+2} - F_i F_j; i, j \in \mathbb{Z}_{\geq 0}, i + j = r - 1\}.$$

Here $|H| = \lceil \frac{r}{2} \rceil$ by Lemma 4. \square

6 Further research

Using our results, it would not be hard to describe the exact positions where the first $\lceil \frac{r}{2} \rceil$ largest values in the r th row of Stern's diatomic sequence appear.

To determine $L_m(r)$ for $1 \leq m \leq \lceil \frac{r}{2} \rceil$, we needed to calculate and (sometimes) compare the values of $K(\underline{l})$ for $\underline{l} \in E_{r,0} \cup E_{r,1} \cup E_{r,2}$. To go a step further, i.e., to determine $L_m(r)$ for some $m > \lceil \frac{r}{2} \rceil$, we would probably need to calculate and compare the values of $K(\underline{l})$ for $\underline{l} \in \bigcup_{i=0}^3 E_{r,i}$. For example, we have ordered $\{K(\underline{l}); \underline{l} \in E_{r,0} \cup E_{r,1}\}$, but we have not yet ordered $\{K(\underline{l}); \underline{l} \in E_{r,2}\}$.

Instead of studying Stern's diatomic array, which starts with the 0th row 1, 1, we could study the following generalization. Start with the 0th row a, b , where $a, b \in \mathbb{R}$, and in each step construct a new row by copying the last row, and writing between each two consecutive elements their sum. One could try to understand the largest values in the r th row of this array.

7 Acknowledgments

The work presented here was carried out at the University of Salzburg, where I was supported by the Austrian Science Fund (FWF): P24574.

References

- [1] C. Defant, Upper bounds for Stern's diatomic sequence and related sequences, preprint, <http://arxiv.org/abs/1506.07824>, 2015.
- [2] L. E. Dickson, *History of the Theory of Numbers, Vol. I: Divisibility and Primality*. Chelsea Publishing Co., 1966.
- [3] J. Lansing, Largest values for the Stern sequence, *J. Integer Seq.*, **17** (2014), [Article 14.7.5](#).
- [4] D. H. Lehmer, On Stern's diatomic series, *Amer. Math. Monthly*, **36** (1929), 59–67.
- [5] E. Lucas, Sur les suites de Farey, *Bull. Soc. Math. France*, **6** (1878), 118–119.
- [6] G. Ramharter, Extremal values of continuants, *Proc. Amer. Math. Soc.*, **89** (1983), 189–201.
- [7] M. Stern, Über eine zahlentheoretische Funktion, *J. Reine Angew. Math.*, **55** (1858), 193–220.
- [8] Wolfram Research, Inc., Mathematica, Version 10.0, 2014.

2010 *Mathematics Subject Classification*: Primary 11A55; Secondary 11B39.

Keywords: Stern sequence, alternating binary expansion, continuant.

(Concerned with sequence [A002487](#).)

Received September 13 2016; revised version received November 15 2016. Published in *Journal of Integer Sequences*, December 27 2016.

Return to [Journal of Integer Sequences home page](#).