



The Fibonacci Sequence and Schreier-Zeckendorf Sets

Hùng Việt Chu

Department of Mathematics

University of Illinois at Urbana-Champaign

Champaign, IL 61820

USA

hungchu2@illinois.edu

Abstract

A finite subset of the natural numbers is *weak-Schreier* if $\min S \geq |S|$, *strong-Schreier* if $\min S > |S|$, and *maximal* if $\min S = |S|$. Let M_n be the number of weak-Schreier sets with n being the largest element and $(F_n)_{n \geq -1}$ denote the Fibonacci sequence. A finite set is said to be Zeckendorf if it does not contain two consecutive natural numbers. Let E_n be the number of Zeckendorf subsets of $\{1, 2, \dots, n\}$. It is well-known that $E_n = F_{n+2}$. In this paper, we first show four other ways to generate the Fibonacci sequence from counting Schreier sets. For example, let C_n be the number of weak-Schreier subsets of $\{1, 2, \dots, n\}$. Then $C_n = F_{n+2}$. To understand why $C_n = E_n$, we provide a bijective mapping to prove the equality directly. Next, we prove linear recurrence relations among the number of Schreier-Zeckendorf sets. Lastly, we discover the Fibonacci sequence by counting the number of subsets of $\{1, 2, \dots, n\}$ such that two consecutive elements in increasing order always differ by an odd number.

1 Background and main results

Let the Fibonacci sequence be $F_{-1} = 1$, $F_0 = 0$, and $F_m = F_{m-1} + F_{m-2}$ for all $m \geq 1$. We only concern ourselves with finite subsets of natural numbers greater than 0 and use \mathbb{N} for the set $\{1, 2, 3, \dots\}$. We define a set to be

- *weak-Schreier* if $\min S \geq |S|$,

- *strong-Schreier* if $\min S > |S|$ and
- *maximal* if $\min S = |S|$,

where $|S|$ is the cardinality of set S . Schreier sets are named after Schreier who defined them to solve a problem in Banach space theory in 1930 [10]. These sets were also independently discovered in combinatorics and are connected to Ramsey-type theorems for subsets of \mathbb{N} . For each $n \in \mathbb{N}$, let M_n be the number of weak-Schreier sets with n being the largest element. In notation,

$$M_n = |\{S \subseteq \mathbb{N} : \min S \geq |S| \text{ and } \max S = n\}|.$$

The first few values of M_n are 1, 1, 2, 3, 5, 8, 13, 21, 34, \dots ; indeed, it is known that $M_n = F_n$ for all n [12]. However, the author is unable to locate the first person to prove this result. If we look at either strong-Schreier sets or maximal sets instead, we can also generate the Fibonacci sequence. Let

- A_n be the number of strong-Schreier sets S with $\max S = n$,
- B_n be the number of maximal sets S with $\max S = n$,
- C_n be the number of weak-Schreier subsets of $\{1, 2, \dots, n\}$ (including the empty set),
- D_n be the number of strong-Schreier subsets of $\{1, 2, \dots, n\}$ (including the empty set).

For our sequence $(C_n)_{n \geq 1}$ and $(D_n)_{n \geq 1}$, we relax the condition about the maximum of our sets. Clearly, for each $n \in \mathbb{N}$, $M_n = A_n + B_n$, $C_n = \sum_{k=1}^n M_k + 1$ and $D_n = \sum_{k=1}^n A_k + 1$.

Theorem 1. *For each $n \in \mathbb{N}$, we have $A_n = F_{n-1}$, $B_n = F_{n-2}$, $C_n = F_{n+2}$ and $D_n = F_{n+1}$*

The Fibonacci representation of natural numbers was first studied by Ostrowski [9] and Lekkerkerker [8]. In 1972, Zeckendorf proved that every positive integer can be uniquely written as a sum of non-consecutive Fibonacci numbers [11]. Since then, many papers have generalized this result and explored properties of the Zeckendorf decomposition: see [1, 2, 3, 4, 5, 6, 8]. We instead focus on the important requirement for uniqueness of the Zeckendorf decomposition; that is, our set contains no two consecutive Fibonacci numbers. We give the same definition for natural numbers.

Definition 2. A finite set of natural numbers is Zeckendorf if the set does not contain two consecutive natural numbers.

Let E_n be the number of subsets of $\{1, 2, \dots, n\}$ that satisfy the Zeckendorf condition. It is well-known that $E_n = F_{n+2}$.

Two different ways of counting subsets of $\{1, 2, \dots, n\}$ give the same number; that is, $C_n = E_n$. To understand the connection, we construct a bijective mapping to show that $C_n = E_n$ directly. Our proof is independent of the fact that $C_n = E_n = F_{n+2}$ and thus, provides insight into the seemingly mysterious equality.

Theorem 3. For each $n \in \mathbb{N}$, $C_n = E_n$.

Next, a natural question is about sequences formed by the number of sets that satisfy both the Schreier and the Zeckendorf conditions. In particular, we say that a set satisfies the k -Zeckendorf condition if two arbitrary numbers in the set are at least k apart. We discover linear recurrence relations among the number of sets satisfying both the Schreier and the k -Zeckendorf conditions.

For each $n \in \mathbb{N}$, let $H_{k,n}$ be the number of subsets of $\{1, 2, \dots, n\}$ that

- (1) satisfy the k -Zeckendorf condition;
- (2) contain n ; and
- (3) are weak-Schreier.

Theorem 4. Fix $k \in \mathbb{N}_{\geq 2}$. We have

$$H_{k,n} = \begin{cases} 1, & \text{if } 1 \leq n \leq k+1; \\ H_{k,n-1} + H_{k,n-(k+1)}, & \text{if } n > k+1. \end{cases}$$

Using the exact same argument as in the proof of Theorem 4, we can also deduce the following theorems regarding strong and maximal Schreier sets. For each $n \in \mathbb{N}$, let $I_{k,n}$ be the number of subsets of $\{1, 2, \dots, n\}$ that (1) satisfy the k -Zeckendorf condition, (2) contain n , and (3) are strong-Schreier.

Theorem 5. Fix $k \in \mathbb{N}_{\geq 2}$. We have

$$I_{k,n} = \begin{cases} 0, & \text{if } n = 1; \\ 1, & \text{if } 2 \leq n \leq k+2; \\ I_{k,n-1} + I_{k,n-(k+1)}, & \text{if } n > k+2. \end{cases}$$

For each $n \in \mathbb{N}$, let $J_{k,n}$ be the number of subsets of $\{1, 2, \dots, n\}$ that

- (1) satisfy the k -Zeckendorf condition;
- (2) contain n ; and
- (3) are maximal.

Theorem 6. Fix $k \in \mathbb{N}_{\geq 2}$. We have

$$J_{k,n} = \begin{cases} 1, & \text{if } n = 1; \\ 0, & \text{if } 2 \leq n \leq k+1; \\ 1, & \text{if } k+1 < n \leq 2k+2; \\ J_{k,n-1} + J_{k,n-(k+1)}, & \text{if } n > 2k+2. \end{cases}$$

We give the following definition that is useful for the statement of our last result.

Definition 7. Let $A = \{a_1, a_2, \dots, a_k\}$ ($a_1 < a_2 < \dots < a_k$) for some $k \in \mathbb{N}_{\geq 2}$. The difference set of A is $\{a_2 - a_1, a_3 - a_2, \dots, a_k - a_{k-1}\}$. The empty set and a set with exactly one element do not have a difference set.

We end with the following small result.

Theorem 8. Fix $n \in \mathbb{N}$. The number of subsets of $\{1, 2, \dots, n\}$

1. that contain n and whose difference sets contain only odd numbers is F_{n+1} ,
2. whose difference sets contain only odd numbers (the empty set and sets with exactly one element vacuously satisfy this requirement) is $F_{n+3} - 1$.

2 Proof of Theorem 1

Proof of Theorem 1. We first prove item (1). Simple computation gives $A_1 = 0 = F_0$, $A_2 = 1 = F_1$, $A_3 = 1 = F_2$, $A_4 = 2 = F_3$, and $A_5 = 3 = F_4$. It suffices to prove that $A_n + A_{n+1} = A_{n+2}$ for $n \geq 4$. Fix $n \geq 4$ and let us find a formula for A_n . The minimum number k in our sets can take values from 1 to n . For each value of k , there are $n - k - 1$ numbers strictly between k and n . Because our sets are strong-Schreier, they contain at most $k - 3$ numbers out of these $n - k - 1$ numbers. Hence, our formula for A_n is

$$A_n = \sum_{k=1}^{n-1} \sum_{j=0}^{k-3} \binom{n-k-1}{j} + 1.$$

Note that the number 1 in our formula accounts for the set $\{n\}$. It remains to show that $A_n + A_{n+1} = A_{n+2}$ or equivalently, $A_{n+2} - A_{n+1} = A_n$ for $n \geq 4$. We have

$$\begin{aligned} A_{n+2} - A_{n+1} &= \sum_{k=1}^{n+1} \sum_{j=0}^{k-3} \binom{n-k+1}{j} - \sum_{k=1}^n \sum_{j=0}^{k-3} \binom{n-k}{j} \\ &= \sum_{k=1}^n \sum_{j=0}^{k-3} \left(\binom{n-k+1}{j} - \binom{n-k}{j} \right) + \sum_{j=0}^{n-2} \binom{0}{j} \\ &= \sum_{k=1}^n \sum_{j=1}^{k-3} \binom{n-k}{j-1} + 1. \end{aligned}$$

Therefore,

$$\begin{aligned} A_{n+2} - A_{n+1} - A_n &= \sum_{k=1}^n \sum_{j=1}^{k-3} \binom{n-k}{j-1} - \sum_{k=1}^{n-1} \sum_{j=0}^{k-3} \binom{n-k-1}{j} \\ &= \sum_{k=4}^n \sum_{j=1}^{k-3} \binom{n-k}{j-1} - \sum_{k=3}^{n-1} \sum_{j=0}^{k-3} \binom{n-k-1}{j} = 0. \end{aligned}$$

The last equality is because for each $4 \leq t \leq n$, we have $\sum_{j=1}^{t-3} \binom{n-t}{j-1} = \sum_{j=0}^{(t-1)-3} \binom{n-(t-1)-1}{j}$. Hence, $A_{n+2} = A_{n+1} + A_n$ and we are done.

Next, we prove item (2), which follows immediately from item (1). We know that

$$B_n = M_n - A_n = F_n - F_{n-1} = F_{n-2}.$$

We prove item (3). Fix $n \geq 1$. We have

$$C_n = \sum_{k=1}^n M_k + 1 = \sum_{k=1}^n F_k + 1 = (F_{n+2} - 1) + 1 = F_{n+2},$$

as desired. The number 1 accounts for the empty set. The fact that $\sum_{k=1}^n F_k = F_{n+2} - 1$ is due to Lucas [7, p. 4].

Similarly, we prove item (4). Fix $n \geq 1$. We have

$$D_n = \sum_{k=1}^n A_k + 1 = \sum_{k=1}^n F_{k-1} + 1 = (F_{n+1} - 1) + 1 = F_{n+1}.$$

We complete our proof of Theorem 1. □

Let L_n^w be the number of weak-Schreier sets as subsets of $\{1, 2, \dots, n\}$ with an even maximum.

Corollary 9. *For each $n \in \mathbb{N}$,*

$$L_n^w = \begin{cases} F_n, & \text{if } n \text{ is odd;} \\ F_{n+1}, & \text{if } n \text{ is even.} \end{cases}$$

Proof. We have

$$L_n^w = \sum_{\substack{1 \leq k \leq n \\ 2|k}} M_k + 1 = \sum_{\substack{1 \leq k \leq n \\ 2|k}} F_k + 1.$$

The number 1 accounts for the empty set.

If n is even,

$$L_n^w = \sum_{\substack{1 \leq k \leq n \\ 2 \nmid k}} F_k + 1 = (F_{n+1} - 1) + 1 = F_{n+1}.$$

If n is odd,

$$L_n^w = \sum_{\substack{1 \leq k \leq n \\ 2 \nmid k}} F_k + 1 = (F_n - 1) + 1 = F_n.$$

□

Let L_n^s be the number of strong-Schreier sets as subsets of $\{1, 2, \dots, n\}$ with an odd maximum.

Corollary 10. *For each $n \in \mathbb{N}$,*

$$L_n^s = \begin{cases} F_n, & \text{if } n \text{ is odd;} \\ F_{n-1}, & \text{if } n \text{ is even.} \end{cases}$$

Proof. We have

$$L_n^s = \sum_{\substack{1 \leq k \leq n \\ 2 \nmid k}} A_k + 1 = \sum_{1 \leq k \leq n, 2 \nmid k} F_{k-1} + 1.$$

If n is even,

$$L_n^s = \sum_{\substack{1 \leq k \leq n \\ 2 \nmid k}} F_{k-1} + 1 = (F_{n-1} - 1) + 1 = F_{n-1}.$$

If n is odd,

$$L_n^s = \sum_{\substack{1 \leq k \leq n \\ 2 \nmid k}} F_{k-1} + 1 = (F_n - 1) + 1 = F_n.$$

□

3 Proof of Theorem 3 — Explanation of the mysterious identity

Recall that C_n is the number of weak-Schreier sets as subsets of $\{1, 2, \dots, n\}$, while E_n is the number of subsets of $\{1, 2, \dots, n\}$ that do not contain two consecutive numbers. At the first glance, C_n and E_n are little related, so it is surprising to see that $C_n = E_n$ for all $n \in \mathbb{N}$.

For each $n \in \mathbb{N}$, let X_n denote the set of weak-Schreier sets as subsets of $\{1, 2, \dots, n\}$ and let Y_n denote the set of subsets of $\{1, 2, \dots, n\}$ that do not contain two consecutive numbers. In this section, we construct a bijective function $f : X_n \rightarrow Y_n$ to prove that $|X_n| = |Y_n|$.

Proof of Theorem 3. Fix $n \in \mathbb{N}$. Let $A = \{a_1, a_2, \dots, a_{k-1}, a_k\}$ ($a_1 < a_2 < \dots < a_k$) be a weak-Schreier subset of $\{1, 2, \dots, n\}$. Our mapping f acts on A as follows

$$f(A) = f(\{a_1, a_2, \dots, a_{k-1}, a_k\}) = \{a_1 - (k-1), a_2 - (k-2), \dots, a_{k-1} - 1, a_k\}.$$

Define $f(\emptyset) = \emptyset$. To show that f is well-defined, we show that $\{a_1 - (k-1), a_2 - (k-2), \dots, a_{k-1} - 1, a_k\}$ is in Y_n . Because A is weak-Schreier, $k \leq a_1 < a_2 < \dots < a_k$. Hence,

$$1 \leq a_1 - (k-1) < a_2 - (k-2) < \dots < a_{k-1} - 1 < a_k \leq n.$$

Let $t_i = a_i - (k-i)$ for $1 \leq i \leq k$. If $k = 1$, then $\{t_1\}$ is clearly in Y_n . If $k \geq 2$, then for each $2 \leq i \leq k$, we have

$$t_i - t_{i-1} = (a_i - (k-i)) - (a_{i-1} - (k-(i-1))) = (a_i - a_{i-1}) + 1 \geq 2.$$

Therefore, $\{t_1, t_2, \dots, t_k\} \in Y_n$. So, f is well-defined.

Next, we prove that f is injective. Suppose that $f(A) = f(B)$. Let $A = \{a_1, a_2, \dots, a_k\}$ and $B = \{b_1, b_2, \dots, b_k\}$, where $a_1 < a_2 < \dots < a_k$ and $b_1 < b_2 < \dots < b_k$. Because

$$\begin{aligned} a_1 - (k-1) &< a_2 - (k-2) < \dots < a_{k-1} - 1 < a_k, \\ b_1 - (k-1) &< b_2 - (k-2) < \dots < b_{k-1} - 1 < b_k, \end{aligned}$$

we know that $f(A) = f(B)$ implies $a_i - (k-i) = b_i - (k-i)$ for all $1 \leq i \leq k$. Hence, $a_i = b_i$, which shows that $A = B$. Therefore, f is injective.

Finally, we prove that f is surjective. Let $C = \{c_1, c_2, \dots, c_k\} \in Y_n$ be chosen, where $c_1 < c_2 < \dots < c_k$. We claim that

$$D = \{c_1 + (k-1), c_2 + (k-2), \dots, c_{k-1} + 1, c_k\}$$

satisfies $f(D) = C$ and $D \in X_n$. Because C do not contain two consecutive numbers, we know that

$$k \leq c_1 + (k-1) < c_2 + (k-2) < \dots < c_{k-1} + 1 < c_k \leq n.$$

Hence, $D \in X_n$.

We have shown that f is both well-defined and bijective. Therefore, $|X| = |Y|$ or $C_n = E_n$, as desired. \square

Remark 11. We would like to discuss the motivation for the bijection f used in the proof of Theorem 3. Let A be a Schreier set. The map f serves to increase the gap between adjacent elements of A by 1, thus fulfilling the Zeckendorf condition that adjacent elements differ by at least 2. Furthermore, the weak-Schreier condition that $\min A \geq |A|$ ensures that the resulting set is in $\{1, 2, \dots, n\}$.

4 Proof of Theorem 4

Before we prove Theorem 4, we need a simple proposition.

Proposition 12. *For $n, k \in \mathbb{Z}$, the following claims hold.*

1. If $\lfloor \frac{n-2}{k+1} \rfloor = \lfloor \frac{n-k-2}{k+1} \rfloor$, then $\lfloor \frac{n-1}{k+1} \rfloor = \lfloor \frac{n-2}{k+1} \rfloor + 1$.
2. If $\lfloor \frac{n-2}{k+1} \rfloor > \lfloor \frac{n-k-2}{k+1} \rfloor$, then $\lfloor \frac{n-1}{k+1} \rfloor < \lfloor \frac{n-2}{k+1} \rfloor + 1$.
3. If $\lfloor \frac{n-k-2}{k+1} \rfloor = \lfloor \frac{n-2}{k+1} \rfloor$, then $\frac{n-k-2}{k+1} = \lfloor \frac{n-2}{k+1} \rfloor$.

Proof. We prove claim (1). We have

$$\left\lfloor \frac{n-2}{k+1} \right\rfloor = \left\lfloor \frac{n-k-2}{k+1} \right\rfloor = \left\lfloor \frac{n-1}{k+1} - 1 \right\rfloor = \left\lfloor \frac{n-1}{k+1} \right\rfloor - 1.$$

Therefore,

$$\left\lfloor \frac{n-1}{k+1} \right\rfloor = \left\lfloor \frac{n-2}{k+1} \right\rfloor + 1.$$

Next, we prove claim (2). We have

$$\left\lfloor \frac{n-2}{k+1} \right\rfloor > \left\lfloor \frac{n-k-2}{k+1} \right\rfloor = \left\lfloor \frac{n-1}{k+1} - 1 \right\rfloor = \left\lfloor \frac{n-1}{k+1} \right\rfloor - 1.$$

Therefore,

$$\left\lfloor \frac{n-1}{k+1} \right\rfloor < \left\lfloor \frac{n-2}{k+1} \right\rfloor + 1.$$

Lastly, we prove claim (3). Write $n - k - 2 = (k + 1)p + q$ for some $0 \leq q \leq k$. Then

$$\frac{n-2}{k+1} = \frac{(k+1)p + q + k}{k+1} = p + \frac{q+k}{k+1} = p + 1 + \frac{q-1}{k+1}.$$

If $q \geq 1$, then $\lfloor \frac{n-2}{k+1} \rfloor = p + 1 > p = \lfloor \frac{n-k-2}{k+1} \rfloor$, a contradiction. So, $q = 0$, implying that $\frac{n-k-2}{k+1} = \lfloor \frac{n-2}{k+1} \rfloor$. \square

The following lemma is from [6, Lemma 2.1] by Kologl et al.

Lemma 13. *The number of solutions to $y_1 + \dots + y_p = n$ with $y_i \geq c_i$ (each c_i a non-negative integer) is $\binom{n - (c_1 + \dots + c_p) + p - 1}{p - 1}$.*

Proof of Theorem 4. Fix $k \geq 2$. We now find a formula for $H_{k,n}$ for all $n \in \mathbb{N}$. Fix $1 \leq \ell \leq n - 1$. Suppose that the set $\{a_1, \dots, a_\ell, n\}$ satisfies all of our requirements. (For $\ell = 0$, we have the set $\{n\}$.) In particular,

1. $a_1 \geq \ell + 1$,

2. $d_i = a_{i+1} - a_i \geq k$ and $d_\ell = n - a_\ell \geq k$.

Note that

$$a_1 + \sum_{i=1}^{\ell} d_i = n. \quad (1)$$

By Lemma 13, the number of sets satisfying Equation (1) is

$$\binom{n - (\ell + 1 + k\ell) + (\ell + 1) - 1}{(\ell + 1) - 1} = \binom{n - k\ell - 1}{\ell}.$$

Therefore, the number of sets containing n that are k -Zeckendorf and weak-Schreier is

$$H_{k,n} = \sum_{\ell=1}^{\lfloor \frac{n-1}{k+1} \rfloor} \binom{n - k\ell - 1}{\ell} + 1.$$

The number 1 accounts for the set $\{n\}$ and we only let ℓ run up to $\lfloor \frac{n-1}{k+1} \rfloor$ to make sure that $n - k\ell - 1 \geq \ell$. It can be easily verified that $H_{k,n} = 1$ for $1 \leq n \leq k + 1$ because $\lfloor \frac{n-1}{k+1} \rfloor = 0$ for $1 \leq n \leq k + 1$. It suffices to show that for $n \geq k + 2$, $H_{k,n} = H_{k,n-1} + H_{k,n-(k+1)}$. Equivalently,

$$\sum_{\ell=1}^{\lfloor \frac{n-1}{k+1} \rfloor} \binom{n - k\ell - 1}{\ell} = \sum_{\ell=1}^{\lfloor \frac{n-2}{k+1} \rfloor} \binom{n - k\ell - 2}{\ell} + \sum_{\ell=1}^{\lfloor \frac{n-(k+1)-1}{k+1} \rfloor} \binom{n - k\ell - 1 - (k+1)}{\ell} + 1. \quad (2)$$

Equivalently, noting that the +1 term cancels with the $\ell = 1$ term in the left hand side summation

$$\begin{aligned} & \sum_{\ell=2}^{\lfloor \frac{n-2}{k+1} \rfloor} \left(\binom{n - k\ell - 1}{\ell} - \binom{n - k\ell - 2}{\ell} \right) + \sum_{\substack{\lfloor \frac{n-1}{k+1} \rfloor \\ \lfloor \frac{n-2}{k+1} \rfloor + 1}} \binom{n - k\ell - 1}{\ell} \\ &= \sum_{\ell=1}^{\lfloor \frac{n-k-2}{k+1} \rfloor} \binom{n - k(\ell + 1) - 2}{\ell}. \end{aligned} \quad (3)$$

We can simplify Equation (3) further by applying the binomial coefficient recurrence

$$\sum_{\ell=2}^{\lfloor \frac{n-2}{k+1} \rfloor} \binom{n - k\ell - 2}{\ell - 1} + \sum_{\substack{\lfloor \frac{n-1}{k+1} \rfloor \\ \lfloor \frac{n-2}{k+1} \rfloor + 1}} \binom{n - k\ell - 1}{\ell} = \sum_{\ell=1}^{\lfloor \frac{n-k-2}{k+1} \rfloor} \binom{n - k(\ell + 1) - 2}{\ell}.$$

Reindexing ℓ in the first summation, we have

$$\sum_{\ell=1}^{\lfloor \frac{n-2}{k+1} \rfloor - 1} \binom{n - k(\ell + 1) - 2}{\ell} + \sum_{\lfloor \frac{n-2}{k+1} \rfloor + 1}^{\lfloor \frac{n-1}{k+1} \rfloor} \binom{n - k\ell - 1}{\ell} = \sum_{\ell=1}^{\lfloor \frac{n-k-2}{k+1} \rfloor} \binom{n - k(\ell + 1) - 2}{\ell}.$$

Subtract the first summation from both sides to have

$$\sum_{\lfloor \frac{n-2}{k+1} \rfloor + 1}^{\lfloor \frac{n-1}{k+1} \rfloor} \binom{n - k\ell - 1}{\ell} = \sum_{\ell=\lfloor \frac{n-2}{k+1} \rfloor}^{\lfloor \frac{n-k-2}{k+1} \rfloor} \binom{n - k(\ell + 1) - 2}{\ell}. \quad (4)$$

We now prove that Equation (4) is correct, which implies that Equation (2) is correct.

Case 1: $\lfloor \frac{n-k-2}{k+1} \rfloor < \lfloor \frac{n-2}{k+1} \rfloor$. Then $\lfloor \frac{n-2}{k+1} \rfloor + 1 > \lfloor \frac{n-1}{k+1} \rfloor$ by Proposition 12. Therefore, two sides of Equation (4) are identically 0.

Case 2: $\lfloor \frac{n-k-2}{k+1} \rfloor = \lfloor \frac{n-2}{k+1} \rfloor$. Then $\lfloor \frac{n-2}{k+1} \rfloor + 1 = \lfloor \frac{n-1}{k+1} \rfloor$ and $\frac{n-k-2}{k+1} = \lfloor \frac{n-2}{k+1} \rfloor$ by Proposition 12. Therefore, the left side of Equation (4) is

$$\binom{n - k(\lfloor \frac{n-2}{k+1} \rfloor + 1) - 1}{\lfloor \frac{n-2}{k+1} \rfloor + 1} = 1$$

because $\frac{n-k-2}{k+1} = \lfloor \frac{n-2}{k+1} \rfloor$. Similarly, the right side is also equal to 1.

In both cases, Equation (4) is correct. This completes our proof. \square

5 Proof of Theorem 8—A new way to generate the Fibonacci sequence

Proof of Theorem 8. First, we prove item (1). Let P_n be the number of subsets of $\{1, 2, \dots, n\}$ that contain n and whose difference sets contain only odd numbers.

Base cases: For $n = 1$, we have $\{1\}$ to be the only subset of $\{1\}$ that satisfies our requirement. So, $P_1 = 1 = F_2$. For $n = 2$, we have $\{2\}$ and $\{1, 2\}$ to be the only two subsets of $\{1, 2\}$ that satisfy our requirement. So, $P_2 = 2 = F_3$.

Inductive hypothesis: Suppose that there exists $k \geq 2$ such that for all $n \leq k$, $P_n = F_{n+1}$. We show that $P_{k+1} = F_{k+2}$. Let O_n denote the set of subsets of $\{1, 2, \dots, n\}$ that satisfy our requirement. Observe that unioning a set in O_{n-1-2i} (for $i \geq 0$) with n produces a set in O_n and any set in O_n is of the form of a set in O_{n-1-2i} plus the element n . Therefore,

$$P_{k+1} = |O_{k+1}| = \sum_{\substack{1 \leq i \leq k \\ 2 \nmid i}} |O_{k+1-i}| + 1 = P_k + \sum_{\substack{3 \leq i \leq k \\ 2 \nmid i}} |O_{k+1-i}| + 1.$$

The number 1 accounts for the set $\{n\}$. If k is odd,

$$\begin{aligned} \sum_{\substack{3 \leq i \leq k \\ 2 \nmid i}} |O_{k+1-i}| &= |O_1| + |O_3| + \cdots + |O_{k-2}| \\ &= |F_2| + |F_4| + \cdots + |F_{k-1}| = F_k - 1 = P_{k-1} - 1. \end{aligned}$$

If k is even,

$$\begin{aligned} \sum_{\substack{3 \leq i \leq k \\ 2 \nmid i}} |O_{k+1-i}| &= |O_2| + |O_4| + \cdots + |O_{k-2}| \\ &= |F_3| + |F_5| + \cdots + |F_{k-1}| = F_k - 1 = P_{k-1} - 1. \end{aligned}$$

In both cases, we have $\sum_{\substack{3 \leq i \leq k \\ 2 \nmid i}} |O_{k+1-i}| = P_{k-1} - 1$. Therefore, $P_{k+1} = P_k + P_{k-1} = F_{k+1} + F_k = F_{k+2}$, as desired.

Next, we prove item (2). Let Q_n be the number of subsets of $\{1, 2, \dots, n\}$ whose difference sets contain only odd numbers is Q_n (the empty set and sets with exactly one element vacuously satisfy this requirement). Note that by definition of P_n and Q_n , we have

$$Q_n = 1 + \sum_{k=1}^n |P_k| = 1 + \sum_{k=1}^n F_{k+1} = \sum_{k=1}^{n+1} F_k = F_{n+3} - 1,$$

as desired. (The +1 before the first summation accounts for the empty set.) □

6 Acknowledgments

The author would like to thank the anonymous referee and the editor for various helpful comments that clarify several points made in this paper. Thanks to Kevin Beanland at Washington and Lee University for useful comments on earlier drafts.

References

- [1] E. Burger, D. Clyde, C. Colbert, G. Shin, and Z. Wang, Canonical diophantine representations of natural numbers with respect to quadratic “bases”, *J. Number Theory* **133** (2013), 1372–1388.
- [2] O. Beckwith, A. Bower, L. Gaudet, R. Insoft, S. Li, S. Miller, and P. Tosteson, The average gap distribution for generalized Zeckendorf decompositions, *Fibonacci Quart.* **51** (2013), 13–27.
- [3] A. Best, P. Dynes, X. Edelsbrunner, B. McDonald, S. Miller, C. Turnage-Butterbaugh, and M. Weinstein, Benford behavior of Zeckendorf decompositions, *Fibonacci Quart.* **52** (2014), 35–46.

- [4] P. Demontigny, T. Do, A. Kulkarni, S. Miller, D. Moon, and U. Varma, Generalizing Zeckendorf's theorem to f -decompositions, *J. Number Theory* **141** (2014), 136–158.
- [5] T. J. Keller, Generalizations of Zeckendorf's theorem, *Fibonacci Quart.* **10** (1972), 95–102.
- [6] G. S. Kopp, M. Kologlu, S. Miller, and Y. Wang, On the number of summands in Zeckendorf decompositions, *Fibonacci Quart.* **49** (2011), 116–130.
- [7] E. Lucas. *Theorie Des Nombres*, Gauthier-Villars, 1891.
- [8] C. G. Lekkerkerker, Voorstelling van natuurlyke getallen door een som van getallen van Fibonacci, *Simon Stevin* **29** (1951–1952), 190–195.
- [9] A. Ostrowski, *Bemerkungen zur Theorie der diophantischen Approximationen*, Hambg. Abh. **1** (1922), 77–98.
- [10] J. Schreier, Ein Gegenbeispiel zur Theorie der schwachen Konvergenz, *Studia Math.* **2** (1962), 58–62.
- [11] E. Zeckendorf, Representation des nombres naturels par une somme de nombres de Fibonacci ou de nombres de Lucas, *Bull. Soc. Roy. Sci. Liege* **41** (1972), 179–182.
- [12] Unknown author, Jozef Schreier, Schreier sets and the Fibonacci sequence, 2012. Available at <https://outofthenormmaths.wordpress.com/2012/05/13/jozef-schreier-schreier-sets-and-the-fibonacci-sequence/>.

2010 *Mathematics Subject Classification*: 11B39.

Keywords: Fibonacci sequence, linear recurrence, combinatorics, Schreier set.

(Concerned with sequences [A000045](#).)

Received June 26 2019; revised version received August 31 2019; September 2 2019. Published in *Journal of Integer Sequences*, September 3 2019.

Return to [Journal of Integer Sequences home page](#).