Journal of Integer Sequences, Vol. 24 (2021),
Article 21.7.2

Redundance in the Signed m-Bonacci Numeration System

Zuzana Masáková and Élise Vandomme
Department of Mathematics FNSPE
Czech Technical University in Prague
Trojanova 13
120 00 Praha 2
Czech Republic
zuzana.masakova@fjfi.cvut.cz

Abstract

We study redundance in the signed m-bonacci numeration system. We determine the number of representations of 0 in this system and describe the automaton accepting the language of such representations. We reveal a surprising connection between Fibonacci and Tribonacci numbers.

1 Introduction

Integers are usually represented as sums of powers of a base $q \in \mathbb{N}$ for $q \geq 2$. In particular, every $x \in \mathbb{N}$ can be written as $x = \sum_{j=1}^{n} a_j q^{j-1}$. If $a_j \in \{0, 1, \ldots, q-1\}$ and $a_n \neq 0$, then such a representation of x is unique. One can replace the sequence $(q^{j-1})_{j\geq 1}$ of powers of the base by any linear recurrence sequence $(G_j)_{j\geq 1}$. For details about such representations for a general linear recurrence sequence (G_j) and conditions on uniqueness, see [20] or [18, Chapter 7].

Perhaps the first example of a possibility of representing integers in these so-called linear numeration systems is the well-known Zeckendorf numeration system [16], where for (G_j) we take the sequence of Fibonacci numbers $G_j = F_j$, where $F_1 = 1$, $F_2 = 2$, $F_{j+2} = F_{j+1} + F_j$ for $j \geq 1$. The representation of an integer x in the form $x = \sum_{j=1}^{n} a_j F^j$ is unique if $a_j \in \{0, 1\}$ where $a_j \cdot a_{j-1} = 0$, i.e., the sequence of coefficients does not contain consecutive 1's. This representation can be obtained by the greedy algorithm. Relaxing the condition on consecutive 1's, the system becomes redundant even with the alphabet of digits $\{0, 1\}$. The function $R(n)$ expressing the number of representations of $n \in \mathbb{N}$ has been studied by several authors. Berstel [2]

provided a formula for calculating $R(n)$ based on the greedy representation of n. A different way of computation was given by Edson and Zamboni [5]. Kocábová et al. [14] and Stockmeyer [22] studied the function $R(n)$ in more detail.

A similar study was performed for different linear recurrence sequences [12]. The m-bonacci sequence as a natural generalization of the Fibonacci case $m = 2$ was considered by Kocábová et al. [15] and Edson and Zamboni [6]. Second-order linear recurrences were studied by Edson [4]. In all these cases, the alphabet of digits was chosen to be the minimal one allowing representation of every integer. In the case of the m-bonacci system, this means the alphabet $\{0, 1\}$.

One can extend the study even more, to considering redundance when the alphabet contains more than the necessary number of digits. Considering such representations may have practical consequences, e.g., for the possibility to define algorithms for parallel addition or algorithms for fast multiplication using non-adjacent form or representations of minimal weight. Parallel addition for a numeration system with integer base $q \geq 3$ was given by Avizienis [1]. For a signed binary system (with the alphabet $\{-1, 0, 1\}$), the algorithm was described by Chow and Robertson [3]. An algorithm for parallel addition in the Fibonacci system with alphabet $\{-1, 0, 1\}$ was given by Frougny et al. [9]. Minimal weight expansions were studied, for example, by Heuberger [13] and Grabner [11]. Similar questions are examined by Frougny and Steiner [10] in systems with real algebraic base β which are directly connected to linear numeration systems [8].

In this paper we study the linear numeration system based on the m-bonacci sequence over the alphabet of digits $\{-1, 0, 1\}$. We show that the number $r(k)$ of representations of 0 in the m-bonacci system over $\{-1, 0, 1\}$ of length at most k is also a linear recurrence sequence and we determine the corresponding recurrence of order $m + 1$. The proof of this result uses the m-bonacci numeration system with the alphabet $\{0, 1, 2\}$. We also present the automaton accepting precisely the representations of 0 over $\{-1, 0, 1\}$.

2 Representations

Definition 1. Let $m \in \mathbb{N}$ with $m \geq 2$. Let $(T_k)_{k\geq 1}$ denote the m-bonacci sequence defined by

$$T_k = 2^{k-1} \text{ for } k \in \{1, \ldots, m\} \text{ and } T_k = \sum_{j=1}^{m} T_{k-j} \text{ for } k > m.$$

Definition 2. Let $A \subset \mathbb{N}$ be an alphabet. A positive integer n is *represented* by $(a_\ell a_{\ell-1} \cdots a_1)$ if $n = \sum_{j=1}^{\ell} a_j T_j$ with $a_j \in A$ and $a_\ell \neq 0$. In that case, ℓ is the *length* of the representation. The number n represented by $(a_\ell a_{\ell-1} \cdots a_1)$ is called the *value* of $(a_\ell a_{\ell-1} \cdots a_1)$ and is denoted by $\mathrm{val}(a_\ell a_{\ell-1} \cdots a_1)$. Let $p_A(n)$ denote the number of representations of n over the alphabet A and by $p_A(n, i)$ the number of representations of n of length exactly i over the alphabet A. If the alphabet is clear from the context, we simply write $p(n)$ and $p(n, i)$.

Our aim is to enumerate the representations of 0 in the m-bonacci system with the alphabet $B = \{-1, 0, 1\}$. Conveniently, we transfer the question to determine

the number of representations in the system with the alphabet $A = \{0, 1, 2\}$ of the number N_k, where $(N_k)_{k\geq 1}$ is the running sum of the sequence $(T_k)_{k\geq 1}$, i.e.,

$$N_k = \sum_{j=1}^{k} T_j \quad \text{for } k \geq 1.$$

We set $N_0 := 0$. In fact, adding 1 to every digit in a representation of 0 over $B = \{-1, 0, 1\}$ of length k, we obtain a representation of N_k over the alphabet $A = \{0, 1, 2\}$. The exact relation of the two systems with alphabets A and B is explained in Section 4.

First, let us present some computer calculations of $p(N_k)$ for the m-bonacci system with $m = 2, 3, 4$, and 5. We count the number of representations of N_k in the m-bonacci numeration system using the alphabet $A = \{0, 1, 2\}$.

$m = 2$		$m = 3$		$m = 4$		$m = 5$	
N_i	$p(N_i)$	N_i	$p(N_i)$	N_i	$p(N_i)$	N_i	$p(N_i)$
1	1	1	1	1	1	1	1
3	2	3	1	3	1	3	1
6	4	7	2	7	1	7	1
11	7	14	4	15	2	15	1
19	13	27	6	30	4	31	2
32	24	51	9	59	6	62	4
53	44	95	15	115	8	123	6
87	81	176	25	223	11	243	8
142	149	325	40	431	17	479	10
231	274	599	64	832	27	943	13
375	504	1103	104	1605	41	1855	19
608	927	2030	169	3095	60	3648	29
985	1705	3735	273	5967	88	7173	43
1595	3136	6871	441	11503	132	14103	61
2582	5768	12639	714	22174	200	27727	84
4179	10609	23248	1156	42743	301	54511	116
6763	19513	42761	1870	82391	449	107167	164
10944	35890	78651	3025	158815	669	210686	236
17709	66012	144663	4895	306127	1001	414199	340
28655	121415	266078	7921	590080	1502	814295	485

Remark 3. Note the surprising fact in the column of results for the case $m = 2$: The number of representations of $N_k = \sum_{j=1}^{k} F_j = F_{k+2} - 2$ is equal to the k-th Tribonacci number T_k. This observation was first made by Vlašic [23]. This statement is included in Proposition 4 as the case $m = 2$.

3 Representations in the m-bonacci system over $\{0, 1, 2\}$

Proposition 4. *Let $A = \{0, 1, 2\}$, $k, m \in \mathbb{N}$ with $m \geq 2$ and $k \geq 1$. The number $p(N_k)$ of representations of N_k in the m-bonacci numeration system over the alphabet A satisfies the recurrence relation*

$$p_A(N_k) = p_A(N_{k-1}) + p_A(N_{k-m}) + p_A(N_{k-m-1}) \text{ for } k > m + 1 \tag{1}$$

with $p_A(N_k) = 1$ for $k \in \{0, \ldots, m-1\}$, $p_A(N_m) = 2$ and $p_A(N_{m+1}) = 4$.

The proof of this result relies on the following lemma.

Lemma 5. *Let $k \in \mathbb{N}$. The following inequalities hold:*

$$N_k < T_k + T_{k+1} \leq T_{k+2} \text{ for } k \geq 1, \tag{2}$$

$$2N_{k-2} < N_k \text{ for } k \geq 3, \tag{3}$$

$$N_k < 2T_{k+1} \text{ for } k \geq 1. \tag{4}$$

Proof. For the first inequality, we proceed by induction. It is easy to check that $N_1 < T_1 + T_2 \leq T_3$. Let $k \geq 2$ and assume the results hold for $k - 1$. We have by definition and induction hypothesis that $N_k = T_k + N_{k-1} < T_k + T_{k+1} \leq T_{k+2}$.

All the other inequalities follow from the first one. For the second inequality with $k \geq 3$, we have

$$N_k = T_k + T_{k-1} + N_{k-2} > N_{k-2} + T_{k-1} + N_{k-2} > 2N_{k-2}.$$

For the last inequality, we just note that $T_k < T_{k+1}$ for all $k \geq 1$. □

Proof of Proposition 4. First note that (2) and (3) imply that every representation of N_k has length in $\{k-1, k, k+1\}$. So

$$p(N_k) = p(N_k, k+1) + p(N_k, k) + p(N_k, k-1) \text{ for } k \geq 1. \tag{5}$$

Obviously, by definition of N_k, (1^k) is one of the representations of N_k.

We first take care of the initial conditions. For $k \in \{1, \ldots, m-1\}$, the number $N_k = 2^k - 1$ has a unique representation that is (1^k). For $k = m$, we have $N_m = 2^m - 1 = T_{m+1}$. Therefore, N_m can be represented only by (1^m) or (10^m). Finally, for $k = m + 1$, we can use the m-bonacci recurrence to write

$$N_{m+1} = \sum_{j=1}^{m+1} T_j = T_{m+2} + T_1 = 2T_{m+1} = 2\sum_{j=1}^{m} T_j.$$

Thus the representations of N_{k+1} are four, namely (1^{m+1}), $(10^m 1)$, (20^m), and (2^m). It is easy to check that these are the only ones. We can summarize it in Table 1.

For $k \geq m + 2$, we will distinguish three cases according to the possible lengths of the representations.

k	$p(N_k)$	$p(N_k, k+1)$	$p(N_k, k)$	$p(N_k, k-1)$
1	1	0	1	0
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$m-1$	1	0	1	0
m	2	1	1	0
$m+1$	4	1	2	1

Table 1: Initial conditions for the sequences $p(N_k)$, $p(N_k, k+1)$, $p(N_k, k)$, and $p(N_k, k-1)$.

Case 1: consider the representations $(a_{k+1} \cdots a_1)$ of N_k of length $k+1$ and assume $k \geq m$. In that case, $a_{k+1} \neq 2$ because otherwise $N_k = \text{val}(2a_k \cdots a_1) \geq 2T_{k+1}$, which is in contradiction with (4). So $a_{k+1} = 1$ and $\text{val}(a_k \cdots a_1) = N_k - T_{k+1} = N_k - \sum_{j=0}^{m-1} T_{k-j} = N_{k-m}$. This means that $p(N_k, k+1)$ is equal to the number of representations of N_{k-m} of length at most k. Since $k-m+1 < k$, this number is equal to $p(N_{k-m})$. Thus

$$p(N_k, k+1) = p(N_{k-m}) \text{ for } k \geq m. \tag{6}$$

Case 2: consider the representations $(a_k \cdots a_1)$ of N_k of length k and assume $k \geq m+1$. If $a_k = 2$, then $\text{val}(a_{k-1} \cdots a_1) = N_k - 2T_k = N_{k-m-1}$. The number of such representations is equal to the number of representations of N_{k-m-1}, and since $k-m < k-1$, this is equal to $p(N_{k-m-1})$. If $a_k = 1$, then $\text{val}(a_{k-1} \cdots a_1) = N_k - T_k = N_{k-1}$. So the number of such representations is equal to $p(N_{k-1}, k-1) + p(N_{k-1}, k-2)$. In conclusion, we get

$$\begin{aligned} p(N_k, k) &= p(N_{k-m-1}) + p(N_{k-1}, k-1) + p(N_{k-1}, k-2) \\ &= p(N_{k-m-1}) + p(N_{k-1}) - p(N_{k-1}, k). \end{aligned}$$

Since $k \geq m+1$, we can use (6) to obtain $p(N_{k-1}, k) = p(N_{k-m-1})$ and thus

$$p(N_k, k) = p(N_{k-1}) \text{ for } k \geq m+1. \tag{7}$$

Case 3: consider the representations $(a_{k-1} \cdots a_1)$ of N_k of length $k-1$ and assume $k \geq m+2$. We have $a_{k-1} = a_{k-2} = \cdots = a_{k-m+1} = 2$. Indeed, let $\ell \in \{1, \ldots, m-1\}$. If the prefix of length $\ell - 1$ of the representation is a concatenation of 2's (i.e., $a_{k-1} = \cdots = a_{k+1-\ell} = 2$), then $a_{k-\ell} = 2$. Otherwise, $a_{k-\ell}$ is equal to 1 or 0 and

$$\begin{aligned} 2N_{k-\ell-1} \geq \text{val}(a_{k-\ell-1} \cdots a_1) &= N_k - 2\sum_{j=1}^{\ell-1} T_{k-j} - a_{k-\ell} T_{k-\ell} \geq \\ &\geq N_k - 2\sum_{j=1}^{\ell-1} T_{k-j} - T_{k-\ell} = N_{k-\ell-1} + T_k - \sum_{j=1}^{\ell-1} T_{k-j} = \\ &= N_{k-\ell-1} + \sum_{j=k-m}^{k-\ell} T_j > 2N_{k-\ell-1} \end{aligned}$$

which is a contradiction. Note that for the last inequality we have used (2) which is justified since $k-\ell-1 \geq k-m \geq 2$.

So the representation of N_k is of the form $(a_{k-1}\cdots a_1) = (2^{m-1}a_{k-m}\cdots a_1)$ and

$$\mathrm{val}(a_{k-m}\cdots a_1) = N_k - 2\sum_{j=k-m+1}^{k-1} T_j = N_{k-m} + T_k - \sum_{j=1}^{m-1} T_{k-j} = N_{k-m} + T_{k-m}.$$

If $a_{k-m} = 2$, then $\mathrm{val}(a_{k-m-1}\cdots a_1) = N_{k-m-1}$. The number of representations of N_k of this form is $p(N_{k-m-1}, k-m-1) + p(N_{k-m-1}, k-m-2)$. If $a_{k-m} = 1$, then $\mathrm{val}(a_{k-m-1}\cdots a_1) = N_{k-m}$, and the number of corresponding representations is $p(N_{k-m}, k-m-1)$. Finally, $a_{k-m} = 0$ is not possible because it would imply

$$2N_{k-m-1} \geq \mathrm{val}(a_{k-m-1}\cdots a_1) = N_{k-m} + T_{k-m} = N_{k-m-1} + 2T_{k-m}$$

which is in contradiction with (4). Here we use that $k-m-1 \geq 1$ from the assumption. Putting everything together, we obtain

$$\begin{aligned} p(N_k, k-1) &= p(N_{k-m-1}, k-m-1) + p(N_{k-m-1}, k-m-2) + p(N_{k-m}, k-m-1) \\ &= p(N_{k-m-1}) - p(N_{k-m-1}, k-m) + p(N_{k-m}, k-m-1). \end{aligned} \tag{8}$$

Now suppose that $k \in \{m+2, \ldots, 2m\}$. Then $1 \leq k-m-1 \leq m-1$ and from the initial conditions in Table 1 we have $p(N_{k-m-1}, k-m) = 0$ and $p(N_{k-m}, k-m-1) = 0$. It follows from (8) that $p(N_k, k-1) = p(N_{k-m-1})$. With (6) and (7), it implies that

$$p(N_k) = p(N_{k-1}) + p(N_{k-m}) + p(N_{k-m-1}) \text{ for } k \in \{m+2, \ldots, 2m\}. \tag{9}$$

Now assume that $k > 2m$. Using (6) and (7) we deduce from (8) that

$$\begin{aligned} p(N_k, k-1) &= p(N_{k-m-1}) - p(N_{k-m-1}, k-m) + \\ &\quad + p(N_{k-m}) - p(N_{k-m}, k-m+1) - p(N_{k-m}, k-m) = \\ &= p(N_{k-m-1}) - p(N_{k-2m-1}) + p(N_{k-m}) - p(N_{k-2m}) - p(N_{k-m-1}) \\ &= p(N_{k-m}) - p(N_{k-2m}) - p(N_{k-2m-1}). \end{aligned}$$

Therefore, for $k > 2m$, we have

$$p(N_k) = p(N_{k-1}) + p(N_{k-m}) + p(N_{k-m}) - p(N_{k-2m}) - p(N_{k-2m-1}).$$

To conclude the argument, an easy induction on k shows the claim

$$p(N_k) = p(N_{k-1}) + p(N_{k-m}) + p(N_{k-m-1}) \text{ for } k > m+1,$$

since the base cases are treated in (9). □

4 Representations of zero in the signed m-bonacci system

We are interested in the representations of 0 over the balanced alphabet $B = \{-1, 0, 1\}$. We let $\mathcal{Z}$ denote the set of such representations. For a fixed $k \geq 1$,



Figure 1: Automaton accepting representations of 0 over the alphabet $\{-1, 0, 1\}$ in the Fibonacci numeration system.

we further let $\mathcal{Z}_k$ denote the set of representations of 0 of length at most k. Let us describe the number of such representations, i.e., $r(k) = \#\mathcal{Z}_k$. In order to determine $r(k)$ for $k \geq 2m+1$, let us present the correspondence between the m-bonacci numeration systems with the alphabet of digits $A = \{0, 1, 2\}$ and $B = \{-1, 0, 1\}$ and then derive the recurrence for the sequence $(r(k))_{k\geq 1}$.

Proposition 6. *Let $A = \{0, 1, 2\}$ and $B = \{-1, 0, 1\}$, $k, m \in \mathbb{N}$ with $m \geq 2$ and $k \geq 1$. The number $r(k)$ of representations of 0 of length at most k in the m-bonacci numeration system over the alphabet B is given by*

$$r(k) = \sum_{j=0}^{k} p_B(0, j) = p_A(N_k) - p_A(N_{k-m}) = p_A(N_{k-1}) + p_A(N_{k-m-1}) \text{ for } k \geq m+1$$

and $r(k) = p_A(N_k) = 1$ for $k \in \{0, \ldots, m\}$.

Proof. Any representation $(b_k \cdots b_1)$ of 0 over $B = \{-1, 0, 1\}$ (with possible leading zeros) is in one-to-one correspondence with a representation $(a_k \cdots a_1)$ of N_k of length at most k over the alphabet $A = \{0, 1, 2\}$, setting $a_j = b_j + 1$ for $j = 1, \ldots, k$. By (5), we have $r(k) = p_A(N_k, k) + p_A(N_k, k-1) = p_A(N_k) - p_A(N_k, k+1)$. Using (6), we have $r(k) = p_A(N_k) - p_A(N_{k-m})$. The last equality in the statement of the proposition follows from the recurrence on $p_A(N_k)$. □

Corollary 7. *The number $r(k)$ of representations of 0 of length at most k in the m-bonacci numeration system over the alphabet $B = \{-1, 0, 1\}$ satisfies*

$$r(k) = r(k-1) + r(k-m) + r(k-m-1) \text{ for } k \geq m+1 \tag{10}$$

and $r(k) = 1$ for $k \in \{0, \ldots, m\}$.

Remark 8. For further use, let us determine a set of initial values of the sequence $(r(k))_{k\geq 0}$. Obviously, the only representation of zero of length $k \leq m$ is (0^k), which is in accordance with the initial conditions of the recurrence (10). If $k = m+1$, we have, besides the representation (0^{m+1}) also $(1\bar{1}^m)$ and $(\bar{1}1^m)$, where we use $\bar{1}$ for -1 for lucidity of notation. We have $r(m+1) = 3 = r(m) + r(1) + r(0)$. For $m+2 \leq k \leq 2m$ we have from (10) that $r(k) = r(k-1) + 2$, i.e., $r(m+j) = 2j+1$ for $j = 1, \ldots, m$.

An automaton accepting the representations of zero in the signed Fibonacci numeration system ($m = 2$, digits $-1, 0, 1$) was described by Frougny [18, Chap. 7];



Figure 2: Automata accepting representations of 0 over the alphabet $\{-1, 0, 1\}$ in the m-bonacci numeration system with $m \geq 2$.

see Figure 1. The number of representations of 0 of length at most k is exactly the number of representations with k symbols where we allow leading zeros. Hence it corresponds to the number of length-k accepting paths in the automaton.

Here, we present an automaton recognizing representations of zero in the signed m-bonacci numeration system for $m \geq 2$.

Proposition 9. *Let $\mathcal{A} = (Q, B, \delta, I, T)$ be a deterministic finite automaton with the set of states $Q = \{i, u_1, \ldots, u_m, v_1, \ldots, v_m\}$, alphabet $B = \{-1, 0, 1\}$, the set of initial and terminal states $I = T = \{i\}$. Define the transition function $\delta : Q \times \{-1, 0, 1\} \to Q$,*

$$\begin{aligned} (i, 0) &\mapsto i, \quad (i, -1) \mapsto u_1, \quad (i, 1) \mapsto v_1 \\ (u_j, 1) &\mapsto u_{j+1}, \quad (u_m, 0) \mapsto u_1, \quad (u_m, 1) \mapsto i \\ (v_j, -1) &\mapsto v_{j+1}, \quad (v_m, 0) \mapsto v_1, \quad (v_m, -1) \mapsto i, \end{aligned}$$

see Figure 2. Then $\mathcal{A}$ recognizes precisely the representations of 0 in the m-bonacci numeration system over the alphabet B.

In order to justify that the accepting paths correspond precisely to the representations of 0, we study the adjacency matrix $\mathbb{A}$ of the automaton $\mathcal{A}$. In an adjacency matrix of the automaton, the element $\mathbb{A}_{j,k}$ is defined as 1 if there is an edge from the state j to the state k, and 0 otherwise. Fixing the ordering of states of $\mathcal{A}$ from 1 to $2m + 1$ as $\{i, u_1, \ldots, u_m, v_1, \ldots, v_m\}$, we have

$$\mathbb{A} = \begin{pmatrix} 1 & v^T & v^T \\ u & C & O \\ u & O & C \end{pmatrix},$$

where $v^T = (10 \ldots 0) \in \mathbb{Z}^m$, $u^T = (0 \ldots 01) \in \mathbb{Z}^m$, $O \in \mathbb{Z}^{m \times m}$ is the zero matrix and

$$C = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & & \vdots \\ \vdots & & \ddots & & 0 \\ 0 & & & 0 & 1 \\ 1 & 0 & \cdots & 0 & 0 \end{pmatrix} \in \mathbb{Z}^{m \times m}$$

is the companion matrix of the polynomial $x^m - 1$.

Lemma 10. *The characteristic polynomial of* $\mathbb{A}$ *is equal to*

$$f(x) = (x^m - 1)(x^{m+1} - x^m - x - 1) = x^{2m+1} - x^{2m} - 2x^{m+1} + x + 1.$$

Proof. For the computation of the characteristic polynomial $f(x) = -\det(\mathbb{A} - xI)$, we use the following statement about determinants of block matrices [21].

Let E, F, G, H *be complex matrices of dimensions* $p \times p$, $p \times q$, $q \times p$, $q \times q$, *with* H *invertible. Then for the determinant of* $\begin{pmatrix} E & F \\ G & H \end{pmatrix}$ *we have*

$$\det \begin{pmatrix} E & F \\ G & H \end{pmatrix} = \det(E - FH^{-1}G)\det(H)\,. \tag{11}$$

We use the statement (11) in two steps. First set

$$E = \begin{pmatrix} 1-x & v^T \\ u & H \end{pmatrix},\ F = \begin{pmatrix} v^T \\ O \end{pmatrix},\ G = \begin{pmatrix} u & O \end{pmatrix},$$

$$H = C - xI = \begin{pmatrix} -x & 1 & 0 & \cdots & 0 \\ 0 & -x & 1 & & \vdots \\ \vdots & & \ddots & & 0 \\ 0 & & & -x & 1 \\ 1 & 0 & \cdots & 0 & -x \end{pmatrix}.$$

Note that $\det(H) = (-1)^m(x^m - 1)$. Also note that the product $FH^{-1}G = (h_{j,k}) \in \mathbb{Z}^{(m+1)\times(m+1)}$ is a matrix that is null everywhere except for $c_{1,1} = h_{1,m}$ with $(h_{j,k}) = H^{-1} = \det(H)^{-1}\mathrm{adj}(H)$ where $\mathrm{adj}(H)$ is the adjugate matrix. Since the adjugate matrix is computed as the transpose of the cofactor matrix, we have

$$c_{1,1} = h_{1,m} = \frac{1}{(-1)^m(x^m-1)}(-1)^{m+1}\det\begin{pmatrix} 1 & 0 & \dots & 0 \\ -x & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & -x & 1 \end{pmatrix} = \frac{-1}{x^m - 1}\,.$$

At this stage, we have

$$\det(\mathbb{A} - xI) = \det(E - FH^{-1}G)\det(H) = \det(H)\det\begin{pmatrix} 1 - x - c_{1,1} & v^T \\ u & H \end{pmatrix}.$$

We use (11) again, with redefined blocks E, F, G, $E = (1 - x - c_{1,1})$, namely $F = v^T$, $G = u$. Realizing that $FH^{-1}G = (c_{1,1})$, we derive

$$\begin{aligned} -f(x) = \det(\mathbb{A} - xI) &= \det(H)^2(1 - x - 2c_{1,1}) = \\ &= (x^m - 1)^2\big(1 - x - \frac{-2}{x^m - 1}\big) = -(x^m - 1)(x^{m+1} - x^m - x - 1), \end{aligned}$$

which proves the lemma. □

Let us now demonstrate that the automaton presented in Figure 2 recognizes precisely the representations of 0 in the m-bonacci system over the alphabet $B = \{-1, 0, 1\}$.

Proof of Proposition 9. The language accepted by the automaton $\mathcal{A}$ depicted in Figure 2 can be described by the regular expression

$$0^*\big(1(\bar{1}^{m-1}0)^*\bar{1}^m + \bar{1}(1^{m-1}0)^*1^m\big)^*0^*$$

It can be directly checked with the use of the m-bonacci recurrence that every word of the above form corresponds to a representation of 0 in the sense of Definition 2. For every fixed k we thus have $\mathcal{L}_k(\mathcal{A}) \subseteq \mathcal{Z}_k$. In order to check the opposite inclusion, it suffices to show that

$$\#\mathcal{L}_k(\mathcal{A}) = \#\mathcal{Z}_k = r(k).$$

Consider the adjacency matrix $\mathbb{A}$ of the automaton $\mathcal{A}$. An accepting path in the automaton is a path both starting and ending in the initial state i. The number of such paths of length k, say $s(k)$, is equal to $(\mathbb{A}^k)_{11}$. We thus have $s(k) := \#\mathcal{L}_k(\mathcal{A}) = (\mathbb{A}^k)_{11}$. We will show that $s(k) = r(k)$ by verifying that both sequences satisfy the same linear recurrence of order $2m+1$ with the same initial conditions $s(j) = r(j)$ for $j = 0, \ldots, 2m$.

The characteristic polynomial f of the matrix $\mathbb{A}$ is computed in Lemma 10. By the Cayley-Hamilton theorem, we have

$$f(\mathbb{A}) = \mathbb{A}^{2m+1} - \mathbb{A}^{2m} - 2\mathbb{A}^{m+1} + \mathbb{A} + I = O,$$

where $I, O \in \mathbb{Z}^{(2m+1)\times(2m+1)}$ are the identity and zero matrix, respectively. Multiplying the above by $\mathbb{A}^k$ for $k \geq 0$, and considering the sequence $s(k) = (\mathbb{A}^k)_{11}$, we have

$$s(k+2m+1) - s(k+2m) - 2s(k+m+1) + s(k+1) + s(k) = 0 \quad \text{for } k \geq 0.$$

Let us verify that the sequence $r(k)$ satisfies the same recurrence. From Corollary 7 we have for $k \geq 0$ that

$$\begin{aligned} r(k+m+1) - r(k+m) - r(k+1) - r(k) &= 0, \\ r(k+2m+1) - r(k+2m) - r(k+m+1) - r(k+m) &= 0. \end{aligned}$$

Subtracting the two, we obtain

$$r(k+2m+1) - r(k+2m) - 2r(k+m+1) + r(k+1) + r(k) = 0 \quad \text{for } k \geq 0,$$

as desired. The initial values $r(0), \ldots, r(2m)$ are determined in Remark 8 as $r(j) = 1$ for $0 \leq j \leq m$ and $r(j+m) = 2j+1$ for $0 \leq j \leq m$. It remains to determine $s(0), \ldots, s(2m)$. For that, we check directly the automaton $\mathcal{A}$. We see that $s(k) = r(k)$ for $0 \leq k \leq 2m$ which implies that $s(k) = r(k)$ for every $k \geq 0$. □



Figure 3: Zeros of the polynomial $\chi(x) = x^{m+1} - x^m - x - 1$ for $m = 3$, $m = 7$ and $m = 32$. The zeros λ with $\Re\lambda > 0$ are of modulus greater than 1, the zeros with negative real part lie inside the unit disk.

5 Characteristic polynomial

When studying growth of the sequences $p_A(N_k)$, $r(k)$ counting representations in the m-bonacci system, one is interested in the zeros of the characteristic polynomial of the corresponding recurrence relation. From Proposition 4 and Corollary 7 we see that both $p_A(N_k)$ and $r(k)$ satisfy the same recurrence with characteristic polynomial $f(x) = x^{m+1} - x^m - x - 1$. Quadrinomials whose coefficients take values only in $\{\pm 1\}$ have been considered by several authors, e.g., Ljunggren [17], Finch and Jones [7], or Mills [19]. It is not difficult to derive the following result.

Proposition 11. *Let $m \geq 2$. Then $\chi(x) = x^{m+1} - x^m - x - 1$ is irreducible over $\mathbb{Q}$ if and only if $m \not\equiv 3 \pmod 4$. If $m = 4k + 3$ for $k \geq 1$, then we have $\chi(x) = (x^2 + 1)\psi(x)$ where for $k = 1$ we have $\psi(x) = (x^3 - x^2 + 1)(x^3 - x - 1)$, and for $k \geq 2$, $\psi \in \mathbb{Z}[x]$ is irreducible over $\mathbb{Q}$.*

Proof. It follows from Theorem 2 in the paper by Finch and Jones [7] that the quadrinomial $\chi(x) = x^{m+1} - x^m - x - 1$ is irreducible over $\mathbb{Q}$ if and only if $m \not\equiv 3 \pmod 4$. The factorization of the polynomial χ in the reducible case can be deduced from the results of Ljunggren [17] and Mills [19]. □

6 Acknowledgments

This work was supported by the project CZ.02.1.01/0.0/0.0/16_019/0000778 from European Regional Development Fund.

References

[1] A. Avižienis, Signed-digit number representations for fast parallel arithmetic, *IRE Trans.* **EC-10** (1961), 389–400.

[2] J. Berstel, An exercise on Fibonacci representations, *Theor. Inform. Appl.* **35** (2001), 491–498.

[3] C. Y. Chow and J. E. Robertson, Logical design of a redundant binary adder, In *1978 IEEE 4th Symposium on Computer Arithmetic (ARITH)*, pp. 109–115. Institute of Electrical and Electronics Engineers, 1978.

[4] M. Edson, Calculating the numbers of representations and the Garsia entropy in linear numeration systems, *Monatsh. Math.* **169** (2013), 161–185.

[5] M. Edson and L. Q. Zamboni, On representations of positive integers in the Fibonacci base, *Theoret. Comput. Sci.* **326** (2004), 241–260.

[6] M. Edson and L. Q. Zamboni, On the number of partitions of an integer in the m-bonacci base, *Ann. Inst. Fourier (Grenoble)* **56** (2006), 2271–2283.

[7] C. Finch and L. Jones, On the irreducibility of $\{-1, 0, 1\}$-quadrinomials, *Integers* **6** (2006), Paper #A16.

[8] C. Frougny, Systèmes de numération linéaires et θ-représentations, *Theoret. Comput. Sci.* **94** (1992), 223–236.

[9] C. Frougny, E. Pelantová, and M. Svobodová, Parallel addition in non-standard numeration systems, *Theoret. Comput. Sci.* **412** (2011), 5714–5727.

[10] C. Frougny and W. Steiner, Minimal weight expansions in Pisot bases, *J. Math. Cryptol.* **2** (2008), 365–392.

[11] P. J. Grabner and C. Heuberger, On the number of optimal base 2 representations of integers, *Des. Codes Cryptogr.* **40** (2006), 25–39.

[12] P. J. Grabner, P. Kirschenhofer, and R. F. Tichy, Combinatorial and arithmetical properties of linear numeration systems, *Combinatorica* **22** (2002), 245–267.

[13] C. Heuberger, Minimal expansions in redundant number systems and shortest paths in graphs, *Computing* **63** (1999), 341–349.

[14] P. Kocábová, Z. Masáková, and E. Pelantová, Integers with a maximal number of Fibonacci representations, *Theor. Inform. Appl.* **39** (2005), 343–359.

[15] P. Kocábová, Z. Masáková, and E. Pelantová, Ambiguity in the m-bonacci numeration system, *Discrete Math. Theor. Comput. Sci.* **9** (2007), 109–123.

[16] C. G. Lekkerkerker, Representation of natural numbers as a sum of Fibonacci numbers, *Simon Stevin* **29** (1952), 190–195.

[17] W. Ljunggren, On the irreducibility of certain trinomials and quadrinomials, *Math. Scand.* **8** (1960), 65–70.

[18] M. Lothaire, *Algebraic Combinatorics on Words*, Vol. 90 of *Encyclopedia of Mathematics and its Applications*, Cambridge University Press, 2002.

[19] W. H. Mills, The factorization of certain quadrinomials, *Math. Scand.* **57** (1985), 44–50.

[20] A. Pethő and R. F. Tichy, On digit expansions with respect to linear recurrences, *J. Number Theory* **33** (1989), 243–256.

[21] J. Silvester, Determinants of block matrices, *Math. Gaz.* **84** (2000), 460–467.

[22] P. K. Stockmeyer, A smooth tight upper bound for the Fibonacci representation function $R(N)$, *Fibonacci Quart.* **46/47** (2008/09), 103–106.

[23] O. Vlašic, *Redundance in numeration systems*, Bachelor's project, Czech Technical University in Prague, 2016. In Czech.

2020 *Mathematics Subject Classification*: Primary 11A67; Secondary 11B39.
Keywords: linear numeration system, m-bonacci number.

Received May 6 2021; revised version received June 2 2021; June 4 2021. Published in *Journal of Integer Sequences*, June 4 2021.

Return to Journal of Integer Sequences home page.