



# Digraphs With Exactly One Eulerian Tour

Luz Grisales  
Massachusetts Institute of Technology  
Cambridge, MA 02139  
USA  
[luzg@mit.edu](mailto:luzg@mit.edu)

Antoine Labelle  
McGill University  
Montréal, QC H3A 0G4  
Canada  
[antoine.labelle@mail.mcgill.ca](mailto:antoine.labelle@mail.mcgill.ca)

Rodrigo Posada  
Massachusetts Institute of Technology  
Cambridge, MA 02139  
USA  
[rposada@mit.edu](mailto:rposada@mit.edu)

Stoyan Dimitrov  
University of Illinois at Chicago  
Chicago, IL, 60607  
USA  
[sdimit6@uic.edu](mailto:sdimit6@uic.edu)

## Abstract

We give two combinatorial proofs of the fact that the number of loopless directed graphs (digraphs) with  $n$  non-isolated vertices and with exactly one Eulerian tour up to a cyclic shift is  $\frac{1}{2}(n-1)!C_n$ , where  $C_n$  denotes the  $n$ -th Catalan number. We construct a bijection with a set of labeled rooted plane trees and with a set of valid parenthesis arrangements.

# 1 Introduction and main facts

Richard Stanley has a list containing nearly 250 problems and facts, for which he asks of combinatorial proofs [1]. This work describes two such proofs for one of the problems in the list, namely Problem 199, without a known combinatorial proof. First, we recall some definitions following [2].

A (finite) directed graph (or digraph)  $D$  consists of a vertex set  $V = \{v_1, \dots, v_n\}$  and an edge set  $E = \{e_1, \dots, e_q\}$ , together with a function  $\phi : V \rightarrow V$  determining the direction of each edge. If  $\phi(e) = (u, v)$ , then we think of  $e$  as an arrow from  $u$  to  $v$ . We will call  $u$  initial vertex and  $v$  final vertex. The outdegree of a vertex  $v$ , denoted  $\text{outdeg}(v)$ , is the number of edges of  $D$  with initial vertex  $v$ . Similarly, the indegree of  $v$ , denoted  $\text{indeg}(v)$ , is the number of edges of  $D$  with final vertex  $v$ . A loop is an edge  $e$  for which  $\phi(e) = (v, v)$  for some vertex  $v$ . A digraph is balanced if  $\text{indeg}(v) = \text{outdeg}(v)$  for each of its vertices  $v$ . Let  $[n] := \{1, 2, \dots, n\}$ . An oriented path in a digraph  $D$  is a sequence of vertices  $v_1, \dots, v_m$ , where  $(v_i, v_{i+1})$  is an edge of  $D$  for each  $i \in [m-1]$ . If the vertices  $v_1, \dots, v_m$  are all different, then we call the path *simple*. If we have a simple path  $v_1, \dots, v_{m-1}$ , then we call the path  $v_1, \dots, v_{m-1}, v_1$  an *oriented simple cycle*.

**Definition 1.** An *Eulerian tour* in a directed graph  $D$  is a sequence of vertices  $a_1 a_2 \cdots a_k$  such that  $(a_1, a_2), (a_2, a_3), \dots, (a_{k-1}, a_k), (a_k, a_1)$  are all the distinct directed edges of  $D$ .

Any cyclic shift  $a_i a_{i+1} \cdots a_k a_1 \cdots a_{i-1}$  of an Eulerian tour is also an Eulerian tour and we will say that these tours are *equivalent up to a cyclic shift*.

**Definition 2.** A *uni-Eulerian digraph* is a digraph which has no isolated vertices and contains exactly one Eulerian tour (and its equivalents under cyclic shifts).

Our goal is to prove the following claim.

**Theorem 3.** If  $A_n$  is the set of loopless uni-Eulerian digraphs on the vertex set  $[n]$ , then  $|A_n| = \frac{1}{2}(n-1)!C_n$  (sequence [A102693](#) in the OEIS [5]), where  $C_n = \frac{1}{n+1}\binom{2n}{n}$  denotes the  $n$ -th Catalan number.

For example,  $|A_3| = 5$ . Indeed, there are two such digraphs that look like triangles and three that consist of two 2-cycles with a common vertex (see Figure 1). The related OEIS sequence, [A102693](#), was created by Richard Stanley. As a reference, he points out to an unpublished work of him. Thus, we can assume that Theorem 3 was first proved there.

It is not difficult to show that every uni-Eulerian graph must be connected and balanced [2, Theorem 10.1]. The BEST theorem that we recall below gives us a formula for the total number of Eulerian tours in a digraph. In order to understand this result, one should be familiar with the term *oriented tree* (see Figure 2). An oriented tree with root  $v$  is a finite digraph  $T$  with  $v$  as one of its vertices, such that there is a unique directed path from any other vertex of  $T$  to  $v$ . This means that the underlying undirected graph (after we erase all the arrows of the edges of  $T$ ) is a tree.




Figure 1: The five digraphs with three vertices and a unique Eulerian tour.




Figure 2: Example of an oriented tree.

**Theorem 4** ([2], Theorem 10.2). *Let  $D$  be a connected balanced digraph with vertex set  $V$ . Fix an edge  $e$  in  $D$  and let  $v$  be the initial vertex of that edge. Let  $\tau(D, v)$  denote the number of oriented (spanning) subtrees of  $D$  with root  $v$ , and let  $\epsilon(D, e)$  denote the number of Eulerian tours of  $D$  starting with the edge  $e$ . Then*

$$\epsilon(D, e) = \tau(D, v) \prod_{u \in V} (\text{outdeg}(u) - 1)!$$

**Corollary 5** (from Theorem 4). *A digraph  $D \in A_n$  if and only if*

1. *For every vertex  $v$ ,  $D$  has exactly one oriented (spanning) subtree with root  $v$ .*
2. *The outdegree of an arbitrary vertex of  $D$  is 1 or 2.*

Note that if the two conditions of Corollary 5 hold, then  $\epsilon(D, e) = 1$  for every edge  $e$  of the digraph  $D$  and thus  $D$  has a unique Eulerian tour. Using Corollary 5, we will characterize the digraphs in  $A_n$  by two other conditions that will be used later.

**Lemma 6.** *A digraph  $D \in A_n$  if and only if*

- i) *There exists a unique oriented simple path between any two vertices of  $D$ .*
- ii) *Every vertex of  $D$  is part of exactly one or two simple oriented cycles.*

**Corollary 7.** *If  $D \in A_n$ , then no pair of cycles in  $D$  have an edge in common.*

*Proof.* Assume  $C_1$  and  $C_2$  are two different cycles in  $D$ , which have the edge  $(u, v)$  in common. Then, we will have at least two oriented paths from  $v$  to  $u$  in  $D$ —one following  $C_1$  and another one following  $C_2$ . This is a contradiction with Corollary 5.  $\square$

*Proof of Lemma 6.* [first part:  $D \in A_n \implies$  conditions  $i$ ) and  $ii$ ] Let  $D \in A_n$ . Let  $u$  and  $v$  be two arbitrary vertices in  $D$ . By condition (1) of Corollary 5, there exists exactly one oriented spanning subtree  $T_u$  of  $D$  with root  $u$ . We know that  $v$  has to be a vertex of  $T_u$  and that there exists a unique simple path  $\mathcal{P}$  from  $v$  to  $u$  in  $T_u$ , which is a simple oriented path in  $D$ . Assume that there exists another path  $\mathcal{P}' \neq \mathcal{P}$  between  $v$  and  $u$  in  $D$ . Begin from  $v$  and follow  $\mathcal{P}'$ . Let  $f'$  be the first edge in  $\mathcal{P}'$ , which is not part of  $\mathcal{P}$  and let  $f$  be the edge in  $\mathcal{P}$  with the same initial vertex as  $f'$ . If you delete  $f$  from  $T_u$  and add  $f'$  to it, you will obtain a graph  $T'_u$ . One can easily see that  $T'_u$  is an oriented spanning subtree of  $D$ , different from  $T_u$  (see Figure 3). This is a contradiction. Thus, we showed that condition  $i$ ) holds.

By Corollary 7, we know that no pair of cycles in  $D$  have an edge in common. Therefore, Corollary 5 implies that each vertex of  $D$  can be part of at most two simple oriented cycles since its outdegree is 1 or 2. It remains to show that each vertex of  $D$  is part of at least one such cycle. Let  $v$  be an arbitrary vertex of  $D$  and let  $(u, v)$  be an edge of  $D$  (such an edge exists since the indegree of  $v$  is 1 or 2). We showed that there is a unique oriented simple path between  $v$  and  $u$ . This path together with the edge  $(u, v)$  forms a simple oriented cycle. Thus condition  $ii$ ) holds.




Figure 3: The tree  $T_u$  in the first part of the proof of Lemma 6;  $u = 1$ ,  $v = 11$ ,  $\mathcal{P} = 11, 9, 6, 5, 1$ ,  $\mathcal{P}' = 11, 9, 6, 2, 1$ ,  $f' = (6, 2)$  and  $f = (6, 5)$ . Delete  $f$  and add  $f'$  to obtain another tree  $T'_u$ .

[second part: conditions  $i$ ) and  $ii$ )  $\implies D \in A_n$ ] Take a digraph  $D$  for which conditions  $i$ ) and  $ii$ ) hold. We have to show that conditions (1) and (2) from Corollary 5 also hold. Let  $v$  be an arbitrary vertex of  $D$ . Condition  $ii$ ) implies that  $\text{outdeg}(v) \geq 1$ . We will show that  $\text{outdeg}(v) < 3$ . By Corollary 7 no edge of  $D$  can be part of two different simple cycles. Assume that  $\text{outdeg}(v) \geq 3$  and let  $(v, u_1)$ ,  $(v, u_2)$  and  $(v, u_3)$  are three different edges of  $D$ .

We know that there exist simple paths between  $u_i$  and  $v$ , for  $i = 1, 2, 3$ . Thus,  $v$  participates in simple cycles through  $u_i$ , for  $i = 1, 2, 3$  and no two of these simple cycles share an edge. Therefore these three cycles are different. This is a contradiction with condition *ii*).

It remains to show that condition (1) from Corollary 5 holds. Take an arbitrary vertex  $v$  of  $D$ . We have a unique oriented simple path from each of the other vertices of  $D$  to  $v$ . Take the union of these paths. The graph that you will obtain is an oriented spanning subtree  $T_v$  of  $D$  with root  $v$ . Assume that there exists another such subtree  $T'_v$ . Then, since  $T_v \neq T'_v$ , we must have a vertex  $w$  in  $D$ , for which the unique oriented path from  $w$  to  $v$  in  $T'_v$  is different from the unique oriented path from  $w$  to  $v$  in  $T_v$ . These are two different oriented paths from  $w$  to  $v$  in  $D$ , which is a contradiction.  $\square$

## 2 Bijection with a set of labeled rooted plane trees

We will construct a bijection between the digraphs in  $A_n$  and a set of labeled rooted plane trees on  $n + 1$  vertices. Let  $U_n$  be the set of the unlabeled rooted plane trees with  $n + 1$  vertices. It is well-known that  $|U_n| = C_n$  (see [3, Theorem 1.5.1]). Let  $L_n$  be the set of the labeled rooted plane trees with  $n + 1$  vertices such that the root is always labeled as 0 and the left-most child of the root is always labeled as 1. We have  $|L_n| = (n - 1)!|U_n| = (n - 1)!C_n$ . Finally, let  $L'_n \subset L_n$  be the set of the labeled trees in  $L_n$ , such that the vertex 2 is in the subtree with root 1.

First, we define a map  $f$  over  $L_n$ , such that  $f(L'_n) = L_n \setminus L'_n$  and  $f(f(T)) = T$ , i.e., an involution. This shows that  $|L'_n| = \frac{|L_n|}{2} = |A_n|$ . Then, we define a map  $g : L'_n \rightarrow A_n$ , which is shown to be a bijection. Below, we will denote by  $T_{(x,j)}$  the  $j$ -th child (from left to right) of the vertex  $x$  of a tree  $T \in L_n$ .

**Definition 8.** Let  $f : L_n \rightarrow L_n$  be a map, which switches the places of the subtree with root 1 (excluding 1) and the subtree with root 0 (excluding 0 and the subtree with root 1), for every tree in  $L_n$  (see Figure 4). Formally, if  $T \in L_n$ , then  $f(T)$  has the following properties. For every  $j \geq 1$ :

- If the vertex  $v$  is the  $(j + 1)$ -th child of the vertex 0 in  $T$ , then  $v$  is the  $j$ -th child of vertex 1 in  $f(T)$ , i.e.,  $T_{(0,j+1)} = f(T)_{(1,j)}$ .
- If the vertex  $v$  is the  $j$ -th child of the vertex 1 in  $T$ , then  $v$  is the  $(j + 1)$ -th child of vertex 0 in  $f(T)$ , i.e.,  $T_{(1,j)} = f(T)_{(0,j+1)}$ .
- All the other directed edges are left the same for both trees, i.e.,  $T_{(u,j)} = f(T)_{(u,j)}$  for  $u \notin \{0, 1\}$ .

It is easy to check that  $f(L'_n) = L_n \setminus L'_n$  and  $f(f(T)) = T$ , i.e.,  $f^{-1} = f$ .

**Definition 9.** Let  $g : L'_n \rightarrow A_n$  be a map, such that if  $T \in L'_n$ ,  $g(T) = D'$  is a digraph with  $V(D') = [n]$  and  $E(D')$ , such that if  $x$  is a vertex of  $T$  with  $r$  children, then:




Figure 4: Example of the action of the map  $f$ .

1. For every  $i \in [1, r)$ ,  $(T_{(x,i)}, T_{(x,i+1)}) \in E(D')$ .
2. If  $x = 0$ , then  $(T_{(x,r)}, T_{(x,1)}) \in E(D')$ .
3. If  $x \neq 0$ , then  $(T_{(x,r)}, x) \in E(D')$  and  $(x, T_{(x,1)}) \in E(D')$ .




Figure 5: Example of the action of the map  $g$ .

**Lemma 10.** For every  $T \in L'_n$ ,  $g(T) \in A_n$ , i.e.,  $g(T)$  has exactly one Eulerian tour.

*Proof.* First, note that every vertex  $x$  of  $g(T)$  belongs to one or two cycles:

- The cycle where  $x$  and its parent from  $T$  both belong.
- In case  $x$  has children in  $T$ , the cycle formed by  $x$  and its children in  $g(T)$ .

By Lemma 6, it remains to show that there exists a unique oriented simple path between any two vertices of  $g(T)$ , for arbitrary  $T \in L'_n$ . To see this, observe that if  $(v, w) \in E(T)$  and  $v, w \neq 0$  or if both  $v$  and  $w$  are children of the root 0 in  $T$ , then we have a unique oriented simple path between  $v$  and  $w$  in  $g(T)$ , which is part of a single cycle. For instance, the edge  $(1, 3)$  in the graph  $T$  shown at Figure 5 corresponds to the oriented simple path

1273 in  $g(T)$ , whereas the path 514 in  $g(T)$  corresponds to the pair of children 4 and 5 of the root of  $T$ .

We will show that since we have a unique non-oriented path  $\mathcal{P}$  between any two vertices  $u$  and  $v$  in  $T$ , where  $u, v \neq 0$ , we will also have a unique oriented simple path  $\mathcal{P}^{\text{or}}$  between  $u$  and  $v$  in  $g(T)$ . If the vertex 0 is part of  $\mathcal{P}$ , then we must have vertices  $h_1$  and  $h_2$  in  $T$ , such that the edges  $(h_1, 0)$  and  $(0, h_2)$  are part of  $\mathcal{P}$  (since  $v, w \neq 0$ ). Hence,  $h_1$  and  $h_2$  are two children of the root 0. Replace the edges  $(h_1, 0)$  and  $(0, h_2)$  of  $\mathcal{P}$  with the unique oriented simple path between  $h_1$  and  $h_2$  in  $g(T)$ . Replace all the other edges of  $\mathcal{P}$  with the corresponding oriented simple paths to obtain  $\mathcal{P}^{\text{or}}$ . For example, the unique path  $\mathcal{P}$  between 3 and 8 in the graph  $T$  on Figure 5 is comprised of the edges  $(3, 1)$ ,  $(1, 7)$ ,  $(7, 8)$ . The oriented paths corresponding to these edges are 31, 127 and 798, respectively. The union of these paths, namely 312798, gives the unique path  $\mathcal{P}^{\text{or}}$  between 3 and 8 in  $g(T)$ . Another example is the path 71056 in  $T$ , which transforms to the path 731456 in  $g(T)$ .  $\square$

**Lemma 11.** *For every digraph  $D \in A_n$ , there exists a unique labeled tree  $T \in L'_n$ , for which  $g(T) = D$ . i.e.,  $g$  has an inverse.*

*Proof.* Let  $D$  be an arbitrary digraph in  $A_n$ . Below, we describe the procedure  $\text{buildSubtree}(r, C, T, D)$  that will be used to obtain the tree  $T$ , for which  $g(T) = D$ . The first argument,  $r$ , is a vertex in  $D$  and the second argument,  $C$ , is a cycle in  $D$  that contains  $r$ .

```

1  buildSubtree( $r, C, T, D$ ):
2      For each  $v \in C$ :
3          if  $v \neq r$ :
4              add an edge  $(r, v)$  to  $E(T)$ .
5              if  $v$  is part of a cycle  $C_1 \neq C$ :
6                  buildSubtree( $v, C_1, T, D$ ).
```

Initially, let  $T$  be an empty tree with  $n + 1$  vertices, i.e., let  $V(T) = \{0, 1, \dots, n\}$  and let  $E(T) = \emptyset$ . Take the vertex with label 1 in  $D$ . By Lemma 6, this vertex belongs to one or two simple oriented cycles. We consider each case separately.

Suppose that the vertex belongs to one such cycle and let  $C$  denotes this cycle. Then, run  $\text{buildSubtree}(1, C, T, D)$  and add an edge  $(0, 1)$  to  $E(T)$ . One can easily show that the resulting graph,  $T$ , is a tree in  $L'_n$ . First,  $T$  is connected since the execution of  $\text{buildSubtree}(1, C, T, D)$  will reach every vertex of  $D$  and connect this vertex to an already reached vertex. In addition, if we have two different paths between two vertices  $u$  and  $v$  of  $T$ , then we will be able to find two simple oriented paths between  $u$  and  $v$  in  $D$ , which contradicts Lemma 6. Finally,  $n \geq 2$  and the only vertex of  $T$ , which is not in the subtree with root 1, is the vertex 0. Thus, 2 is in that subtree and  $T \in L'_n$ .

Now, suppose that the vertex 1 belongs to two different cycles  $C_1$  and  $C_2$ . Then, find the unique oriented simple path between 1 and 2 in  $D$ . This path has to have an edge in common with either  $C_1$  or  $C_2$ , but not with both. Otherwise, we will have a contradiction with

Corollary 7. Without loss of generality, let this be  $C_2$ . Execute  $\text{buildSubtree}(0, C_1, T, D)$ . The graph  $T$ , obtained at the end, will be a tree in  $L'_n$  (see Figure 6).  $\square$




Figure 6: Example of the action of the inverse map  $g^{-1}$ .

Lemmas 10 and 11 imply that the map  $g$  is a bijection.

### 3 A bijection with parentheses arrangements

In this section, we give a second combinatorial proof of Theorem 3, via a bijection between the digraphs in  $A_n$  and a set of valid parentheses arrangements. Suppose that you have  $n$  pairs of opening and closing parenthesis, such that the two parenthesis in each pair are labeled with the numbers in  $[n]$ . A valid labeled parentheses arrangement is an ordering of these  $2n$  parenthesis, such that we cannot have two interlaced pairs, e.g.,  $(i(j)_i)_j$  for some  $i, j \in [n]$ . A valid unlabeled parentheses arrangement is a sequence of unlabeled opening and closing parentheses that can be obtained by forgetting the labels of a valid labeled arrangement. One can easily check that a sequence of  $n$  opening and  $n$  closing parentheses is valid if and only if every prefix of the sequence has at least as many opening parentheses as closing parentheses. The number of valid arrangements of  $n$  unlabeled pairs of parentheses is  $C_n$  [3, Theorem 1.5.1]. Thus the number of such arrangements for labeled pairs is  $n!C_n$ .

To construct a bijection with the set of digraphs  $A_n$ , let us first note that one can assume that all vertices of a uni-Eulerian digraph have indegree and outdegree 2, provided that we allow digraphs with loops.

**Lemma 12.** *If  $B_n$  is the set of all uni-Eulerian digraphs on the vertex set  $[n]$  (possibly with loops) with all vertices of indegree and outdegree 2, then  $|B_n| = |A_n|$*

*Proof.* Given a digraph  $D$  in  $A_n$ , Corollary 5 implies that all vertices have outdegree 1 or 2 and the same indegree. Moreover, the single Eulerian tour of  $D$  passes exactly once through each vertex of outdegree 1. Hence, adding a loop to every vertex of outdegree 1 gives an element of  $B_n$ .

Conversely, given a digraph  $D'$  in  $B_n$ , deleting all loops gives an element of  $A_n$ . Indeed, the loopless digraph still has a unique Eulerian tour, which is just the tour for  $D'$  without



the loops (the uniqueness follows because adding back loops must give an Eulerian tour for  $D'$ ). These two maps are inverses of each other and thus give a bijection between  $A_n$  and  $B_n$ .  $\square$

Now, let  $B_n^*$  be the set of digraphs in  $B_n$  together with an identified edge. Since all digraphs in  $B_n$  have  $2n$  edges, we have  $|B_n^*| = 2n|B_n| = 2n|A_n|$ . We will give a bijection between  $B_n^*$  and the set of valid arrangements of  $n$  labeled pairs of parentheses, which will show that  $2n|A_n| = n!C_n$ , that is,  $|A_n| = \frac{1}{2}(n-1)!C_n$ .

**Theorem 13.** *There exists a bijection between  $B_n^*$  and the set of valid arrangements of  $n$  labeled pairs of parentheses.*

*Proof.* [first part: Digraphs in  $B_n^* \rightarrow$  valid parentheses arrangements].

Let  $D$  be a digraph in  $B_n^*$  with identified edge  $e$ . We define a parentheses arrangement  $h(D)$  as follows:

Following the unique Eulerian tour of  $D$ , starting at  $e$ , open the  $i$ -th pair of parentheses when you pass through the vertex  $i$  for the first time and close the  $i$ -th pair of parentheses when you pass through the vertex  $i$  for the second time (see Figure 7 below).




Figure 7: The digraph in  $B_3^*$  (with the edge  $2 \rightarrow 1$  being identified) that yields the string  $(1)1(2(3)3)2$

To show that the resulting string of parentheses is valid, we have to show that we cannot have two interlaced pairs of parentheses, e.g.,

$$\cdots (i \cdots (j \cdots )_i \cdots )_j \cdots .$$

In other words, the unique tour cannot have the form

$$i \xrightarrow{a} j \xrightarrow{b} i \xrightarrow{c} j \xrightarrow{d} i$$

for some walks  $a, b, c, d$ . But this is clearly impossible, because otherwise we would have a second Eulerian tour  $i \xrightarrow{a} j \xrightarrow{d} i \xrightarrow{c} j \xrightarrow{b} i$ .

[second part: Valid parentheses arrangements  $\rightarrow$  digraphs in  $B_n^*$ ] Given a valid parentheses arrangement  $w = (x \cdots )_y$ , we obtain a digraph  $h^{-1}(w) \in B_n^*$  by putting an edge between the corresponding vertices of any pair of consecutive parentheses (from the first parenthesis to the second) and an edge from  $y$  to  $x$ . The identified edge is  $y \rightarrow x$ .

Clearly, every vertex in  $h^{-1}(w)$  has indegree and outdegree 2 and that there exists an Eulerian tour  $T$ , given by the order of the parentheses' labels in  $w$ . Hence, we just have to show that  $T$  is the unique Eulerian tour of  $h^{-1}(w)$ . Let  $i \in [n]$  and let

$$w = \cdots ?_{\ell} ( (j \cdots )_i ?_k \cdots ,$$

where  $?$  represents either a closing or an opening parenthesis (if  $( (i$  and  $)_i$  are consecutive we let  $j = i$ , if  $( (i$  is the first parenthesis of  $w$  we let  $\ell$  be the label of the last one and if  $)_i$  is the last parenthesis, we let  $k$  be the label of the first one). We have to show that if an Eulerian tour enters the vertex  $i$  for the first time from  $\ell$ , this tour must exit the vertex  $i$  towards  $j$  and not  $k$ . Indeed, if this is true for all  $i$ , then the Eulerian tour is entirely determined by its first edge. Thus, this tour and  $T$  are equal up to a cyclic shift. Suppose, for the sake of contradiction, that there exists an Eulerian tour  $T'$  of  $h^{-1}(w)$ , which exits  $i$  towards  $k$ , after entering  $i$  for the first time, through  $\ell$ .

Note that, by the properties of valid parentheses arrangements, the two parentheses corresponding to any vertex  $v \neq i$  are either both between  $( (i$  and  $)_i$  (then we will say that  $v$  is of type  $A$ ) or both outside (type  $B$ ). Clearly, all edges of the graph  $h^{-1}(w)$  with initial vertex of type  $A$  (respectively  $B$ ) have a final vertex either  $i$  or of type  $A$  (respectively  $B$ ), so the only way to go from a vertex of type  $A$  to a vertex of type  $B$  is through  $i$  and vice-versa. Therefore, since  $k$  is of type  $B$ , we must re-enter  $i$  in  $T'$  through a vertex of type  $B$ , in order to access vertices of type  $A$ . The only way to do so, however, is through the edge  $\ell \rightarrow i$ , which was already used in  $T'$ . This is a contradiction, so the uniqueness of the Eulerian tour is proved. The two described maps  $h$  and  $h^{-1}$  are obviously inverses of each other, so the proof is complete.  $\square$

## 4 Acknowledgments

We are thankful to Jan Kynčel for the suggestions in the mathoverflow post related to the considered problem [4].

## References

- [1] R. P. Stanley, "Bijective proof problems.", 2009, <https://klein.mit.edu/~rstan/bij.pdf>.
- [2] R. P. Stanley, *Algebraic Combinatorics*. Springer, 2013.
- [3] R. P. Stanley, *Catalan Numbers*. Cambridge University Press, 2015.
- [4] J. Kynčel, mathoverflow answer, 2020, <https://mathoverflow.net/questions/366383/digraphs-with-exactly-one-eulerian-tour>.

[5] OEIS Foundation Inc., The On-Line Encyclopedia of Integer Sequences, 2020, <https://oeis.org>.

---

2020 *Mathematics Subject Classification*: Primary 05A19; Secondary 05C20.

*Keywords*: bijective proof, directed graph, Eulerian tour, Catalan number, labeled tree.

---

(Concerned with sequence [A102693](#).)

---

Received May 26 2021; revised versions received June 4 2021; August 24 2021; February 10 2022. Published in *Journal of Integer Sequences*, March 27 2022.

---

Return to [Journal of Integer Sequences home page](#).