

Fixed Points and Cycles of the Kaprekar Transformation: 1. Odd Bases

Anthony Kay
72 Tiverton Road
Loughborough
Leicestershire LE11 2RY
United Kingdom
anthony.kay@sky.com

Katrina Downes-Ward

Abstract

We extend Yamagami and Matsui's theory of the Kaprekar transformation in base 3 to higher odd bases. The structure of primitive proper, primitive non-proper and general cycles appears in all odd bases, but many other features differ according to whether the base is congruent to 1 or 3 modulo 4. In the latter case, cycles are derived from subgroups and cosets in multiplicative groups modulo odd numbers. We examine cycles and fixed points in bases 5 and 7 in some detail, and make some broad observations relating to higher bases. There also exist many cycles outside the primitive/general structure.

1 Introduction

The *Kaprekar transformation* $T_{b,n}$ takes an integer x which has $n \geq 2$ digits when represented in base $b \geq 2$, rearranges the digits in descending and ascending orders to produce new integers x_{\downarrow} and x_{\uparrow} , and subtracts the latter from the former. Thus, given $a_0, a_1, \dots, a_{n-1} \in \mathbb{Z}$ with

$$0 \leq a_0 \leq a_1 \leq \dots \leq a_{n-1} < b,$$

we take an integer

$$x = \sum_{j=0}^{n-1} a_j b^{\alpha_j}$$

where the set $\{\alpha_0, \alpha_1, \dots, \alpha_{n-1}\}$ is some permutation of the set $\{0, 1, \dots, n-1\}$, and the transformation is then defined by

$$T_{b,n} : x \mapsto x_{\downarrow} - x_{\uparrow} := \sum_{j=0}^{n-1} a_j b^j - \sum_{j=0}^{n-1} a_{n-1-j} b^j. \quad (1)$$

Clearly the result of the transformation depends only on how many occurrences there are of each of the digits $0, 1, \dots, b-1$ in the set $\{a_0, a_1, \dots, a_{n-1}\}$. So if there are k_i occurrences of the digit i in the base- b expression of an integer x , i.e., in the set $\{a_0, a_1, \dots, a_{n-1}\}$, we define the *Kaprekar index* of x in base b to be the b -tuple $(k_0, k_1, \dots, k_{b-1})$, generalizing the definition introduced by Yamagami and Matsui [11] for base 3. We shall also use the shorthand notation \mathbf{k} for this b -tuple, and refer to k_i as the *i -component* of \mathbf{k} for each $i \in \{0, 1, \dots, b-1\}$; components of Kaprekar indices have previously been referred to as *digit counters* by Lapenta, Ludington, and Prichett [5, 8]. For example, the integer represented as 30133103 in base 4 has the Kaprekar index $(2, 2, 0, 4)$, with 0-component $k_0 = 2$, etc.; applying the Kaprekar transformation to this integer involves doing the subtraction in base 4,

$$T_{4,8} : 30133103 \mapsto 33331100 - 00113333 = 33211101, \quad (2)$$

in which the result has the Kaprekar index $(1, 4, 1, 2)$; and we would have obtained the same result if we had started with any other integer with a base-4 representation consisting of two 0's, two 1's, no 2's and four 3's.

The Kaprekar transformation can be iterated: if $T_{b,n} : x \mapsto T_{b,n}(x)$, we define $T_{b,n}^m$ recursively by

$$T_{b,n}^{m+1} : x \mapsto T_{b,n}(T_{b,n}^m(x)).$$

Since there are finitely many integers with n -digit representations in base b (for any given $b, n \in \mathbb{N}$), then for any x there must be integers $d(x) \geq 0$ and $l(x) \geq 1$ such that

$$T_{b,n}^{d(x)+l(x)}(x) = T_{b,n}^{d(x)}(x). \quad (3)$$

Taking $d(x)$ and $l(x)$ to be minimal for (3) to be satisfied, we see that iterations of the Kaprekar transformation must lead ultimately to a *cycle* of length $l(x)$, with

$$T_{b,n}^{m+l(x)}(x) = T_{b,n}^m(x)$$

for all $m \geq d(x)$. If $l(x) = 1$, we have a *fixed point*, with $T_{b,n}^m(x) = T_{b,n}^{d(x)}(x)$ for all $m \geq d(x)$. The notation and terminology is applied to Kaprekar indices as well as to integers: for example, we may observe that in base 2 the integer 1001 is a fixed point because

$$T_{2,4}(1001) = 1100 - 0011 = 1001,$$

a result which can be written in terms of Kaprekar indices as

$$T_{2,4}(2, 2) = (2, 2)$$

so that the Kaprekar index $(2, 2)$ is a fixed point in base 2. The ordered set of integers, or of their Kaprekar indices, produced by iterating the Kaprekar transformation from an integer x with Kaprekar index \mathbf{k} , is called a *Kaprekar sequence*. The integer $T_{b,n}(x)$ and its Kaprekar index $T_{b,n}(\mathbf{k})$ is called the *successor* of x and \mathbf{k} , respectively. As shorthand, we shall use primes to denote successors: $\mathbf{k}' = T_{b,n}(\mathbf{k})$, with components $\mathbf{k}' = (k'_0, k'_1, \dots, k'_{b-1})$, and multiple primes can be used to denote further terms in the Kaprekar sequence. A formula for \mathbf{k}' , or any of its components, in terms of the components of \mathbf{k} is called a *succession formula*. We shall also avoid the cumbersome $T_{b,n}^m$ notation by simply using the symbol \mapsto between successive members of a Kaprekar sequence. For example, readers may wish to check that further iterations from the calculation in (2) yield the sequence of integers,

$$\begin{aligned} 30133103 &\mapsto 33211101 \mapsto 32033211 \mapsto 32133111 \mapsto \\ &22202112 \mapsto 21033222 \mapsto 31333311 \mapsto \dots \end{aligned}$$

with its corresponding sequence of Kaprekar indices,

$$\begin{aligned} (2, 2, 0, 4) &\mapsto (1, 4, 1, 2) \mapsto (1, 2, 2, 3) \mapsto (0, 4, 1, 3) \mapsto \\ &(1, 2, 5, 0) \mapsto (1, 1, 4, 2) \mapsto (0, 3, 0, 5) \mapsto \dots \end{aligned}$$

In the companion paper [4], we show that in base 4 the integer 31333311, obtained after 6 iterations, is in fact the first member of the above sequence to be in a cycle, which is a cycle of length 3. Thus for the starting integer $x = 30133103$ we have $d(x) = 6$ and $l(x) = 3$. We also find that the Kaprekar index $(1, 1, 4, 2)$, obtained after only 5 iterations, is in a cycle of Kaprekar indices, but at that stage this index corresponds to an integer not yet in the cycle; it commonly occurs that the sequence of Kaprekar indices enters a cycle one term before the sequence of integers, since each index corresponds to many integers.

It is important to retain leading zeroes in all calculations, so that the digit-count n remains fixed while iterating with the Kaprekar transformation. For example, in base ten with $n = 3$, $T_{3,10}(877) = 877 - 778 = 99$ which we write with three digits as 099, so that the next iteration gives $T_{3,10}^2(877) = 990 - 099 = 891$. If the leading zero had been dropped, the second iteration would have given the spurious result $99 - 99 = 00$ (although in fact some authors have operated in this way). Of course, any integer which is a *repdigit* in base b , i.e., with all its digits the same, is transformed to $00 \dots 0$ (with n zeroes), which is a fixed point of the Kaprekar transformation. It has been shown by Yamagami [10] that only with 2-digit integers in an odd base can a non-repdigit be transformed to a repdigit; in all other cases, in particular whenever $n \geq 3$, we regard repdigits as a trivial case, and exclude them from any further consideration. If every non-repdigit with some given digit-count n in some base b iterates to the *same* fixed point or cycle by the Kaprekar transformation, that fixed point or cycle is termed *unanimous*.

The history of the transformation started with the observation by D. R. Kaprekar that when applied to non-trivial 4-digit integers in base ten, it always leads to the fixed point 6174 after no more than seven iterations [2, 3]; so 6174 is a unanimous fixed point for $b = 10, n = 4$. Kaprekar regarded this merely as a curiosity; but other authors have subsequently studied the transformation in various bases b and with various digit-counts n , and with rather more attention on fixed points than on cycles of length $l > 1$. A selection of these studies is listed by Yamagami [10], in a paper which also includes a complete theory of the Kaprekar transformation in base 2: the principal result for base 2 is that every Kaprekar index (k_0, k_1) with $k_0 \leq k_1$ is a fixed point, and that there do not exist cycles of length $l > 1$.

Yamagami and Matsui [11] then considered the much more complicated case of base 3, and produced a complete and rigorous theory which determines every possible fixed point and cycle, for any number of digits. At the same time, Downes-Ward [1] was using a more empirical approach to study cycles in bases 3 to 8, searching for patterns in lists of integers which are fixed points or members of cycles; those lists have been provided by Joseph Myers [6] in the *On-line Encyclopedia of Integer Sequences* (OEIS) [9], and are complete for digit-counts n up to at least 70 for bases 2 to 10. General formulae for members of cycles in base 3 were found by Downes-Ward [1], and we have checked that these formulae are consistent with the theory of Yamagami and Matsui [11].

The patterns found by Downes-Ward [1] in the data for number and lengths of cycles as a function of digit-count n are much more regular in even bases than in odd bases; in a companion paper [4] we formulate a complete theory of the Kaprekar transformation in base 4 and develop some theory that applies more generally to even bases, while in the present paper we show how Yamagami and Matsui's theory for base 3 may be generalized to cover all odd bases. We find different behaviour according to whether $b \equiv 1 \pmod{4}$ or $b \equiv 3 \pmod{4}$, and we illustrate these behaviours by examining bases 5 and 7 in detail.

Thus the remaining sections of this paper are as follows. Section 2 contains a summary of Yamagami and Matsui's results for base 3, with some further theoretical observations which are useful for consideration of higher odd bases. Section 3 contains theory that applies to all odd bases; some of the theorems here can be deduced from the formulae for difference counters and digit counters in the papers by Lapenta, Ludington, and Prichett [5, 8], but we prove them from first principles to maintain a consistent approach. The following two sections contain more specific results for bases $b \equiv 1 \pmod{4}$ and $b \equiv 3 \pmod{4}$ respectively, and we draw some general conclusions regarding odd bases in Section 6. There has been a wide variety of notation and terminology used by different researchers on the Kaprekar transformation; much of our usage is that introduced by Yamagami. One notational convention that we introduce is that the letter j always refers to place in a n -digit string, as in (1), while i always refers to a base- b digit, $0 \leq i \leq b - 1$.

2 Review of Yamagami and Matsui's results for base 3

A complete characterisation of the fixed points and cycles of the Kaprekar transformation in base 3 has been given by Yamagami and Matsui [11]. The elegance of their theory may not be apparent to readers of their paper, due to the rather lengthy proofs required to demonstrate rigorously that no possible fixed points or cycles have been omitted; so the present authors consider it worthwhile to provide a summary of the results and their theoretical background. We do not address the questions of which cycle or fixed point is reached from a given starting integer x , or the number of iterations of the Kaprekar transformation required to enter a cycle or reach a fixed point from x ; Yamagami and Matsui found relatively simple answers to these questions for odd digit-counts n , but the case of even n was found to be too complicated to develop a theory.

Yamagami and Matsui found that all fixed points and cycles in base 3 fit into a classification for which they introduced the terms *singular*, *primitive proper*, *primitive non-proper*, and *general*, with just two exceptions: with $n = 2$, the Kaprekar transformation takes all 2-digit integers to the repdigit 11 in no more than 2 iterations, and hence to the fixed point 00, with Kaprekar index $(2, 0, 0)$; while with $n = 3$, there is a unanimous cycle of length 2,

$$022 \mapsto 121 \mapsto 022,$$

with Kaprekar indices

$$(1, 0, 2) \mapsto (0, 2, 1) \mapsto (1, 0, 2).$$

The latter case was classified by Yamagami and Matsui as a singular cycle, but does not exactly fit into the pattern of singular cycles described below. For $n \geq 4$, fixed points in base 3 can simply be treated as cycles of length 1, with no separate theory required (in contrast to the situation in even bases, considered in our companion paper [4]).

The singular cycles are of length $l = m + 1$ and digit-count $n = 2^m + 2$ for each $m \in \mathbb{N}$; the Kaprekar index of one member of a cycle is $(0, 2, 2^m)$, and the remaining members are $(1, 2^j, 2^m - 2^j + 1)$ for $j = 1, \dots, m$. So with $m = 1$ we have a cycle of length 2 in 4-digit integers,

$$1221 \mapsto 1012 \mapsto 1221,$$

or in terms of Kaprekar indices,

$$(0, 2, 2) \mapsto (1, 2, 1) \mapsto (0, 2, 2).$$

The cycle of indices in the m 'th case is

$$\begin{aligned} (0, 2, 2^m) &\mapsto (1, 2, 2^m - 2 + 1) \mapsto (1, 2^2, 2^m - 2^2 + 1) \mapsto \dots \\ &\mapsto (1, 2^{m-1}, 2^m - 2^{m-1} + 1) \mapsto (1, 2^m, 1) \mapsto (0, 2, 2^m). \end{aligned}$$

Fundamental to Yamagami and Matsui's theory for the remaining classes of cycle in base 3 are the following definition and theorem.

Definition 1. For any odd integer $r \geq 3$, let $\sigma(r)$ be the least positive integer such that $2^{\sigma(r)} \equiv \pm 1 \pmod{r}$. See [A003558](#) in the OEIS.

Theorem 2. For every odd integer $r \geq 3$, $\sigma(r)$ is a divisor of $\phi(r)/2$, where ϕ is the Euler totient function ([A000010](#) in OEIS).

Euler's totient theorem states that r divides $a^{\phi(r)} - 1$ when $r \in \mathbb{N}$ and a is coprime with r ; setting $a = 2$ confirms the existence of $\sigma(r)$ satisfying Definition 1. Next, let q_0 be any even positive integer that is less than and coprime with a given odd r , and define q_i for $i > 0$ recursively by the rules:

$$q_{i+1} = \begin{cases} 2q_i, & \text{if } q_i < \frac{r}{2}; \\ 2(r - q_i), & \text{if } q_i > \frac{r}{2}. \end{cases} \quad (4)$$

Clearly $q_i \equiv \pm 2^i q_0 \pmod{r}$, so from Definition 1 we have that $q_{\sigma(r)} = q_0$ but $q_i \neq q_0$ for $1 \leq i < \sigma(r)$; note that although Definition 1 appears to allow $q_{\sigma(r)} = r - q_0$, this cannot happen since $r - q_0$ is odd. Thus we have a cycle of q -values of length $\sigma(r)$,

$$q_0 \mapsto q_1 \mapsto \cdots \mapsto q_{\sigma(r)-1} \mapsto q_{\sigma(r)} = q_0.$$

These q -cycles may be understood by reference to \mathbb{Z}_r^\times , the multiplicative group of integers modulo r , consisting of the $\phi(r)$ positive integers less than and coprime with r . The q -cycle starting from $q_0 = r - 1$ is obtained as follows: take the cyclic subgroup generated by 2 in \mathbb{Z}_r^\times , i.e., the ordered set $(1, 2, 2^2 \pmod{r}, \dots, 2^{j-1} \pmod{r})$ for some j which divides $\phi(r)$, and replace any odd members q_i^* with even integers $q_i = r - q_i^*$ (in particular, replace $q_0^* = 1$ with $q_0 = r - 1$). The subgroup may or may not include $r - 1$. If it does, the integer $r - 1$ is reached at the halfway point of generating the subgroup (since $(r - 1)^2 \equiv 1 \pmod{r}$); but the q -cycle is complete at this point. If $r - 1$ is not in the subgroup, the q -cycle is derived from the entire subgroup. Either way, corresponding to the members of the q -cycle, there are equally many odd integers $q_i^* = r - q_i$ in \mathbb{Z}_r^\times , and including the q_i^* together with the q_i accounts for $2\sigma(r)$ members of the group. This may be the entirety of the group; if not, further q -cycles are derived from cosets of our subgroup in \mathbb{Z}_r^\times , and Lagrange's Theorem then yields the result in Theorem 2. The number of q -cycles for any given odd r is then

$$N_q(r) = \frac{\phi(r)}{2\sigma(r)}. \quad (5)$$

Yamagami and Matsui found succession formulae for the 1-component of a Kaprekar index, which in the case where $k_0 = 1$ take the form

$$k'_1 = \begin{cases} 2k_1, & \text{if } 1 + k_1 < k_2; \\ 2(k_2 - 1), & \text{if } 1 < k_2 < 1 + k_1. \end{cases} \quad (6)$$

But with $k_0 = 1$, we have $k_2 = n - 1 - k_1$, and by setting $r = n - 2$ we can see that these succession formulae are precisely the rules (4) that govern q -cycles. Hence, corresponding to

each q -cycle there exists a base-3 Kaprekar cycle for integers of digit-count $n = r + 2$, with members of the cycle having Kaprekar indices of the form

$$(1, q_i, r - q_i + 1). \quad (7)$$

Yamagami and Matsui termed these Kaprekar cycles the *primitive proper cycles*. Corresponding to each of these cycles, there exist *primitive non-proper cycles* of integers with digit-count $n = cr + 2$, with members having Kaprekar indices

$$(1, cq_i, cr - cq_i + 1), \quad (8)$$

for every integer $c > 1$. Finally, corresponding to each primitive (proper or non-proper) cycle, there exist *general cycles* of integers with digit-count $n = cr + 2s$, with members having Kaprekar indices

$$(s, cq_i, cr - cq_i + s), \quad (9)$$

for every integer $s > 1$. Being derived from a q -cycle, all these primitive and general cycles have the same length, $\sigma(r)$.

The fixed points of the Kaprekar transformation in base 3 correspond to any q -cycle of length $\sigma(r) = 1$. Only $r = 3$ satisfies this requirement, with the single member of the q -cycle being $q_0 = 2$. So there is the single primitive proper fixed point with Kaprekar index $(1, 2, 2)$ for $n = 5$, primitive non-proper fixed points $(1, 2c, c + 1)$ for $n = 3c + 2$, and general fixed points $(s, 2c, c + s)$ for $n = 3c + 2s$.

While the formulae to generate all cycles and fixed points are simple, if it is required to determine the cycles and fixed points with a given digit-count n , the following procedure needs to be followed. If n is odd with $n \geq 5$, there is at least one primitive proper cycle. The number of such cycles is $N_q(r)$, found from (5) where $r = n - 2$. We have seen that a q -cycle can always be generated using the rules (4) from $q_0 = r - 1$; this q -cycle corresponds to a subgroup of \mathbb{Z}_r^\times . If $N_q(r) > 1$, further q -cycles corresponding to cosets in \mathbb{Z}_r^\times may be found by taking q_0 as the greatest even positive integer less than and coprime with r and not appearing in a cycle already found. The primitive proper cycles of Kaprekar indices are then obtained from the q -cycles by the formula (7). If $n - 2$ is prime, there are no non-proper cycles. If $n - 2$ is composite (odd or even), there are non-proper cycles corresponding to each of its odd factors $r \geq 3$ such that $cr = n - 2$ for some $c \in \mathbb{N}$; having obtained the q -cycle(s) for each such r , the non-proper cycles of Kaprekar indices of form (8) can be calculated using the multiplier c . Clearly the only primitive cycles for even n are non-proper cycles with even c ; but if $n = 2^m + 2$ for any $m \in \mathbb{N}$, which is the case where a singular cycle exists, there are no primitive cycles at all. Finally, there are general cycles for the given n corresponding to the primitive cycles for digit-counts $n - 2t$, for each $t \in \mathbb{N}$ such that $n - 2t \geq 5$ or $n - 2t \geq 8$ in the respective cases of odd or even n (since 5 and 8 are the smallest digit-counts for which primitive cycles exist in the respective odd and even cases).

Hence the enumeration of general cycles for a given n requires the enumeration of primitive cycles for all smaller positive integers n' of the same parity, while the enumeration of

primitive cycles involves the summation of $N_q(r)$ for all odd r that are factors of $n' - 2$. This leads to Yamagami and Matsui's [11] formulae involving double summations for the total number of cycles (including fixed points) with any given n . However, simpler formulae may be found for the enumeration of integers which are members of a cycle with given n . For odd n with $n - 2$ prime, each positive even number less than $n - 2$ is in a q -cycle for $r = n - 2$, and hence appears as the 1-component of the Kaprekar index of a member of a primitive proper cycle; so there are $(n - 3)/2$ members of such cycles. If n is odd but with $n - 2$ composite, each even number less than $n - 2$ still appears as the 1-component of the Kaprekar index of a member of a primitive cycle which may be proper or non-proper, so the formula $(n - 3)/2$ remains valid as the number of members of all primitive cycles with the given digit-count n . The summation to yield $N_K(n)$, the number of integers which are members of any cycle, primitive or general, with odd digit-count n is then

$$N_K(n) = \sum_{\text{odd } n', 5 \leq n' \leq n} \frac{n' - 3}{2} = \sum_{m'=1}^{(n-3)/2} m' = \frac{(n - 3)(n - 1)}{8}. \quad (10)$$

For even n , we note first that any positive integer m may be written in terms of a 2-adic valuation $v(m)$ and an odd part $\rho(m)$ as

$$m = 2^{v(m)}\rho(m). \quad (11)$$

Since there are primitive cycles corresponding to each of the odd factors of $n - 2$, the number of integers which are members of primitive cycles with digit-count n is $(\rho(n - 2) - 1)/2$. The summation to enumerate members of all cycles, primitive or general, must be made over even n' from 8 to n , but does not yield a simple formula for $N_K(n)$ since there is no simple formula for $\rho(n - 2)$. If $n = 2^m + 2$ for any $m \in \mathbb{N}$, there are $m + 1$ members of a singular cycle, but no members of primitive cycles since $\rho(n - 2) = 1$.

A table of q -cycles for all odd r up to 67 is given in [Appendix A](#), and [Figure 1](#) shows correspondingly all the cycle lengths l that exist for each digit-count n up to 70: cycle lengths are in sequence [A165003](#) in OEIS, with digit-counts obtained by converting least members of cycles, [A165002](#) in OEIS, to base 3. For primitive proper cycles, for which the cycle length is $\sigma(r) = \phi(r)/2N_q(r)$ with $r = n - 2$, the greatest possible value of $\sigma(r)$ is when $N_q(r) = 1$ and r is prime so that $\phi(r) = r - 1$. Any non-proper or general cycle of the same length must have a greater digit-count. Thus the leftmost dot on almost every row in [Figure 1](#) represents a primitive proper cycle, and is on the line $l = (n - 3)/2$ if $n - 2$ is prime with $N_q(n - 2) = 1$. To the right of the leftmost dot on any row, there are further dots representing general cycles at every greater odd digit-count, but the first dot for an even digit-count is for a primitive non-proper cycle with $c = 2$; this has $n = 2n_0 - 2$, where n_0 is the digit-count for the proper cycle. The exceptions to the above are dots representing singular cycles at $n = 4, 6, 10, 18, 34$ and 66 for cycle lengths up to 7, and a dot for the special case of $n = 3$; no dot is shown for $n = 2$ because we have excluded fixed points with Kaprekar indices $(n, 0, 0)$ as being trivial, but all 2-digit integers are taken to such a fixed point under the Kaprekar transformation. Apart from these exceptions, there are no dots to the left of the line $l = (n - 3)/2$.




Figure 1: Cycle lengths that exist for digit-counts n up to 70.

We conclude this section with an example of finding all the primitive Kaprekar cycles with a given digit-count: $n = 254$. Since n is even, we are only looking for non-proper cycles. The odd factors of $n - 2 = 252$ are $r = 3, 7, 9, 21$ and 63 , with respective multipliers $c = 84, 36, 28, 12$ and 4 such that $cr = n - 2$. So, from the table in [Appendix A](#) and formula (8), we have:

- Corresponding to $r = 3$, a fixed point with Kaprekar index $(1, 168, 85)$;
- Corresponding to $r = 7$, a cycle of length 3, members having Kaprekar indices

$$(1, 72, 181) \mapsto (1, 144, 109) \mapsto (1, 216, 37) \mapsto (1, 72, 181);$$
- Corresponding to $r = 9$, a cycle of length 3, members having Kaprekar indices

$$(1, 56, 197) \mapsto (1, 112, 141) \mapsto (1, 224, 29) \mapsto (1, 56, 197);$$
- Corresponding to $r = 21$, a cycle of length 6, members having Kaprekar indices

$$(1, 24, 229) \mapsto (1, 48, 205) \mapsto (1, 96, 157) \mapsto (1, 192, 61) \mapsto (1, 120, 133) \mapsto (1, 240, 13) \mapsto (1, 24, 229);$$
- Corresponding to $r = 63$, three cycles of length 6, members having Kaprekar indices

$$(1, 8, 245) \mapsto (1, 16, 237) \mapsto (1, 32, 221) \mapsto (1, 64, 189) \mapsto (1, 128, 125) \mapsto (1, 248, 5) \mapsto (1, 8, 245),$$

$$(1, 40, 213) \mapsto (1, 80, 173) \mapsto (1, 160, 93) \mapsto (1, 184, 69) \mapsto (1, 136, 117) \mapsto \\ (1, 232, 21) \mapsto (1, 40, 213),$$

$$(1, 88, 165) \mapsto (1, 176, 77) \mapsto (1, 152, 101) \mapsto (1, 200, 53) \mapsto (1, 104, 149) \mapsto \\ (1, 208, 45) \mapsto (1, 88, 165).$$

Finding the actual 254-digit integer in a cycle from its Kaprekar index simply requires a base-3 subtraction of integers having the previous Kaprekar index in the cycle. For example, the integer with Kaprekar index $(1, 72, 181)$ in the cycle corresponding to $r = 7$ is found from the Kaprekar index $(1, 216, 37)$ by

$$\begin{array}{r} \underbrace{2 \dots \dots 2}_{37} \quad \underbrace{1 \dots \dots 10}_{216} \\ - 0 \underbrace{1 \dots \dots 1}_{216} \quad \underbrace{2 \dots 2}_{37} \\ = 2 \underbrace{1 \dots 10}_{35} \quad \underbrace{2 \dots 2}_{180} \quad \underbrace{1 \dots 11}_{36}, \end{array}$$

where the result has Kaprekar index $(1, 72, 181)$ as required.

Observe that every multiple of 8 that is less than 252 appears once as the 1-component of a Kaprekar index in the above primitive cycles. With $n = 254$ we have $\rho(n - 2) = 63$ and $v(n - 2) = 2$ according to the definition (11), so there is a member of a primitive cycle corresponding to each even positive integer less than 63; and the 1-components of the Kaprekar indices of these cycle members are even multiples of $2^{v(n-2)} = 4$.

3 Theory on primitive and general cycles in any base

It is useful to reframe some relevant parts of the theory for base 3 before considering general bases. We introduce the notation

$$k_2^- := k_2 - 1. \tag{12}$$

Then for odd digit-counts n in base 3, every Kaprekar index of the form $(1, k_1, 1 + k_2^-)$ with even positive k_1 , so that k_2^- is odd and $k_1 + k_2^- = n - 2$, is in a primitive cycle. Let $c := \gcd(n - 2, k_1) = \gcd(k_1, k_2^-)$ and $r := (n - 2)/c$; then the length l of the primitive cycle is equal to $\sigma(r)$, defined in Definition 1. If $c = 1$, the primitive cycle is proper. Letting $q = k_1/c$, the Kaprekar index $(1, k_1, 1 + k_2^-)$ can be rewritten as $(1, cq, cr - cq + 1)$, and is in a cycle in which the succession of indices is given by a q -cycle derived from a subgroup or coset of the multiplicative group modulo r . With odd n , c is also odd; but non-proper primitive cycles with even n may also be obtained from proper cycles by applying even multipliers c in cycles of Kaprekar indices of form $(1, cq, cr - cq + 1)$. Furthermore, indices of form $(s, cq, cr - cq + s)$, for any $s > 1$, are in associated general cycles. This theory leads to the pattern of cycle lengths vs. digit-counts seen in Figure 1.

We now introduce some terminology for cycles and individual Kaprekar indices, some of which is a generalization to any base $b \geq 3$ of the terminology used by Yamagami and Matsui [11] for base 3. To make the notation tidier, we introduce the symbol

$$B := b - 1$$

for the greatest digit that appears in base b .

Definition 3.

- A Kaprekar index $\mathbf{k} = (k_0, k_1, \dots, k_B)$ is *regular* if $k_i \geq 1$ for $i = 0, 1, 2, \dots, B$ and $k_B > k_0$.
- A cycle is *regular* if every member has a regular Kaprekar index.
- A cycle is *special* if any member has a Kaprekar index that is not regular.

Thus in base 3, Yamagami and Matsui's singular cycles are classified here as *special*, as are the unanimous fixed point with $n = 2$ and the unanimous cycle of length 2 with $n = 3$. However, our concern in this section is to develop theory applying to regular cycles. These cycles and their members may be classified using an extension of Yamagami and Matsui's terminology.

Definition 4.

- A regular Kaprekar index is *primitive* if $k_0 = 1$.
- A cycle is *primitive* if each member has a Kaprekar index that is primitive.
- A regular Kaprekar index is *general* if $k_0 > 1$.
- A cycle is *general* if each member has a general Kaprekar index.

Clearly, the existence of regular cycles depends on whether regular Kaprekar indices \mathbf{k} can be guaranteed to have regular successors \mathbf{k}' . It is therefore useful to have a general formula for the digits of a successor. Such a formula was given by Prichett, Ludington and Lapenta [8, 5]; in a variant of their difference notation, we define

$$d_j := a_{n-1-j} - a_j \quad \text{for } j = 0, 1, \dots, \nu, \tag{13}$$

where $\nu = \lfloor n/2 \rfloor - 1$, and we observe that $d_0 \geq d_1 \geq \dots \geq d_\nu$. We set $\mu = \max\{j : d_j > 0\}$, so either $d_{\mu+1} = 0$ or $\mu = \nu$. Performing the subtraction (1), the digits of $T_{b,n}(x)$ are then found to be [8]

$$d_0, d_1, \dots, d_{\mu-1}, d_\mu - 1, B, \dots, B, B - d_\mu, B - d_{\mu-1}, \dots, B - d_1, b - d_0, \tag{14}$$

where commas have been inserted between digits for clarity since some digits are in the form of expressions involving a subtraction. The digit string (14) may be subdivided into three

sections. On the right, there are $\mu + 1$ instances of digits a_{n-1-j} of x_\uparrow being subtracted from smaller digits a_j of x_\downarrow : the rightmost digit is $b - (a_0 - a_{n-1})$, and then 1 is carried to the left so that each subsequent digit subtraction (going leftward) yields $b - (a_j - a_{n-j-1}) - 1 = B - d_j$. In the centre, there are $n - 2\mu - 2$ instances where $a_{n-1-j} = a_j$, and the subtraction with 1 carried yields B ; but if $\mu = \nu$, then this central string of digits B is absent if n is even, or is reduced to a single such digit if n is odd. On the left, there are $\mu + 1$ subtractions of a_{n-1-j} from larger digits a_j , yielding d_j except that 1 is still carried into the rightmost subtraction of this section.

Suppose that x has a regular Kaprekar index \mathbf{k} , so the base- b representations of x_\downarrow and x_\uparrow consist of blocks of k_i digits i , with $k_i \geq 1$ for every value of i between 0 and B , arranged in descending and ascending order, respectively. Given any i_\uparrow and i_\downarrow with $0 \leq i_\uparrow < i_\downarrow \leq B$, let K_{i_\uparrow} and K_{i_\downarrow} be the number of digits (counting from the left) appearing in the base- b representations of x_\uparrow and x_\downarrow up to the last occurrence of the digit i_\uparrow and i_\downarrow , respectively:

$$K_{i_\uparrow} := \sum_{i=0}^{i_\uparrow} k_i \quad \text{and} \quad K_{i_\downarrow} := \sum_{i=i_\downarrow}^B k_i.$$

So, going along the digit string (14) from the left, the value of the difference d_j decreases by 1 at positions $j + 1 = K_{i_\uparrow}$ and $j + 1 = K_{i_\downarrow}$ for each $i_\uparrow = 0, 1, \dots$ and each $i_\downarrow = B, B - 1, \dots$, but decreases by 2 at a position where $K_{i_\uparrow} = K_{i_\downarrow}$ for some i_\uparrow and i_\downarrow . Thus the differences d_j ($j = 0, 1, \dots, \mu$) take every possible value from B down to 1 if the condition $K_{i_\uparrow} = K_{i_\downarrow}$ never occurs. This motivates the following definition and theorem.

Definition 5. A Kaprekar index $\mathbf{k} = (k_0, k_1, \dots, k_B)$ is deemed to satisfy the *inequality condition* if

$$\sum_{i=0}^{i_\uparrow} k_i \neq \sum_{i=i_\downarrow}^B k_i \tag{15}$$

for every choice of i_\uparrow, i_\downarrow with $0 \leq i_\uparrow < i_\downarrow \leq B$.

Theorem 6. *If a Kaprekar index \mathbf{k} is primitive or general and satisfies the inequality condition, then its successor \mathbf{k}' is primitive or general, respectively. Furthermore, the components of the successor satisfy $k'_i \geq 2$ for $i = 1, 2, \dots, B$.*

Proof. First observe that the central section of digits B in (14) is absent only if the inequality condition is violated, with $K_{i_\uparrow} = K_{i_\downarrow} = n/2$ for some i_\uparrow and $i_\downarrow = i_\uparrow + 1$. The discussion above verifies that every digit from 1 to B is present among the differences d_0, \dots, d_μ if \mathbf{k} is regular and satisfies the inequality condition, and so every digit from 0 to $B - 1$ is present among values of $B - d_j$ ($j = 0, 1, \dots, \mu$). Also, each instance of 0 among values of $B - d_j$ corresponds to an instance of B among the d_j , and $B - d_j = 0$ if and only if $a_j = 0$ (because $k_B > k_0$ in a regular Kaprekar index, so $a_{n-1-j} = B$ whenever $a_j = 0$).

We would now have proved the theorem if it were not for having “pattern-breaking” digits, $d_\mu - 1$ instead of d_μ in the left section of (14), and $b - d_0$ instead of $B - d_0$ as the

rightmost digit. Indeed, we would have each digit from 1 to $B - 1$ appearing in both the left and right sections of (14), so $k'_i \geq 2$ for $i = 1, 2, \dots, B - 1$; at least one digit B in the central section and in the left section, with each of the latter corresponding to a digit 0 in the right section, so that $k_B > k_0 \geq 1$; and $k'_0 = k_0$, so that primitive and general Kaprekar indices have respective primitive and general successors. So it remains to resolve the issue of the two pattern-breaking digits; but the inequality condition ensures that $d_\mu = 1$, so having $d_\mu - 1$ in (14) provides a digit 0 instead of 1; and $d_0 = B$, so having $b - d_0$ in (14) provides a digit 1 instead of 0. Hence the pattern-breaking does not affect the Kaprekar index of the successor, under the given conditions. \square

Note that the inequality condition is sufficient but not necessary in this theorem: the successor of a regular Kaprekar index may be regular and may have $k'_i \geq 2$ for $i = 1, 2, \dots, B$ even when \mathbf{k} does not satisfy the inequality condition. On the other hand, finding a regular Kaprekar index with a regular successor does not imply that they are members of a regular cycle: our next theorem enables us to write down infinitely many more regular cycles once a primitive cycle has been found, but it is of little help in finding a primitive cycle in the first place. It will be useful for this theorem to generalize the notation (12) as

$$k_B^- := k_B - 1,$$

so that a primitive Kaprekar index takes the form,

$$(1, k_1, \dots, k_{B-1}, k_B^- + 1).$$

Theorem 7. *If the primitive Kaprekar index $\mathbf{k} = (1, k_1, \dots, k_{B-1}, k_B^- + 1)$ has a primitive successor $\mathbf{k}' = (1, k'_1, \dots, k'_{B-1}, k_B^- + 1)$, then:*

(i) *the primitive Kaprekar index $(1, ck_1, \dots, ck_{B-1}, ck_B^- + 1)$ has the primitive successor $(1, ck'_1, \dots, ck'_{B-1}, ck_B^- + 1)$, for any integer $c > 1$;*

(ii) *the general Kaprekar index $(s, ck_1, \dots, ck_{B-1}, ck_B^- + s)$ has the general successor $(s, ck'_1, \dots, ck'_{B-1}, ck_B^- + s)$, for any integers $s > 1$ and $c \geq 1$.*

Proof. (i) Multiplying components k_1, \dots, k_{B-1} and k_B^- of the primitive Kaprekar index \mathbf{k} by c means that each digit in the original integers x_\downarrow and x_\uparrow is replaced with c copies of the same digit, except for the first and last digits. So in (14) the first digit d_0 and last digit $b - d_0$ are unchanged, but every other subtraction $a_{n-1-j} - a_j$ is replaced with c copies of the same subtraction. This yields c copies of the same digit in (14), except that there is still only one instance of the pattern-breaking digit $d_\mu - 1$ arising from the last subtraction from the right to have 1 carried. As in the proof of Theorem 6, this compensates for the pattern-breaking rightmost digit $b - d_0$.

(ii) Increasing each of k_0 and k_B by $s - 1$ means that we leave the leftmost and rightmost digits of x_\downarrow and x_\uparrow unchanged, but insert $s - 1$ digits B adjacent to the leftmost B of x_\downarrow and the rightmost B of x_\uparrow , together with $s - 1$ digits 0 adjacent to the rightmost 0 of x_\downarrow and the leftmost 0 of x_\uparrow . The subtractions then yield $s - 1$ further digits B (from $B - 0$) on the left of (14), and $s - 1$ further digits 0 (from $0 - B$ with 1 carried) on the right of (14). \square

Corollary 8. *If the primitive Kaprekar index $(1, k_1, \dots, k_{B-1}, k_B^- + 1)$ is a member of a primitive cycle, then:*

(i) *the Kaprekar index $(1, ck_1, \dots, ck_{B-1}, ck_B^- + 1)$, for any integer $c > 1$, is also a member of a primitive cycle of the same length;*

(ii) *the Kaprekar index $(s, ck_1, \dots, ck_{B-1}, ck_B^- + s)$, for any integers $s > 1$ and $c \geq 1$, is a member of a general cycle of the same length.*

This follows from Theorem 7 because the c -multiplication and the s -addition can be applied to each succession in the original primitive cycle.

Part (i) of Theorem 7 motivates the following definitions, which are again an extension to general bases $b \geq 3$ of a definition given for base 3 by Yamagami and Matsui.

Definition 9.

- A Kaprekar index is *primitive proper* if it is primitive with

$$\gcd(k_1, \dots, k_{B-1}, k_B^-) = 1.$$

- A Kaprekar index is *primitive non-proper* if it is primitive with $\gcd(k_1, \dots, k_{B-1}, k_B^-) = c$ for some integer $c > 1$.

It follows from part (i) of Theorem 7 that if the successor of a primitive proper index is primitive, then it is proper, and if the successor of a primitive non-proper index is primitive, then it is non-proper with the same multiplier c . Hence we can apply similar definitions to cycles.

Definition 10.

- A cycle is *primitive proper* if its members are primitive proper.
- A cycle is *primitive non-proper* if its members are primitive non-proper.

Once we have found a primitive proper cycle, Corollary 8 shows that there are then infinitely many primitive non-proper cycles corresponding to it, and then infinitely many general cycles corresponding to each primitive (proper or non-proper) cycle. In base 3, each primitive proper cycle corresponds in a simple way to a q -cycle derived from a subgroup (generated by 2) or coset in \mathbb{Z}_r^\times . The next theorem suggests a link between q -cycles and primitive proper cycles in any odd base, whereas no similar theorem applies in even bases.

Theorem 11. *In base $b = B + 1$, where B is even, the formula for $k'_{B/2}$ in terms of the components of \mathbf{k} includes a factor of 2 if \mathbf{k} is regular and satisfies the inequality condition.*

Proof. If $d_j = B/2$, then $B - d_j = B/2$ also. Hence each instance of the digit $B/2$ in the right section of (14) corresponds to an instance of $B/2$ in the left section. So the total number of instances, $k'_{B/2}$, must include a factor of 2. \square

In base 3, for which $B/2 = 1$, Yamagami and Matsui's [11] Theorem 1.3 shows the factor of 2 appearing in the formula for k'_1 in every case, not merely those in which \mathbf{k} is regular and satisfies the inequality condition. In particular, when those conditions are satisfied, we have the succession formulae (6) for k'_1 , which relate in a simple way to q -cycles since the latter are generated by the factor 2 in the multiplicative group modulo $n - 2$. However, in higher odd bases, we shall find that the succession formulae retain the connection between q -cycles and cycles in the Kaprekar transformation in bases $b \equiv 3 \pmod{4}$ but not in bases $b \equiv 1 \pmod{4}$.

A further theorem that applies to all odd bases and may be of some use, at least to reduce the burden of calculation, is that the Kaprekar index of a successor \mathbf{k}' displays a symmetry (except between k_0 and k_B) when \mathbf{k} and \mathbf{k}' are both regular:

Theorem 12. *In base $b = B + 1$ where B is even, if \mathbf{k} is regular and satisfies the inequality condition, then its successor has $k'_{B-i} = k'_i$ for $1 \leq i \leq B - 1$.*

Proof. Since each digit d_j in the left section of (14) corresponds to a digit $B - d_j$ in the right section (with the pattern-breaking digits compensating each other as explained above), the count of any digit i equals that of the digit $B - i$. The exception is for digits 0 and B , since the central section of digits B has no corresponding digits 0. \square

We conclude this section by observing that according to Theorem 6 the successor to a regular Kaprekar index satisfying the inequality condition must have a digit-count $n \geq 2b - 1$, since $k'_0 \geq 1$ and $k'_i \geq 2$ for $i = 1, \dots, B$. In fact, a Kaprekar index with $k_i = 2$ for $i = 1, \dots, B$ corresponds to a fixed point in any odd base. This is seen most easily from the example of base 5, where we have the subtraction,

$$\begin{aligned} & 443322110 \\ & - 011223344 \\ & = 432043211, \end{aligned} \tag{16}$$

from which it is clear that in any greater odd base b the fixed-point integer is

$$B \dots 20B \dots 211, \tag{17}$$

where each ellipsis indicates that there is a complete sequence of decreasing digits from $B (= b - 1)$ to 2. We shall see that the fixed point of this form does have the least digit-count of any regular fixed point in bases $b \equiv 1 \pmod{4}$, but that regular fixed points with smaller digit-counts (and not satisfying the inequality condition) exist in bases $b \equiv 3 \pmod{4}$.

4 Base 5 and higher bases $b \equiv 1 \pmod{4}$

Much of this section is focussed on base 5; we have also examined cycles in base 9, enabling us to draw some general conclusions about bases $b \equiv 1 \pmod{4}$, although the only rigorous

theory beyond that in Section 3 is for base 5. Least members of base 5 cycles are given in base ten in [A165041](#) in OEIS, with corresponding cycle lengths in [A165042](#); we convert the least members to base 5 and then compute further cycle members to obtain the cycles displayed below and in the appendices. For base 9 the corresponding OEIS sequences are [A165119](#) (least members of cycles) and [A165120](#) (cycle lengths).

We first display in Figure 2 the base-5 equivalent to Figure 1 for base 3, showing the cycle lengths that exist for each digit-count, although only up to $n = 50$ in the present case. The pattern is similar in the two figures, with points representing primitive proper cycles on or to the right of the line $l = (n - 3)/2$, general cycles represented at intervals of 2 in n to the right of the primitive proper cycles, and points with even n representing non-proper cycles with multiplier $c = 2$ appearing further to the right. However, there are many more exceptions to this pattern, representing special cycles, in base 5 than in base 3; in particular, there are many points to the left of the line $l = (n - 3)/2$ for small values of n in Figure 2.




Figure 2: Cycle lengths that exist for digit-counts n up to 50 in base 5.

A list of special cycles for digit-counts up to $n = 50$ in base 5 is given in [Appendix B](#); there certainly exist special cycles for greater digit-counts. Regular cycles do not exist for $n < 9$, so all cycles (including fixed points) with digit-counts $n \leq 8$ are special; they are unanimous for $3 \leq n \leq 8$, as are the special cycles with $n = 10$ and $n = 12$, but with $n = 2$ both the fixed point 13 and the repdigit fixed point 00 can be reached from different non-trivial starting integers. Among the special cycles there is a class of singular cycles, with digit-counts $n = 2^m + 2$ and lengths $l = m + 1$ for integers m as in base 3, but only for $m \geq 3$ in base 5. The structure of these singular cycles is more complicated in base 5 than in base 3: the sequence of Kaprekar indices in these cycles is

$$\begin{aligned}
 (1, 0, 1, 2, 2^m - 2) &\mapsto (1, 2, 2, 2, 2^m - 6 + 1) \mapsto (1, 2^2, 2^2, 2^2, 2^m - 12 + 1) \mapsto \dots \\
 &\mapsto (1, 2^{m-2}, 2^{m-2}, 2^{m-2}, 2^m - 3 \times 2^{m-2} + 1) \mapsto (1, 2^{m-1}, 0, 2^{m-1}, 1) \\
 &\mapsto (0, 2, 2^m - 1, 0, 1) \mapsto (1, 0, 1, 2, 2^m - 2).
 \end{aligned}$$

The cycles with $n = 10$, $n = 18$ and $n = 34$ in [Appendix B](#) are examples of this type. While the remaining special cycles in [Appendix B](#) display many common features, we have not been able to find any further clear patterns analogous to that of the singular cycles.

In base 9 we have also found numerous special cycles. There does not appear to be any equivalent of the singular cycles in bases 3 and 5, but there are interesting classes of fixed points and cycles of length $l = 2$. Specifically, for any integers $t \geq 1$ and $u \geq 0$, there are fixed points with Kaprekar indices

$$(0, 0, t, t + u, 0, t + u, t, 0, t + u) \quad \text{and} \quad (0, t, t + u, 2t + u, 0, 2t + u, t + u, t, t + u),$$

with respective digit-counts $n = 5t + 3u$ and $n = 9t + 5u$. Apart from the repdigit fixed point 00 at $n = 2$, we have not found any other special fixed points. For any integer $t \geq 0$ there is a cycle of length 2,

$$(0, 1, t, t + 2, 0, t + 1, t + 2, 0, t + 1) \mapsto (0, 0, t + 2, t + 1, 0, t + 2, t, 1, t + 1) \mapsto \\ (0, 1, t, t + 2, 0, t + 1, t + 2, 0, t + 1),$$

with digit-count $n = 5t + 7$. There are also many other special cycles of lengths $l \geq 2$ in base 9, which do not appear to fit into this or any other class. We have not formulated any further theory of special cycles or fixed points in bases $b \equiv 1 \pmod{4}$.

To develop theory for regular cycles, we require succession formulae, of which a partial list for base 5 is given in [Appendix C](#). Formula (14) gives the individual digits of a successor, but the practical means of finding a succession formula is to set out the calculation as in the following example, which is for case (v) of [Appendix C](#):

$$\begin{array}{r} \underbrace{4 \cdots 4}_{k_4} \quad \underbrace{3 \cdots \cdots \cdots 3}_{k_3} \quad \underbrace{2 \cdots \cdots \cdots 2}_{k_2} \quad \underbrace{1 \cdots \cdots \cdots 1}_{k_1} \quad 0 \\ - 0 \underbrace{1 \cdots \cdots \cdots 1}_{k_1} \quad \underbrace{2 \cdots \cdots \cdots 2}_{k_2} \quad \underbrace{3 \cdots \cdots \cdots 3}_{k_3} \quad \underbrace{4 \cdots \cdots \cdots 4}_{k_4} \\ = 4 \underbrace{3 \cdots \cdots 3}_{k_4-1} \quad \underbrace{2 \cdots \cdots 2}_{1+k_1-k_4} \quad \underbrace{1 \cdots \cdots 1}_{k_4+k_3-k_1-2} \quad 0 \underbrace{4 \cdots \cdots \cdots 4}_{1+k_1+k_2-k_3-k_4} \quad \underbrace{3 \cdots \cdots \cdots 3}_{k_3+k_4-k_1-1} \quad \underbrace{2 \cdots \cdots \cdots 2}_{1+k_1-k_4} \quad \underbrace{1 \cdots \cdots \cdots 1}_{k_4}. \end{array} \quad (18)$$

The list in [Appendix C](#) covers all cases where \mathbf{k} is primitive (and also some other cases, in particular where $k_4 = 1 = k_0$); from [Theorem 7](#), succession formulae when \mathbf{k} is general follow from primitive cases, so all regular Kaprekar indices are covered. Referring to these formulae, we can prove the next two theorems.

Theorem 13. *Every member of a regular cycle in base 5 has a Kaprekar index satisfying the inequality condition.*

Proof. In each case in [Appendix C](#) where the inequality condition is not satisfied, i.e. where the conditions on k_1, k_2, k_3 and k_4 include an equality, the successor \mathbf{k}' is not regular: either $k'_0 = 0$ or $k'_2 = 0$ or $k'_4 = 1 = k'_0$. \square

Note that this theorem is not trivial: in base 7 we shall see that the inequality condition is not necessary for membership of a regular cycle. The inequality condition is not sufficient for a regular Kaprekar index to be a cycle member in any base $b \geq 5$; but stricter conditions that do appear to be sufficient in base 5 only are displayed below.

An immediate consequence of this theorem, according to Theorem 6, is that every member of a regular cycle in base 5 has $k_i \geq 2$ for $i = 1, 2, 3, 4$. The case where $k_i = 2$ for $1 \leq i \leq 4$ is the fixed point (16), and we have a further theorem regarding this fixed point.

Theorem 14. *The only primitive proper fixed point in base 5 is that displayed in (16) above, with Kaprekar index $\mathbf{k} = (1, 2, 2, 2, 2)$.*

Proof. Setting $k'_i = k_i$ for $i = 0, 1, 2, 3, 4$ in each of the succession formulae in Appendix C, we obtain an inconsistency in every case except (v). In that case, setting $k'_i = k_i$ yields that $k_1 = k_2 = k_3 = 2(k_4 - 1)$, which allows just the Kaprekar index (1, 2, 2, 2, 2) or any primitive non-proper index derived from it according to Theorem 7(i). \square

Based on a survey of regular cycles and fixed points with digit-counts up to $n = 35$ in base 9, we conjecture that the last two theorems apply in all bases $b \equiv 1 \pmod{4}$: that all members of regular cycles in these bases satisfy the inequality condition, and so have $k_i \geq 2$ for $1 \leq i \leq b - 1$, implying that regular cycles cannot exist with digit-count $n \leq 2b - 1$; and that the only primitive proper fixed point in any such base is (17). The theorems for base 5 were proved by exhaustive consideration of succession formulae for that particular base, so a more elegant method would need to be devised in order to prove the conjectures for higher bases. And in any case there is evidence from base 9 of complications that may cast doubt on the truth of such conjectures: there is a set of “pseudo-regular” cycles of length $l = 2$ in which one member has $k_4 = 0$ (so is not regular), the other member does not satisfy the inequality condition, but the structure of primitive and general cycles specified in Corollary 8 is retained: see the note at the end of Appendix E.

Given that the inequality condition is definitely satisfied by cycle members in base 5, the components of their Kaprekar indices are constrained by Theorems 11 and 12 to satisfy the requirements,

$$k_2 \text{ is even; } \quad k_3 = k_1 \geq 2. \quad (19)$$

If we restrict attention to primitive Kaprekar indices, for which $k_0 = 1$, then once the digit-count n is fixed, only two free parameters remain in an index. These can be taken as k_1 and k_2 , and we then have $k_4 = n - 2k_1 - k_2 - 1$. If we restrict attention further to odd values of n , the Kaprekar indices under consideration are either proper, or non-proper with an odd multiplier c . Then, given the conditions (19), a further condition that suffices to ensure that a Kaprekar index satisfies the inequality condition is as follows:

$$\text{either } k_1 \text{ is even, or } k_1 \text{ is odd and } k_1 > k_4. \quad (20)$$

To see why conditions (20) apply, note first that k_4 is even when n is odd. Then if k_1 is even, the left side of (15) is odd (since $k_0 = 1$) while the right side is even, guaranteeing

the inequality; while if k_1 is odd the further condition $k_1 > k_4$ precludes equalities such as $k_0 + k_1 = k_4$. In higher bases, the free parameters are k_i for $i = 1, 2, \dots, B/2$, and it remains true that the inequality condition is satisfied if these are all even; but we have not determined more general principles governing when the inequality condition is satisfied.

In base 3, all primitive cycles could be derived from q -cycles, as tabulated in [Appendix A](#) for $r \leq 67$, because of the simple succession formulae (6) for a single free parameter. In base 5, having established that (19) applies throughout any regular cycle, we only require succession formulae for the two free parameters k'_1 and k'_2 , which are as follows (where we have taken all cases from [Appendix C](#) in which \mathbf{k} is regular and satisfies the inequality condition, with $k_3 = k_1$):

$$k'_1 = \begin{cases} 2k_4 - 2, & \text{if } k_4 < 1 + k_1 \text{ and } k_4 < 1 + k_2; \\ k_2 + k_4 - 1, & \text{if } 1 + k_2 < k_4 < 1 + k_1; \\ 2k_1, & \text{if } 1 + k_1 < k_4 < 1 + k_2; \\ 2k_1 + k_2 - k_4 + 1, & \text{if } 1 + k_1 < k_4 \text{ and } 1 + k_2 < k_4 < 1 + k_1 + k_2; \\ k_4 - k_2 - 1, & \text{if } 1 + k_1 + k_2 < k_4 < 1 + 2k_1 + k_2; \\ 2k_1, & \text{if } 1 + 2k_1 + k_2 < k_4. \end{cases} \quad (21)$$

$$k'_2 = \begin{cases} 2(k_1 - k_4 + 1), & \text{if } k_4 < 1 + k_1; \\ 2(k_4 - k_1 - 1), & \text{if } 1 + k_1 < k_4 < 1 + k_1 + k_2; \\ 2k_2, & \text{if } 1 + k_1 + k_2 < k_4. \end{cases} \quad (22)$$

We may observe the factors of 2 appearing in all the formulae for k'_2 , in accord with [Theorem 11](#). However, unlike in base 3, there is no simple equivalence between succession formulae and the q -cycle formulae (4): whereas in base 3 we have $k'_1 \equiv \pm 2k_1 \pmod{(n-2)}$ in all cases, in base 5 the first two cases in (22) yield $k'_2 \equiv \pm 2(k_2 + 3k_1) \pmod{(n-2)}$. Hence the link between q -cycles and cycles of the Kaprekar transformation is broken in base 5 (and higher bases $b \equiv 1 \pmod{4}$), because the succession formulae do not correspond to simple multiplication by $\pm 2 \pmod{(n-2)}$ of a component of \mathbf{k} . So whereas in base 3 the length l of a primitive proper cycle is strictly equal to $\sigma(n-2)$, this does not apply in bases $b \equiv 1 \pmod{4}$; although in base 5 there is empirical evidence (from checking all primitive proper cycles with $n \leq 41$) that l is either equal to or is a divisor of $\sigma(n-2)$ (with equality applying whenever $\sigma(n-2)$ is prime and $n > 9$, due to [Theorem 14](#)).

Since the primitive proper cycles cannot be simply derived from q -cycles in bases $b \equiv 1 \pmod{4}$, we have listed them for $n \leq 33$ in [Appendix D](#) for base 5 and [Appendix E](#) for base 9. Comparing with values of $\sigma(r)$ in [Appendix A](#), we may observe that in the majority of cases, the lengths l of the primitive cycles in base 5 are equal to $\sigma(n-2)$, even where that number is not prime; but there are also several cases where l is a smaller factor of $\sigma(n-2)$, for example with $n = 23$ for which $\sigma(n-2) = 6$ and both the possibilities $l = 2$ and $l = 3$ appear among the primitive cycles listed in [Appendix D](#). On the other hand, the lengths of primitive cycles in base 9 bear no relation to $\sigma(n-2)$ in most cases.

In base 3, enumerating cycles for any given n leads to complicated expressions involving double summations, whereas enumerating the integers which are members of cycles leads to closed-form expressions in many cases. The same applies in base 5, although the calculations to enumerate members of cycles are more complicated. We conjecture, on the basis of evidence from all base-5 primitive cycles with $n \leq 41$, that every primitive Kaprekar index \mathbf{k} with odd n in base 5 which satisfies conditions (19) and (20) (which are sufficient but not necessary to satisfy the inequality condition) represents a member of a primitive cycle. [In base 9 we have not been able to find conditions similar to (19) and (20) that guarantee membership of a cycle, which is why we have not attempted any enumeration of cycle members for bases higher than 5.] Then $k_0 = 1$, $k_3 = k_1$ and $k_4 \geq 2$, so to enumerate primitive cycle members we need to count the pairs of positive integers (k_1, k_2) satisfying $2k_1 + k_2 + 3 \leq n$, with k_2 even and k_1 satisfying (20). The result is given in the following theorem.

Theorem 15. *In base 5, the number of members of primitive cycles with odd digit-count $n \geq 9$ is:*

$$\begin{aligned} & ((n-5)^2 - 4)/12 \text{ members if } n \equiv 1 \pmod{6} \text{ or } n \equiv 3 \pmod{6}; \\ & (n-5)^2/12 \text{ members if } n \equiv 5 \pmod{6}. \end{aligned}$$

Proof. Given n and k_2 , there are $(n-5-k_2)/2$ integers k_1 satisfying $2k_1 \leq n-3-k_2$ with $k_1 \geq 2$. Given n , the allowed values of k_2 are the positive even integers up to $n-7$ (since $k_0 = 1$ and $k_i \geq 2$ for $i = 1, 3, 4$). So the number of pairs (k_1, k_2) satisfying the requirement $2k_1 + k_2 \leq n-3$ and condition (19) is (setting $k_2 = 2i$ and $m = (n-5)/2$)

$$\sum_{i=1}^{(n-7)/2} \frac{n-5-2i}{2} = \sum_{i=1}^{m-1} (m-i) = \frac{m(m-1)}{2} = \frac{(n-5)(n-7)}{8}. \quad (23)$$

To satisfy condition (20) we need to subtract the number of *non-compliant pairs*, those in which k_1 is odd and does not satisfy $k_1 > k_4$, so does not satisfy $3k_1 \leq n-1-k_2$; but not counting pairs with $k_1 = 1$, which have already been excluded. Now, for any integer $K \geq 3$ there are $\lfloor (K-3)/6 \rfloor$ odd integers k_1 such that $k_1 \geq 3$ and $3k_1 \leq K$ (where $\lfloor \cdot \rfloor$ is the floor function); so, given n and k_2 , there are $\lfloor (n-4-k_2)/6 \rfloor$ non-compliant pairs. Thus the number of non-compliant pairs for a given n is

$$\sum_{\text{even } k_2, 2 \leq k_2 \leq n-7} \left\lfloor \frac{n-4-k_2}{6} \right\rfloor.$$

Case (i): if $n \equiv 1 \pmod{6}$, let $n = 6t + 1$. As k_2 decreases through even integers from $n-7$ to 2, values of $(n-4-k_2)$ increase through odd integers from 3 to $6(t-1) + 1$, yielding a set of values of $\lfloor (n-4-k_2)/6 \rfloor$ including two zeroes, three of each integer from 1 to $t-2$, and one instance of the integer $t-1$. The sum of all these is

$$3 \sum_{i=1}^{t-2} i + (t-1) = 3 \frac{(t-2)(t-1)}{2} + (t-1) = \frac{(n-9)(n-7)}{24}$$

non-compliant pairs, where we have substituted $t = (n - 1)/6$ at the last step.

Case (ii): if $n \equiv 3 \pmod{6}$, let $n = 6t + 3$. As k_2 decreases through even integers from $n - 7$ to 2, values of $(n - 4 - k_2)$ increase through odd integers from 3 to $6(t - 1) + 3$, yielding a set of values of $\lfloor (n - 4 - k_2)/6 \rfloor$ including two zeroes, three of each integer from 1 to $t - 2$, and two instances of the integer $t - 1$. The sum of all these is

$$3 \sum_{i=1}^{t-2} i + 2(t-1) = 3 \frac{(t-2)(t-1)}{2} + 2(t-1) = \frac{(n-9)(n-7)}{24}$$

non-compliant pairs.

Case (iii): if $n \equiv 5 \pmod{6}$, let $n = 6t + 5$. As k_2 decreases through even integers from $n - 7$ to 2, values of $(n - 4 - k_2)$ increase through odd integers from 3 to $6(t - 1) + 5$, yielding a set of values of $\lfloor (n - 4 - k_2)/6 \rfloor$ including two zeroes and three of each integer from 1 to $t - 1$. The sum of all these is

$$3 \sum_{i=1}^{t-1} i = 3 \frac{t(t-1)}{2} = \frac{(n-11)(n-5)}{24}$$

non-compliant pairs.

The numbers of non-compliant pairs in each case above are to be subtracted from the result (23) which does not account for condition (20), and the results for the three cases are as stated in the theorem. \square

Counting the members of primitive *proper* cycles for each n in [Appendix C](#), the enumerations agree with the formulae in [Theorem 15](#) for members of *all* primitive cycles except in two cases, $n = 23$ and $n = 29$. [Theorem 15](#) includes non-proper cycles with odd multipliers c ; applying the multiplier $c = 3$ to the proper fixed point with $n = 9$ and the proper cycle of length $l = 3$ with $n = 11$, we obtain a non-proper fixed point with $n = 23$ and a non-proper cycle with $n = 29$, respectively. Obviously, non-proper cycles feature more frequently at higher digit-counts n . To count members of primitive cycles with even n , an argument similar to that used in base 3 indicates that we should replace $(n - 5)$ with $\rho(n - 2) - 3$ in the formulae in [Theorem 15](#), where ρ denotes the odd part.

To count members of all regular cycles (i.e. both primitive and general cycles) for some n , we require a summation of numbers of members of primitive cycles with any $n' \leq n$ of the same parity, similar to that in (10). As in base 3, results in closed form are only available for odd n . From [Theorem 15](#), adding the numbers of members of primitive cycles with digit-counts n , $n - 2$ and $n - 4$, for any odd $n \geq 13$, yields a total of $(n - 7)^2/4$ members with these three digit-counts. This result can then be used with standard summation formulae to yield the formulae for $N_K(n)$, the number of integers with digit-count n which are members

of a regular cycle, for any odd $n \geq 9$:

$$\begin{aligned} N_K(n) &= \frac{1}{72}(n-1)(n-4)(n-7) && \text{if } n \equiv 1 \pmod{6}; \\ N_K(n) &= \frac{1}{72}(n-3)(n^2-9n+12) && \text{if } n \equiv 3 \pmod{6}; \\ N_K(n) &= \frac{1}{72}(n-5)(n^2-7n+4) && \text{if } n \equiv 5 \pmod{6}. \end{aligned}$$

Since the distribution of special cycles is somewhat irregular, including several for odd values of n (see [Appendix B](#)), we cannot provide completely reliable formulae for the total number of cycle members for any n ; although since special cycles with odd n are comparatively scarce, the above formulae for $N_K(n)$ are accurate for this purpose in most cases where n is odd.

5 Base 7 and higher bases $b \equiv 3 \pmod{4}$

As in previous sections, we start by noting special cycles, but only in base 7, since we have not made a detailed study of the cycles that exist for any base $b > 10$; we have used data for base-7 cycles from sequences [A165080](#) (least members of cycles) and [A165081](#) (cycle lengths) in the OEIS. As in base 3, the fixed point 00 is unanimous for $n = 2$. For each digit-count $3 \leq n \leq 9$ there is a unanimous special cycle: for $n = 3$ this has length $l = 2$,

$$264 \mapsto 336 \mapsto 264;$$

for $n = 4$ it has length $l = 3$,

$$3054 \mapsto 5052 \mapsto 5232 \mapsto 3054;$$

and for $5 \leq n \leq 9$ the cycles are longer. For $n \geq 10$ there exist three classes of special cycle of length $l = 2$:

(a) For each $t \geq 1$, there is a special cycle with digit-count $n = 2t + 8$, with sequence of Kaprekar indices

$$(1, 1, t+1, 2, t, 3, 0) \mapsto (0, 1, t+1, 2, t+1, 1, 2) \mapsto (1, 1, t+1, 2, t, 3, 0); \quad (24)$$

(b) For each $t \geq 0$, there is a special cycle with digit-count $n = 5t + 10$, with sequence of Kaprekar indices

$$\begin{aligned} (1, t, t+3, 0, t+2, t+2, t+2) \mapsto & (0, t+1, 2, 2t+3, 1, t, t+3) \mapsto \\ & (1, t, t+3, 0, t+2, t+2, t+2); \end{aligned} \quad (25)$$

(c) For each $t \geq 2$, there is a special cycle with digit-count $n = 5t + 10$, with sequence of Kaprekar indices

$$\begin{aligned} (1, t+1, t+1, 2, t, t+3, t+2) \mapsto & (0, t+3, 1, 2t+2, 2, t+1, t+1) \mapsto \\ & (1, t+1, t+1, 2, t, t+3, t+2). \end{aligned} \quad (26)$$

A survey up to $n = 53$ has revealed numerous special cycles not falling into these three classes, and these are listed in [Appendix F](#); except for those with digit-counts $n = 3, 4, 10, 19$ and 51 , they all have lengths $l \geq 5$.

Turning now to regular cycles, we find that, in contrast to bases $b \equiv 1 \pmod{4}$, the regular cycles in bases $b \equiv 3 \pmod{4}$ are determined by the q -cycles arising from subgroups and cosets of \mathbb{Z}_r^\times ; but the Kaprekar indices of cycle members do not always satisfy the inequality condition, and a wider variety of further cycles can be derived from a primitive cycle than those specified by the c -multiples and s -additions of [Corollary 8](#). In more detail:

- Corresponding to each q -cycle, there exists a primitive proper base-7 cycle of digit-count $n = 3r + 2$ with members having Kaprekar indices

$$(1, q_i, r - q_i, q_i, r - q_i, q_i, r - q_i + 1) \quad (27)$$

(compare [\(7\)](#) in base 3).

- There is a further degree of generalization beyond that in formulae [\(8\)](#) and [\(9\)](#) in base 3: corresponding to each primitive proper cycle of type [\(27\)](#) with $r = (n - 2)/3$, there exist cycles of digit-count $n = 3cr + 2t + 2s$ with members having Kaprekar indices

$$(s, cq_i, c(r - q_i) + t, cq_i, c(r - q_i) + t, cq_i, c(r - q_i) + s) \quad (28)$$

for each integer $c \geq 1$, $s \geq 1$ and $t \geq 0$.

- This pattern applies for all bases $b = 4m + 3, m \geq 1$; there is a cycle of digit-count $n = (2m + 1)r + 2$ with members having Kaprekar indices

$$(1, q_i, r - q_i, \dots, q_i, r - q_i, q_i, r - q_i + 1), \quad (29)$$

from which are derived cycles of digit-count $n = (2m + 1)cr + 2t + 2s$ with members having Kaprekar indices

$$(s, cq_i, c(r - q_i) + t, \dots, cq_i, c(r - q_i) + t, cq_i, c(r - q_i) + s) \quad (30)$$

for each integer $c \geq 1$, $s \geq 1$ and $t \geq 0$, where q_i is in a q -cycle as listed in [Appendix A](#).

- There can exist further cycles derived from a cycle of type [\(29\)](#), with the same patterns of c -multiples and s -additions but different patterns of t -additions to components of its members. Several such generalizations in base 7 are noted below.

Note that if t is coprime with c , a Kaprekar index of form [\(30\)](#) may satisfy the definition of a primitive proper index, even with $c \geq 1$. Thus it is useful to introduce the following terminology in bases $b = 4m + 3, m \geq 1$.

Definition 16.

- A Kaprekar index with odd digit-count n in a base $b = 4m + 3$ is *fundamental* if it has the form

$$(1, q, r - q, \dots, q, r - q, q, r - q + 1)$$

with $r = (n - 2)/(2m + 1)$, for some even $q < r$.

- A cycle is *fundamental* if its members have fundamental Kaprekar indices.

This definition is formulated so that a Kaprekar index of the general form (30), or with any other pattern of t -additions to different components, is fundamental only if $c = 1$, $s = 1$ and $t = 0$.

As in base 3, the features described above are explained by considering succession formulae. Suppose that in base $b = 4m + 3$, a Kaprekar index \mathbf{k} satisfies the conditions

$$k_0 = 1, \quad k_j = k_1 \text{ for all odd } j, \quad k_j = k_2 \text{ for even } j \text{ with } 2 \leq j < B, \quad k_B = k_2 + 1, \quad (31)$$

where $B = b - 1$. In particular, a fundamental Kaprekar index of form (29) satisfies conditions (31) with $k_1 = q_i$ and $k_2 = r - q_i$. Since r is odd and q_i is even, $k_1 \neq k_2$ so we have two cases to consider, $k_1 < k_2$ or $k_1 > k_2$. By subtraction calculations similar to (18), we find that the successor Kaprekar index \mathbf{k}' satisfies conditions of the form (31), and that if $k_1 < k_2$, then $k'_1 = 2k_1$ and $k'_2 = k_2 - k_1$, while if $k_1 > k_2$, then $k'_1 = 2k_2$ and $k'_2 = k_1 - k_2$. With the identification $k_1 = q_i$ and $k_2 = r - q_i$, these succession formulae yield $k'_1 = q_{i+1}$ according to the q -cycle rules (4), with $k'_2 = r - q_{i+1}$. This explains the existence of fundamental cycles with values of k_1 following q -cycles. Cycles of form (30) with any positive c and s but with $t = 0$ are then explained by Theorem 7 and Corollary 8, and it remains to explain why cycles of this form with positive t exist. So we suppose that $k_0 = s$, $k_j = k_1$ for all odd j , $k_j = k_2 + t$ for even j with $2 \leq j < B$, and $k_B = k_2 + s$, and do the subtraction first in the case where $k_1 < k_2$:

$$\begin{array}{r}
 \underbrace{6 \dots 6}_{k_2+s} \underbrace{5 \dots 5}_{k_1} \underbrace{4 \dots 4}_{k_2+t} \underbrace{3 \dots 3}_{k_1} \underbrace{2 \dots 2}_{k_2+t} \underbrace{1 \dots 1}_{k_1} \underbrace{0 \dots 0}_s \\
 - \underbrace{0 \dots 0}_s \underbrace{1 \dots 1}_{k_1} \underbrace{2 \dots 2}_{k_2+t} \underbrace{3 \dots 3}_{k_1} \underbrace{4 \dots 4}_{k_2+t} \underbrace{5 \dots 5}_{k_1} \underbrace{6 \dots 6}_{k_2+s} \\
 = \underbrace{6 \dots 6}_s \underbrace{5 \dots 5}_{k_1} \underbrace{4 \dots 4}_{k_2-k_1} \underbrace{3 \dots 3}_{k_1} \underbrace{2 \dots 2}_t \underbrace{1 \dots 1}_{k_1-1} \underbrace{0}_{k_2-k_1} \underbrace{6 \dots 6}_{k_1} \underbrace{5 \dots 5}_{k_1} \underbrace{4 \dots 4}_t \underbrace{3 \dots 3}_{k_1} \underbrace{2 \dots 2}_{k_2-k_1} \underbrace{1 \dots 1}_{k_1} \underbrace{0 \dots 0}_{s-1} 1.
 \end{array} \tag{32}$$

The Kaprekar index of the result is $\mathbf{k}' = (s, 2k_1, k_2 - k_1 + t, 2k_1, k_2 - k_1 + t, 2k_1, k_2 - k_1 + s)$, which is simply the succession formula described above but with t -additions in the required places. A similar calculation in the case where $k_1 > k_2$ confirms that in all cases, addition of t to the even components of a Kaprekar index satisfying conditions (31) carries through to its successor, and so can apply throughout a regular cycle.

Unlike in bases $b \equiv 1 \pmod{4}$, it is not necessarily true that $k_j \geq 2$ for all $j \geq 1$ in Kaprekar indices of members of regular cycles in bases $b \equiv 3 \pmod{4}$. Indeed, since a q -cycle derived from a subgroup, as opposed to a coset, of \mathbb{Z}_r^\times includes the number $r - 1$, any fundamental cycle corresponding to such a q -cycle must include a member in which $k_2 = k_4 = \dots = k_{B-2} = 1$. In particular, the fundamental fixed point has Kaprekar index $(1, 2, 1, \dots, 2, 1, 2, 2)$ and digit-count $n = 6m + 5$ in base $b = 4m + 3$, and all fixed points and cycles with lesser digit-count must be special.

In base 7 we have found some classes of general cycle with t -additions not of the form (30). Derived from the fundamental cycle corresponding to the q -cycle with $r = 5$, there is a class of general cycles of length $l = 2$ with digit-count $n = 15c + 5t + 2s$:

$$(s, 2c + t, 3c + t, 2c, 3c + t, 2c + t, 3c + t + s) \mapsto (s, 4c + t, c, 4c + 2t, c, 4c + t, c + t + s) \mapsto (s, 2c + t, 3c + t, 2c, 3c + t, 2c + t, 3c + t + s),$$

in which the pattern of t -additions to components is the same as in the special cycles (25) and (26); similarly the t -additions in the special cycles of form (24) correspond to the standard pattern of t -additions in regular cycles (28). Derived from the fundamental cycle corresponding to the q -cycle with $r = 7$, there is a class of general cycles of length $l = 3$ with digit-count $n = 21c + 7t + 2s$:

$$\begin{aligned} (s, 2c + t, 5c + 2t, 2c, 5c + 2t, 2c + t, 5c + t + s) \mapsto \\ (s, 4c + t, 3c + t, 4c + 2t, 3c + t, 4c + t, 3c + t + s) \mapsto \\ (s, 6c + 2t, c, 6c + 2t, c, 6c + 2t, c + t + s) \mapsto \\ (s, 2c + t, 5c + 2t, 2c, 5c + 2t, 2c + t, 5c + t + s), \end{aligned}$$

for which we have not found a corresponding class of special cycles. Rather more anomalously, we have found a class of cycles of length $l = 3$ with digit-count $n = 19c + 8t + 2s$:

$$\begin{aligned} (s, 2c, 4c + 3t, 2c, 4c + 3t, 2c, 5c + 2t + s) \mapsto \\ (s, 3c + t, 4c + t, 2c + 2t, 4c + t, 3c + t, 3c + 2t + s) \mapsto \\ (s, 5c + 3t, c, 6c + 2t, c, 5c + 3t, c + s) \mapsto \\ (s, 2c, 4c + 3t, 2c, 4c + 3t, 2c, 5c + 2t + s), \end{aligned} \tag{33}$$

which exists only for $0 \leq t \leq c$. The fundamental cycle of this class has $n = 21$, so should be associated with the q -cycle for $r = 19$, for which $\sigma(r) = 9$; so the length of these cycles is a divisor of $\sigma(r)$, in a way reminiscent of cycles in base 5, with the progression of Kaprekar index components not tied to a q -cycle in any obvious way. Curiously, the cycle of this class with $c = 1, t = 1, s = 1$ is the fundamental cycle corresponding to the q -cycle with $r = 9$, with further cycles of the type (33) with $t = c$ being identified as higher c -multiples of the $r = 9$ fundamental cycle; but no similar identification of those cycles with $t < c$ has been found.

6 Conclusions

The paper by Yamagami and Matsui [11] was a major advance in the theory of the Kaprekar transformation. Not only did they provide a complete theory for the cycles and fixed points in base 3, but also their introduction of the Kaprekar index provided an excellent formalism for the further study of the transformation.

The principal development in the present paper is that we have shown how the structure of primitive proper, primitive non-proper and general cycles, elucidated by Yamagami and Matsui for base 3, can be generalized to all odd bases. We have classified these types of cycle together as regular cycles, and have shown that as well as satisfying the regularity condition of Definition 3, the Kaprekar indices of members of these cycles are governed by Theorems 11 and 12: the $(b-1)/2$ component must be even, and there is symmetry between the i component and the $b-1-i$ component for $1 \leq i \leq b-2$. Once these conditions are satisfied, the inequality condition of Definition 5 may come into play: it appears to be necessary for membership of a cycle in bases $b \equiv 1 \pmod{4}$, although we have not proved this. The inequality condition is satisfied trivially in base 3, but appears to be unimportant in higher bases $b \equiv 3 \pmod{4}$, where we find that there are fundamental cycles determined by subgroups generated by 2, and their cosets, in \mathbb{Z}_{n-2}^\times , the multiplicative group modulo $n-2$. In base 3 the fundamental cycles are the primitive proper cycles, from which the primitive non-proper and general cycles are derived, but in higher bases there is a greater variety of cycles that may be derived from a fundamental cycle. All cycles derived from a fundamental cycle have the same length; if 2 is a generator of the group \mathbb{Z}_{n-2}^\times and $n-2$ is prime, then the length of these cycles is $l = (n-3)/2$. Thus if Artin's Conjecture on primitive roots is true, in particular if 2 is a primitive root modulo infinitely many primes, then there exist arbitrarily long cycles in bases $b \equiv 3 \pmod{4}$. Myers [7] has conjectured that there exist arbitrarily long cycles in any odd base, which remains unproved.

We describe all cycles that do not fit within our classification of regular cycles as special. Special cycles become more prevalent in higher bases. Whereas many of them do not fit within any clear pattern, there are also some infinite classes of special cycle: these include the singular cycles noted by Yamagami and Matsui in base 3, which also appear in a more complicated form in base 5, and which yield a cycle of every possible length $l \in \mathbb{N}$ in these two bases. We have also found several classes of special fixed points and cycles of specific length in bases 7 and 9, but we have not been able to develop any general theory for classes of special cycles.

Overall, there is a picture of increasing complexity as the base b increases. We have established principles that govern the regular cycles in bases $b \equiv 1 \pmod{4}$ and $b \equiv 3 \pmod{4}$, but there are ever more variations on these basic principles in higher bases. There is also a plethora of special cycles, taking forms which are individual to each base.

7 Acknowledgments

This work was begun by KD-W as a final-year undergraduate project under the supervision of AK at Loughborough University, and continued by AK after his retirement. The authors thank Ben Crossley for useful discussions. The work could not have been done without the computational results made freely available by Joseph Myers in the OEIS.

Appendix A: Table of q -cycles

r	$\phi(r)$	$\sigma(r)$	$N(r)$	q -cycle(s)
3	2	1	1	$2 \mapsto 2$
5	4	2	1	$2 \mapsto 4 \mapsto 2$
7	6	3	1	$2 \mapsto 4 \mapsto 6 \mapsto 2$
9	6	3	1	$2 \mapsto 4 \mapsto 8 \mapsto 2$
11	10	5	1	$2 \mapsto 4 \mapsto 8 \mapsto 6 \mapsto 10 \mapsto 2$
13	12	6	1	$2 \mapsto 4 \mapsto 8 \mapsto 10 \mapsto 6 \mapsto 12 \mapsto 2$
15	8	4	1	$2 \mapsto 4 \mapsto 8 \mapsto 14 \mapsto 2$
17	16	4	2	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 2,$ $6 \mapsto 12 \mapsto 10 \mapsto 14 \mapsto 6$
19	18	9	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 6 \mapsto 12 \mapsto 14 \mapsto 10 \mapsto 18 \mapsto 2$
21	12	6	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 10 \mapsto 20 \mapsto 2$
23	22	11	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 14 \mapsto 18 \mapsto 10 \mapsto 20 \mapsto 6 \mapsto 12 \mapsto 22 \mapsto 2$
25	20	10	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 18 \mapsto 14 \mapsto 22 \mapsto 6 \mapsto 12 \mapsto 24 \mapsto 2$
27	18	9	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 22 \mapsto 10 \mapsto 20 \mapsto 14 \mapsto 26 \mapsto 2$
29	28	14	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 26 \mapsto 6 \mapsto 12 \mapsto 24 \mapsto 10 \mapsto 20 \mapsto 18 \mapsto 22 \mapsto$ $14 \mapsto 28 \mapsto 2$
31	30	5	3	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 30 \mapsto 2,$ $6 \mapsto 12 \mapsto 24 \mapsto 14 \mapsto 28 \mapsto 6,$ $10 \mapsto 20 \mapsto 22 \mapsto 18 \mapsto 26 \mapsto 10$
33	20	5	2	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 2,$ $10 \mapsto 20 \mapsto 26 \mapsto 14 \mapsto 28 \mapsto 10$
35	24	12	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 6 \mapsto 12 \mapsto 24 \mapsto 22 \mapsto 26 \mapsto 18 \mapsto 34 \mapsto 2$
37	36	18	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 10 \mapsto 20 \mapsto 34 \mapsto 6 \mapsto 12 \mapsto 24 \mapsto 26 \mapsto$ $22 \mapsto 30 \mapsto 14 \mapsto 28 \mapsto 18 \mapsto 36 \mapsto 2$
39	24	12	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 14 \mapsto 28 \mapsto 22 \mapsto 34 \mapsto 10 \mapsto 20 \mapsto 38 \mapsto 2$
41	40	10	2	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 18 \mapsto 36 \mapsto 10 \mapsto 20 \mapsto 40 \mapsto 2,$ $6 \mapsto 12 \mapsto 24 \mapsto 34 \mapsto 14 \mapsto 28 \mapsto 26 \mapsto 30 \mapsto 22 \mapsto 38 \mapsto 6$
43	42	7	3	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 22 \mapsto 42 \mapsto 2,$ $6 \mapsto 12 \mapsto 24 \mapsto 38 \mapsto 10 \mapsto 20 \mapsto 40 \mapsto 6,$ $14 \mapsto 28 \mapsto 30 \mapsto 26 \mapsto 34 \mapsto 18 \mapsto 36 \mapsto 14$
45	24	12	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 26 \mapsto 38 \mapsto 14 \mapsto 28 \mapsto 34 \mapsto 22 \mapsto 44 \mapsto 2$
47	46	23	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 30 \mapsto 34 \mapsto 26 \mapsto 42 \mapsto 10 \mapsto 20 \mapsto 40 \mapsto$ $14 \mapsto 28 \mapsto 38 \mapsto 18 \mapsto 36 \mapsto 22 \mapsto 44 \mapsto 6 \mapsto 12 \mapsto 24 \mapsto 46 \mapsto 2$
49	42	21	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 34 \mapsto 30 \mapsto 38 \mapsto 22 \mapsto 44 \mapsto 10 \mapsto 20 \mapsto$ $40 \mapsto 18 \mapsto 36 \mapsto 26 \mapsto 46 \mapsto 6 \mapsto 12 \mapsto 24 \mapsto 48 \mapsto 2$
51	32	8	2	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 38 \mapsto 26 \mapsto 50 \mapsto 2,$ $10 \mapsto 20 \mapsto 40 \mapsto 22 \mapsto 44 \mapsto 14 \mapsto 28 \mapsto 46 \mapsto 10$

r	$\phi(r)$	$\sigma(r)$	$N(r)$	q -cycle(s)
53	52	26	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 42 \mapsto 22 \mapsto 44 \mapsto 18 \mapsto 36 \mapsto 34 \mapsto 38 \mapsto 30 \mapsto 46 \mapsto 14 \mapsto 28 \mapsto 50 \mapsto 6 \mapsto 12 \mapsto 24 \mapsto 48 \mapsto 10 \mapsto 20 \mapsto 40 \mapsto 26 \mapsto 52 \mapsto 2$
55	40	20	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 46 \mapsto 18 \mapsto 36 \mapsto 38 \mapsto 34 \mapsto 42 \mapsto 26 \mapsto 52 \mapsto 6 \mapsto 12 \mapsto 24 \mapsto 48 \mapsto 14 \mapsto 28 \mapsto 54 \mapsto 2$
57	36	9	2	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 50 \mapsto 14 \mapsto 28 \mapsto 56 \mapsto 2, 10 \mapsto 20 \mapsto 40 \mapsto 34 \mapsto 46 \mapsto 22 \mapsto 44 \mapsto 26 \mapsto 52 \mapsto 10$
59	58	29	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 54 \mapsto 10 \mapsto 20 \mapsto 40 \mapsto 38 \mapsto 42 \mapsto 34 \mapsto 50 \mapsto 18 \mapsto 36 \mapsto 46 \mapsto 26 \mapsto 52 \mapsto 14 \mapsto 28 \mapsto 56 \mapsto 6 \mapsto 12 \mapsto 24 \mapsto 48 \mapsto 22 \mapsto 44 \mapsto 30 \mapsto 58 \mapsto 2$
61	60	30	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 58 \mapsto 6 \mapsto 12 \mapsto 24 \mapsto 48 \mapsto 26 \mapsto 52 \mapsto 18 \mapsto 36 \mapsto 50 \mapsto 22 \mapsto 44 \mapsto 34 \mapsto 54 \mapsto 14 \mapsto 28 \mapsto 56 \mapsto 10 \mapsto 20 \mapsto 40 \mapsto 42 \mapsto 38 \mapsto 46 \mapsto 30 \mapsto 60 \mapsto 2$
63	36	6	3	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 62 \mapsto 2, 10 \mapsto 20 \mapsto 40 \mapsto 46 \mapsto 34 \mapsto 58 \mapsto 10, 22 \mapsto 44 \mapsto 38 \mapsto 50 \mapsto 26 \mapsto 52 \mapsto 22$
65	48	6	4	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 64 \mapsto 2, 6 \mapsto 12 \mapsto 24 \mapsto 48 \mapsto 34 \mapsto 62 \mapsto 6, 14 \mapsto 28 \mapsto 56 \mapsto 18 \mapsto 36 \mapsto 58 \mapsto 14, 22 \mapsto 44 \mapsto 42 \mapsto 46 \mapsto 38 \mapsto 54 \mapsto 22$
67	66	33	1	$2 \mapsto 4 \mapsto 8 \mapsto 16 \mapsto 32 \mapsto 64 \mapsto 6 \mapsto 12 \mapsto 24 \mapsto 48 \mapsto 38 \mapsto 58 \mapsto 18 \mapsto 36 \mapsto 62 \mapsto 10 \mapsto 20 \mapsto 40 \mapsto 54 \mapsto 26 \mapsto 52 \mapsto 30 \mapsto 60 \mapsto 14 \mapsto 28 \mapsto 56 \mapsto 22 \mapsto 44 \mapsto 46 \mapsto 42 \mapsto 50 \mapsto 34 \mapsto 66 \mapsto 2$

Table 1: For odd r up to 67, the Euler totient $\phi(r)$, the length $\sigma(r)$ and number $N(r)$ of q -cycles is given, together with the q -cycles required to calculate base-3 Kaprekar cycles. Whereas in the main text we have taken q_0 as the largest member of a q -cycle, here we start the cycles with the smallest member.

Appendix B: Special cycles in base 5

$$n = 2, l = 1 : (0, 1, 0, 1, 0) \mapsto (0, 1, 0, 1, 0)$$

$$n = 2, l = 1 : (2, 0, 0, 0, 0) \mapsto (2, 0, 0, 0, 0)$$

[repdigit from non-trivial initial integer]

$$n = 3, l = 2 : (0, 1, 0, 1, 1) \mapsto (0, 0, 2, 0, 1) \mapsto (0, 1, 0, 1, 1)$$

$$n = 4, l = 1 : (1, 0, 1, 2, 0) \mapsto (1, 0, 1, 2, 0)$$

$$n = 5, l = 4 : (1, 0, 1, 2, 1) \mapsto (1, 1, 0, 1, 2) \mapsto (0, 2, 1, 0, 2) \mapsto (0, 1, 2, 1, 1) \mapsto (1, 0, 1, 2, 1)$$

$$n = 6, l = 5 : (1, 0, 1, 2, 2) \mapsto (0, 2, 1, 0, 3) \mapsto (0, 2, 2, 2, 0) \mapsto (0, 1, 2, 1, 2) \mapsto (1, 0, 3, 2, 0) \mapsto (1, 0, 1, 2, 2)$$

$n = 7, l = 4 : (1, 0, 3, 2, 1) \mapsto (1, 2, 0, 2, 2) \mapsto (0, 3, 1, 1, 2) \mapsto (0, 2, 2, 2, 1) \mapsto (1, 0, 3, 2, 1)$
 $n = 8, l = 6 : (1, 0, 1, 2, 4) \mapsto (1, 2, 2, 2, 1) \mapsto (0, 2, 3, 0, 3) \mapsto (0, 2, 2, 2, 2) \mapsto (1, 2, 1, 4, 0) \mapsto (1, 0, 5, 2, 0) \mapsto (1, 0, 1, 2, 4)$
 $n = 10, l = 4 : (1, 0, 1, 2, 6) \mapsto (1, 2, 2, 2, 3) \mapsto (1, 4, 0, 4, 1) \mapsto (0, 2, 7, 0, 1) \mapsto (1, 0, 1, 2, 6)$
 $n = 12, l = 8 : (1, 0, 5, 2, 4) \mapsto (1, 2, 6, 2, 1) \mapsto (0, 2, 3, 0, 7) \mapsto (0, 2, 6, 2, 2) \mapsto (1, 2, 1, 4, 4) \mapsto (0, 4, 1, 2, 5) \mapsto (0, 4, 2, 4, 2) \mapsto (1, 2, 5, 4, 0) \mapsto (1, 0, 5, 2, 4)$
 $n = 14, l = 4 : (1, 4, 1, 6, 2) \mapsto (1, 2, 6, 2, 3) \mapsto (1, 4, 0, 4, 5) \mapsto (0, 5, 1, 3, 5) \mapsto (1, 4, 1, 6, 2)$
 $n = 14, l = 5 : (1, 2, 5, 4, 2) \mapsto (1, 4, 2, 4, 3) \mapsto (1, 4, 4, 4, 1) \mapsto (0, 2, 7, 0, 5) \mapsto (0, 2, 6, 2, 4) \mapsto (1, 2, 5, 4, 2)$
 $n = 15, l = 4 : (0, 2, 2, 2, 9) \mapsto (1, 2, 5, 4, 3) \mapsto (1, 6, 0, 6, 2) \mapsto (0, 3, 9, 1, 2) \mapsto (0, 2, 2, 2, 9)$
 $n = 16, l = 4 : (0, 2, 2, 2, 10) \mapsto (1, 2, 5, 4, 4) \mapsto (1, 6, 2, 6, 1) \mapsto (0, 2, 11, 0, 3) \mapsto (0, 2, 2, 2, 10)$
 $n = 16, l = 5, : (1, 2, 9, 4, 0) \mapsto (1, 0, 5, 2, 8) \mapsto (1, 2, 10, 2, 1) \mapsto (0, 2, 3, 0, 11) \mapsto (0, 2, 6, 2, 6) \mapsto (1, 2, 9, 4, 0)$
 $n = 18, l = 5 : (1, 0, 1, 2, 14) \mapsto (1, 2, 2, 2, 11) \mapsto (1, 4, 4, 4, 5) \mapsto (1, 8, 0, 8, 1) \mapsto (0, 2, 15, 0, 1) \mapsto (1, 0, 1, 2, 14)$
 $n = 19, l = 3 : (0, 5, 7, 3, 4) \mapsto (1, 4, 3, 6, 5) \mapsto (1, 7, 0, 7, 4) \mapsto (0, 5, 7, 3, 4)$
 $n = 22, l = 5 : (1, 6, 3, 8, 4) \mapsto (1, 6, 6, 6, 3) \mapsto (1, 4, 8, 4, 5) \mapsto (1, 8, 0, 8, 5) \mapsto (0, 6, 7, 4, 5) \mapsto (1, 6, 3, 8, 4)$
 $n = 22, l = 5 : (0, 2, 6, 2, 12) \mapsto (1, 2, 13, 4, 2) \mapsto (1, 4, 2, 4, 11) \mapsto (1, 8, 4, 8, 1) \mapsto (0, 2, 15, 0, 5) \mapsto (0, 2, 6, 2, 12)$
 $n = 24, l = 11 : (1, 2, 1, 4, 16) \mapsto (1, 6, 2, 6, 9) \mapsto (0, 8, 3, 6, 7) \mapsto (1, 8, 3, 10, 2) \mapsto (1, 4, 14, 4, 1) \mapsto (0, 2, 7, 0, 15) \mapsto (0, 2, 14, 2, 6) \mapsto (1, 2, 9, 4, 8) \mapsto (1, 6, 10, 6, 1) \mapsto (0, 2, 11, 0, 11) \mapsto (0, 2, 18, 2, 2) \mapsto (1, 2, 1, 4, 16)$
 $n = 26, l = 10 : (1, 0, 1, 2, 22) \mapsto (1, 2, 2, 2, 19) \mapsto (1, 4, 4, 4, 13) \mapsto (1, 8, 8, 8, 1) \mapsto (0, 2, 15, 0, 9) \mapsto (0, 2, 14, 2, 8) \mapsto (1, 2, 13, 4, 6) \mapsto (1, 6, 6, 6, 7) \mapsto (1, 12, 0, 12, 1) \mapsto (0, 2, 23, 0, 1) \mapsto (1, 0, 1, 2, 22)$
 $n = 30, l = 9 : (0, 2, 6, 2, 20) \mapsto (1, 2, 13, 4, 10) \mapsto (1, 6, 14, 6, 3) \mapsto (1, 4, 8, 4, 13) \mapsto (0, 6, 15, 4, 5) \mapsto (1, 6, 3, 8, 12) \mapsto (1, 8, 6, 8, 7) \mapsto (1, 12, 4, 12, 1) \mapsto (0, 2, 23, 0, 5) \mapsto (0, 2, 6, 2, 20)$
 $n = 32, l = 4 : (0, 2, 22, 2, 6) \mapsto (1, 2, 9, 4, 16) \mapsto (1, 6, 18, 6, 1) \mapsto (0, 2, 11, 0, 19) \mapsto (0, 2, 22, 2, 6)$
 $n = 33, l = 4 : (1, 13, 0, 13, 6) \mapsto (0, 7, 15, 5, 6) \mapsto (1, 8, 3, 10, 11) \mapsto (1, 9, 4, 9, 10) \mapsto (1, 13, 0, 13, 6)$
 $n = 34, l = 6 : (1, 0, 1, 2, 30) \mapsto (1, 2, 2, 2, 27) \mapsto (1, 4, 4, 4, 21) \mapsto (1, 8, 8, 8, 9) \mapsto (1, 16, 0, 16, 1) \mapsto (0, 2, 31, 0, 1) \mapsto (1, 0, 1, 2, 30)$
 $n = 38, l = 6 : (1, 6, 3, 8, 20) \mapsto (1, 14, 6, 14, 3) \mapsto (1, 4, 24, 4, 5) \mapsto (1, 8, 0, 8, 21) \mapsto$

$$\begin{aligned}
& (1, 16, 0, 16, 5) \mapsto (0, 6, 23, 4, 5) \mapsto (1, 6, 3, 8, 20) \\
n = 42, l = 12 : & (0, 2, 14, 2, 24) \mapsto (1, 2, 29, 4, 6) \mapsto (1, 6, 6, 6, 23) \mapsto \\
& (1, 12, 12, 12, 5) \mapsto (1, 8, 16, 8, 9) \mapsto (1, 16, 0, 16, 9) \mapsto \\
& (0, 10, 15, 8, 9) \mapsto (1, 14, 3, 16, 8) \mapsto (1, 10, 14, 10, 7) \mapsto \\
& (1, 12, 8, 12, 9) \mapsto (1, 16, 8, 16, 1) \mapsto (0, 2, 31, 0, 9) \mapsto (0, 2, 14, 2, 24) \\
n = 46, l = 5 : & (1, 2, 5, 4, 34) \mapsto (1, 6, 10, 6, 23) \mapsto (1, 12, 20, 12, 1) \mapsto \\
& (0, 2, 23, 0, 21) \mapsto (0, 2, 38, 2, 4) \mapsto (1, 2, 5, 4, 34) \\
n = 47, l = 4 : & (1, 19, 0, 19, 8) \mapsto (0, 9, 23, 7, 8) \mapsto (1, 12, 3, 14, 17) \mapsto \\
& (1, 13, 6, 13, 14) \mapsto (1, 19, 0, 19, 8) \\
n = 50, l = 7 : & (1, 2, 29, 4, 14) \mapsto (1, 6, 22, 6, 15) \mapsto (1, 12, 16, 12, 9) \mapsto \\
& (1, 16, 8, 16, 9) \mapsto (1, 16, 16, 16, 1) \mapsto (0, 2, 31, 0, 17) \mapsto \\
& (0, 2, 30, 2, 16) \mapsto (1, 2, 29, 4, 14)
\end{aligned}$$

Appendix C: Succession formulae in base 5

We list Kaprekar indices of successors \mathbf{k}' to indices \mathbf{k} , where $k_0 = 1$ and the remaining components satisfy the given conditions in each case. The cases cover all possibilities where $k_0 = 1$ and $k_i \geq 1$ for $i = 1, 2, 3, 4$.

- (i) If $k_4 + k_3 + k_2 < 1 + k_1$, then
$$\mathbf{k}' = (1, k_2 + k_4 - 1, 2k_3, k_2 + k_4 - 1, k_1 - k_2 - k_3 - k_4 + 2)$$
- (ii) If $k_4 + k_3 + k_2 = 1 + k_1$, then
$$\mathbf{k}' = (1, k_2 + k_4 - 1, 2k_3, k_2 + k_4 - 1, 1)$$
- (iii) If $k_4 + k_3 < 1 + k_1 < k_4 + k_3 + k_2$, then
$$\mathbf{k}' = (1, k_1 - k_3, 2k_3, k_1 - k_3, k_2 + k_3 + k_4 - k_1)$$
- (iv) If $k_4 + k_3 = 1 + k_1$, then
$$\mathbf{k}' = (0, k_4 + 1, 2k_3 - 1, k_4 - 1, k_2 + 1)$$
- (v) If $k_4 < 1 + k_1 < k_4 + k_3 < 1 + k_1 + k_2$, then
$$\mathbf{k}' = (1, 2k_4 + k_3 - k_1 - 2, 2k_1 - 2k_4 + 2, 2k_4 + k_3 - k_1 - 2, k_1 + k_2 - k_3 - k_4 + 2)$$
- (vi) If $k_4 < 1 + k_1$ and $k_4 + k_3 = 1 + k_1 + k_2$, then
$$\mathbf{k}' = (1, k_4 + k_2 - 1, 2k_1 - 2k_4 + 2, k_4 + k_2 - 1, 1)$$
- (vii) If $k_4 < 1 + k_1$ and $k_4 + k_3 > 1 + k_1 + k_2$, then
$$\mathbf{k}' = (1, k_4 + k_2 - 1, 2k_1 - 2k_4 + 2, k_4 + k_2 - 1, k_4 + k_3 - k_2 - k_1)$$
- (viii) If $k_4 = 1 + k_1$ and $k_3 < k_2$, then
$$\mathbf{k}' = (1, k_1 + k_3, 0, k_1 + k_3, k_2 - k_3 + 1)$$
- (ix) If $k_4 = 1 + k_1$ and $k_2 < k_3$, then
$$\mathbf{k}' = (1, k_1 + k_2, 0, k_1 + k_2, k_3 - k_2 + 1)$$
- (x) If $1 + k_1 < k_4$ and $k_4 + k_3 < 1 + k_1 + k_2$, then
$$\mathbf{k}' = (1, k_1 + k_3, 2k_4 - 2k_1 - 2, k_1 + k_3, k_1 + k_2 - k_3 - k_4 + 2)$$
- (xi) If $1 + k_1 < k_4$ and $k_4 + k_3 = 1 + k_1 + k_2$, then
$$\mathbf{k}' = (1, k_1 + k_3, 2k_4 - 2k_1 - 2, k_1 + k_3, 1)$$

- (xii) If $1 + k_1 < k_4 < 1 + k_1 + k_2 < k_4 + k_3$, then
 $\mathbf{k}' = (1, 2k_1 + k_2 - k_4 + 1, 2k_4 - 2k_1 - 2, 2k_1 + k_2 - k_4 + 1, k_4 + k_3 - k_2 - k_1)$
- (xiii) If $1 + k_1 + k_2 = k_4$, then
 $\mathbf{k}' = (0, k_1 + 2, 2k_2 - 1, k_1, k_3 + 1)$
- (xiv) If $1 + k_1 + k_2 < k_4 < 1 + k_1 + k_2 + k_3$, then
 $\mathbf{k}' = (1, k_4 - k_2 - 1, 2k_2, k_4 - k_2 - 1, k_3 + k_2 + k_1 - k_4 + 2)$
- (xv) If $1 + k_1 + k_2 + k_3 = k_4$, then
 $\mathbf{k}' = (1, k_1 + k_3, 2k_2, k_1 + k_3, 1)$
- (xvi) If $1 + k_1 + k_2 + k_3 < k_4$, then
 $\mathbf{k}' = (1, k_1 + k_3, 2k_2, k_1 + k_3, k_4 - k_3 - k_2 - k_1)$

Appendix D: Primitive proper cycles in base 5

The Kaprekar indices of members of primitive cycles take the form

$$(k_0, k_1, k_2, k_3, k_4) = (1, k_1, k_2, k_1, n - 2k_1 - k_2 - 1),$$

with only two free parameters; so we display cycles of pairs of values (k_1, k_2) as shorthand for the full Kaprekar indices.

- $n = 9, l = 1$: $(2, 2) \mapsto (2, 2)$
- $n = 11, l = 3$: $(2, 2) \mapsto (3, 2) \mapsto (2, 4) \mapsto (2, 2)$
- $n = 13, l = 5$: $(2, 2) \mapsto (3, 4) \mapsto (2, 4) \mapsto (4, 2) \mapsto (2, 6) \mapsto (2, 2)$
- $n = 15, l = 2$: $(2, 4) \mapsto (3, 6) \mapsto (2, 4)$
- $n = 15, l = 6$: $(2, 2) \mapsto (4, 4) \mapsto (2, 6) \mapsto (4, 2) \mapsto (5, 2) \mapsto (2, 8) \mapsto (2, 2)$
- $n = 17, l = 2$: $(2, 6) \mapsto (4, 6) \mapsto (2, 6)$
- $n = 17, l = 2$: $(2, 4) \mapsto (3, 8) \mapsto (2, 4)$
- $n = 17, l = 4$: $(2, 2) \mapsto (4, 4) \mapsto (6, 2) \mapsto (2, 10) \mapsto (2, 2)$
- $n = 17, l = 4$: $(2, 8) \mapsto (4, 2) \mapsto (5, 2) \mapsto (5, 4) \mapsto (2, 8)$
- $n = 19, l = 4$: $(2, 2) \mapsto (4, 4) \mapsto (7, 2) \mapsto (2, 12) \mapsto (2, 2)$
- $n = 19, l = 4$: $(2, 4) \mapsto (4, 8) \mapsto (2, 6) \mapsto (3, 10) \mapsto (2, 4)$
- $n = 19, l = 4$: $(2, 8) \mapsto (4, 6) \mapsto (6, 2) \mapsto (5, 6) \mapsto (2, 8)$
- $n = 19, l = 4$: $(2, 10) \mapsto (4, 2) \mapsto (5, 4) \mapsto (6, 4) \mapsto (2, 10)$
- $n = 21, l = 3$: $(2, 12) \mapsto (4, 2) \mapsto (7, 4) \mapsto (2, 12)$
- $n = 21, l = 9$: $(2, 2) \mapsto (4, 4) \mapsto (5, 6) \mapsto (6, 4) \mapsto (6, 6) \mapsto (2, 10) \mapsto (4, 6) \mapsto (8, 2) \mapsto (2, 14) \mapsto (2, 2)$
- $n = 21, l = 9$: $(2, 4) \mapsto (4, 8) \mapsto (6, 2) \mapsto (7, 2) \mapsto (5, 8) \mapsto (2, 8) \mapsto (4, 10) \mapsto (2, 6) \mapsto (3, 12) \mapsto (2, 4)$
- $n = 23, l = 2$: $(2, 6) \mapsto (4, 12) \mapsto (2, 6)$
- $n = 23, l = 3$: $(2, 14) \mapsto (4, 2) \mapsto (8, 4) \mapsto (2, 14)$
- $n = 23, l = 3$: $(2, 12) \mapsto (4, 6) \mapsto (7, 6) \mapsto (2, 12)$
- $n = 23, l = 6$: $(2, 2) \mapsto (4, 4) \mapsto (5, 8) \mapsto (6, 4) \mapsto (9, 2) \mapsto (2, 16) \mapsto (2, 2)$

$n = 23, l = 6 : (2, 4) \mapsto (4, 8) \mapsto (8, 2) \mapsto (5, 10) \mapsto (2, 8) \mapsto (3, 14) \mapsto (2, 4)$
 $n = 23, l = 6 : (2, 10) \mapsto (4, 10) \mapsto (6, 2) \mapsto (7, 2) \mapsto (7, 4) \mapsto (6, 8) \mapsto (2, 10)$
 $n = 25, l = 11 : (2, 2) \mapsto (4, 4) \mapsto (7, 8) \mapsto (2, 12) \mapsto (4, 10) \mapsto (8, 2) \mapsto (7, 6) \mapsto$
 $(6, 8) \mapsto (6, 6) \mapsto (10, 2) \mapsto (2, 18) \mapsto (2, 2)$
 $n = 25, l = 11 : (2, 4) \mapsto (4, 8) \mapsto (8, 6) \mapsto (2, 14) \mapsto (4, 6) \mapsto (5, 10) \mapsto (6, 4) \mapsto$
 $(9, 2) \mapsto (5, 12) \mapsto (2, 8) \mapsto (3, 16) \mapsto (2, 4)$
 $n = 25, l = 11 : (2, 6) \mapsto (4, 12) \mapsto (6, 2) \mapsto (7, 4) \mapsto (9, 4) \mapsto (2, 16) \mapsto (4, 2) \mapsto$
 $(8, 4) \mapsto (6, 10) \mapsto (2, 10) \mapsto (4, 14) \mapsto (2, 6)$
 $n = 27, l = 10 : (2, 2) \mapsto (4, 4) \mapsto (8, 8) \mapsto (2, 14) \mapsto (4, 10) \mapsto (8, 6) \mapsto (6, 10) \mapsto$
 $(6, 6) \mapsto (11, 2) \mapsto (2, 20) \mapsto (2, 2)$
 $n = 27, l = 10 : (2, 4) \mapsto (4, 8) \mapsto (7, 10) \mapsto (2, 12) \mapsto (4, 14) \mapsto (6, 2) \mapsto (9, 4) \mapsto$
 $(6, 12) \mapsto (2, 10) \mapsto (3, 18) \mapsto (2, 4)$
 $n = 27, l = 10 : (2, 6) \mapsto (4, 12) \mapsto (8, 2) \mapsto (9, 2) \mapsto (7, 8) \mapsto (6, 8) \mapsto (10, 2) \mapsto$
 $(5, 14) \mapsto (2, 8) \mapsto (4, 16) \mapsto (2, 6)$
 $n = 27, l = 10 : (2, 16) \mapsto (4, 6) \mapsto (5, 12) \mapsto (6, 4) \mapsto (7, 6) \mapsto (10, 4) \mapsto (2, 18) \mapsto$
 $(4, 2) \mapsto (8, 4) \mapsto (9, 6) \mapsto (2, 16)$
 $n = 29, l = 9 : (2, 2) \mapsto (4, 4) \mapsto (8, 8) \mapsto (6, 10) \mapsto (10, 2) \mapsto (7, 10) \mapsto (6, 8) \mapsto$
 $(12, 2) \mapsto (2, 22) \mapsto (2, 2)$
 $n = 29, l = 9 : (2, 4) \mapsto (4, 8) \mapsto (5, 14) \mapsto (6, 4) \mapsto (7, 8) \mapsto (10, 4) \mapsto (6, 14) \mapsto$
 $(2, 10) \mapsto (3, 20) \mapsto (2, 4)$
 $n = 29, l = 9 : (2, 6) \mapsto (4, 12) \mapsto (8, 6) \mapsto (10, 6) \mapsto (2, 18) \mapsto (4, 6) \mapsto (7, 12) \mapsto$
 $(2, 12) \mapsto (4, 18) \mapsto (2, 6)$
 $n = 29, l = 9 : (2, 8) \mapsto (4, 16) \mapsto (6, 2) \mapsto (11, 4) \mapsto (2, 20) \mapsto (4, 2) \mapsto (8, 4) \mapsto$
 $(11, 2) \mapsto (5, 16) \mapsto (2, 8)$
 $n = 29, l = 9 : (2, 14) \mapsto (4, 14) \mapsto (8, 2) \mapsto (9, 2) \mapsto (9, 4) \mapsto (9, 8) \mapsto (2, 16) \mapsto$
 $(4, 10) \mapsto (8, 10) \mapsto (2, 14)$
 $n = 31, l = 14 : (2, 2) \mapsto (4, 4) \mapsto (8, 8) \mapsto (10, 6) \mapsto (6, 14) \mapsto (6, 6) \mapsto (7, 10) \mapsto$
 $(10, 4) \mapsto (9, 10) \mapsto (2, 16) \mapsto (4, 14) \mapsto (8, 6) \mapsto (13, 2) \mapsto$
 $(2, 24) \mapsto (2, 2)$
 $n = 31, l = 14 : (2, 4) \mapsto (4, 8) \mapsto (5, 16) \mapsto (6, 4) \mapsto (9, 8) \mapsto (6, 12) \mapsto (10, 2) \mapsto$
 $(9, 6) \mapsto (10, 8) \mapsto (2, 18) \mapsto (4, 10) \mapsto (7, 14) \mapsto (2, 12) \mapsto$
 $(3, 22) \mapsto (2, 4)$
 $n = 31, l = 14 : (2, 6) \mapsto (4, 12) \mapsto (8, 10) \mapsto (6, 10) \mapsto (12, 2) \mapsto (5, 18) \mapsto (2, 8) \mapsto$
 $(4, 16) \mapsto (8, 2) \mapsto (9, 4) \mapsto (11, 4) \mapsto (6, 16) \mapsto (2, 10) \mapsto$
 $(4, 20) \mapsto (2, 6)$
 $n = 31, l = 14 : (2, 14) \mapsto (4, 18) \mapsto (6, 2) \mapsto (12, 4) \mapsto (2, 22) \mapsto (4, 2) \mapsto (8, 4) \mapsto$
 $(11, 2) \mapsto (7, 12) \mapsto (6, 8) \mapsto (11, 6) \mapsto (2, 20) \mapsto (4, 6) \mapsto$
 $(8, 12) \mapsto (2, 14)$
 $n = 33, l = 5 : (2, 2) \mapsto (4, 4) \mapsto (8, 8) \mapsto (14, 2) \mapsto (2, 26) \mapsto (2, 2)$
 $n = 33, l = 5 : (2, 4) \mapsto (4, 8) \mapsto (7, 16) \mapsto (2, 12) \mapsto (3, 24) \mapsto (2, 4)$
 $n = 33, l = 5 : (2, 6) \mapsto (4, 12) \mapsto (8, 14) \mapsto (2, 14) \mapsto (4, 22) \mapsto (2, 6)$
 $n = 33, l = 5 : (2, 8) \mapsto (4, 16) \mapsto (8, 6) \mapsto (13, 2) \mapsto (5, 20) \mapsto (2, 8)$

$$\begin{aligned}
n = 33, l = 5 : & \quad (2, 10) \mapsto (4, 20) \mapsto (6, 2) \mapsto (12, 4) \mapsto (6, 18) \mapsto (2, 10) \\
n = 33, l = 5 : & \quad (2, 16) \mapsto (4, 18) \mapsto (8, 2) \mapsto (11, 4) \mapsto (9, 12) \mapsto (2, 16) \\
n = 33, l = 5 : & \quad (2, 18) \mapsto (4, 14) \mapsto (8, 10) \mapsto (10, 6) \mapsto (10, 10) \mapsto (2, 18) \\
n = 33, l = 5 : & \quad (2, 20) \mapsto (4, 10) \mapsto (5, 18) \mapsto (6, 4) \mapsto (11, 8) \mapsto (2, 20) \\
n = 33, l = 5 : & \quad (2, 22) \mapsto (4, 6) \mapsto (8, 12) \mapsto (6, 10) \mapsto (12, 6) \mapsto (2, 22) \\
n = 33, l = 5 : & \quad (2, 24) \mapsto (4, 2) \mapsto (8, 4) \mapsto (9, 6) \mapsto (13, 4) \mapsto (2, 24) \\
n = 33, l = 5 : & \quad (6, 6) \mapsto (7, 12) \mapsto (10, 4) \mapsto (11, 6) \mapsto (6, 16) \mapsto (6, 6) \\
n = 33, l = 5 : & \quad (6, 8) \mapsto (9, 10) \mapsto (6, 12) \mapsto (12, 2) \mapsto (7, 14) \mapsto (6, 8) \\
n = 33, l = 5 : & \quad (6, 14) \mapsto (10, 2) \mapsto (11, 2) \mapsto (9, 8) \mapsto (10, 8) \mapsto (6, 14)
\end{aligned}$$

Appendix E: Primitive proper cycles in base 9

The Kaprekar indices of members of primitive cycles have $k_0 = 1, k_5 = k_3, k_6 = k_2, k_7 = k_1$ and $k_8 = n - 2(k_1 + k_2 + k_3) - k_4 - 1$, so there are only four free parameters; thus we display cycles of quadruplets of values (k_1, k_2, k_3, k_4) as shorthand for the full Kaprekar indices.

$$\begin{aligned}
n = 17, l = 1 : & \quad (2, 2, 2, 2) \mapsto (2, 2, 2, 2) \\
n = 19, l = 4 : & \quad (2, 2, 2, 2) \mapsto (3, 2, 2, 2) \mapsto (2, 3, 2, 2) \mapsto (2, 2, 2, 4) \mapsto (2, 2, 2, 2) \\
n = 21, l = 7 : & \quad (2, 2, 2, 2) \mapsto (3, 3, 2, 2) \mapsto (2, 3, 2, 4) \mapsto (2, 2, 2, 4) \mapsto (4, 2, 2, 2) \mapsto \\
& \quad (2, 4, 2, 2) \mapsto (2, 2, 2, 6) \mapsto (2, 2, 2, 2) \\
n = 23, l = 3 : & \quad (2, 2, 2, 4) \mapsto (3, 4, 2, 2) \mapsto (2, 3, 2, 6) \mapsto (2, 2, 2, 4) \\
n = 23, l = 8 : & \quad (2, 2, 2, 2) \mapsto (3, 3, 3, 2) \mapsto (2, 4, 2, 4) \mapsto (2, 2, 2, 6) \mapsto (4, 2, 2, 2) \mapsto \\
& \quad (4, 2, 3, 2) \mapsto (2, 5, 2, 2) \mapsto (2, 2, 2, 8) \mapsto (2, 2, 2, 2) \\
n = 25, l = 2 : & \quad (2, 5, 2, 2) \mapsto (3, 2, 4, 4) \mapsto (2, 5, 2, 2) \\
n = 25, l = 3 : & \quad (2, 2, 2, 6) \mapsto (4, 4, 2, 2) \mapsto (2, 4, 2, 6) \mapsto (2, 2, 2, 6) \\
n = 25, l = 6 : & \quad (2, 2, 2, 2) \mapsto (3, 3, 3, 4) \mapsto (2, 4, 2, 4) \mapsto (4, 2, 4, 2) \mapsto (2, 6, 2, 2) \mapsto \\
& \quad (2, 2, 2, 10) \mapsto (2, 2, 2, 2) \\
n = 27, l = 5 : & \quad (2, 2, 2, 4) \mapsto (3, 3, 3, 6) \mapsto (2, 4, 2, 4) \mapsto (3, 6, 2, 2) \mapsto (2, 3, 2, 10) \mapsto \\
& \quad (2, 2, 2, 4) \\
n = 29, l = 5 : & \quad (2, 2, 2, 4) \mapsto (3, 3, 3, 8) \mapsto (2, 4, 2, 4) \mapsto (3, 7, 2, 2) \mapsto (2, 3, 2, 12) \mapsto \\
& \quad (2, 2, 2, 4) \\
n = 29, l = 7 : & \quad (2, 2, 2, 2) \mapsto (3, 4, 4, 4) \mapsto (2, 5, 2, 6) \mapsto (4, 2, 4, 4) \mapsto (6, 2, 4, 2) \mapsto \\
& \quad (2, 8, 2, 2) \mapsto (2, 2, 2, 14) \mapsto (2, 2, 2, 2) \\
n = 29, l = 8 : & \quad (2, 2, 2, 10) \mapsto (4, 4, 2, 2) \mapsto (5, 3, 4, 2) \mapsto (2, 7, 2, 4) \mapsto \\
& \quad (2, 2, 2, 12) \mapsto (4, 2, 2, 2) \mapsto (5, 4, 3, 2) \mapsto (2, 6, 2, 6) \mapsto \\
& \quad (2, 2, 2, 10) \\
n = 31, l = 5 : & \quad (2, 5, 2, 8) \mapsto (4, 2, 4, 4) \mapsto (7, 2, 2, 6) \mapsto (2, 7, 2, 2) \mapsto (3, 5, 4, 4) \mapsto \\
& \quad (2, 5, 2, 8) \\
n = 31, l = 8 : & \quad (2, 2, 2, 12) \mapsto (4, 4, 2, 2) \mapsto (6, 4, 2, 4) \mapsto (2, 6, 2, 6) \mapsto (4, 2, 4, 6) \mapsto \\
& \quad (6, 2, 4, 2) \mapsto (5, 4, 4, 2) \mapsto (2, 7, 2, 6) \mapsto (2, 2, 2, 12) \\
n = 33, l = 5 : & \quad (2, 2, 2, 2) \mapsto (4, 4, 4, 4) \mapsto (6, 2, 6, 2) \mapsto (2, 10, 2, 2) \mapsto \\
& \quad (2, 2, 2, 18) \mapsto (2, 2, 2, 2)
\end{aligned}$$

$$\begin{aligned}
n = 33, l = 5 : & \quad (2, 2, 2, 10) \mapsto (4, 4, 4, 6) \mapsto (2, 6, 2, 6) \mapsto (4, 6, 4, 2) \mapsto \\
& \quad (2, 6, 2, 10) \mapsto (2, 2, 2, 10) \\
n = 33, l = 5 : & \quad (2, 2, 2, 12) \mapsto (4, 4, 4, 2) \mapsto (5, 2, 6, 2) \mapsto (5, 5, 4, 2) \mapsto \\
& \quad (2, 7, 2, 8) \mapsto (2, 2, 2, 12)
\end{aligned}$$

There is also a cycle with $n = 23, l = 2$: $(2, 5, 2, 0) \mapsto (3, 1, 4, 4) \mapsto (2, 5, 2, 0)$, which does not qualify as a regular cycle since $k_4 = 0$ in one member, but in all other respects behaves as a primitive proper cycle, having primitive non-proper and general cycles derived from it according to Corollary 8.

Appendix F: Special cycles in base 7

There are three classes of special cycles described in Section 5, which exist for digit-counts $n = 2m + 8, m \geq 1$; $n = 5m + 10, m \geq 0$; and $n = 5m + 10, m \geq 2$, respectively. We list here all special cycles in base 7 with $n \leq 53$ that are *not* in one of these three classes. Some of these special cycles are very long, even for fairly low digit-counts; if the length is greater than 6, we only list the Kaprekar index of a single member, which in each case is the smallest integer among the members of the cycle.

$$\begin{aligned}
n = 2, l = 1 : & \quad (2, 0, 0, 0, 0, 0, 0) \mapsto (2, 0, 0, 0, 0, 0, 0) \\
& \quad [\text{repdigit from non-trivial initial integer}] \\
n = 3, l = 2 : & \quad (0, 0, 1, 0, 1, 0, 1) \mapsto (0, 0, 0, 2, 0, 0, 1) \mapsto (0, 0, 1, 0, 1, 0, 1) \\
n = 4, l = 3 : & \quad (1, 0, 0, 1, 1, 1, 0) \mapsto (1, 0, 1, 0, 0, 2, 0) \mapsto (0, 0, 2, 1, 0, 1, 0) \mapsto \\
& \quad (1, 0, 0, 1, 1, 1, 0) \\
n = 5, l = 5 : & \quad (1, 0, 0, 1, 1, 1, 1) \mapsto (0, 2, 0, 0, 1, 0, 2) \mapsto (0, 1, 1, 0, 1, 1, 1) \mapsto \\
& \quad (0, 0, 2, 1, 0, 1, 1) \mapsto (0, 0, 1, 2, 1, 0, 1) \mapsto (1, 0, 0, 1, 1, 1, 1) \\
n = 6, l = 6 : & \quad (1, 0, 1, 1, 2, 1, 0) \mapsto (1, 0, 2, 0, 1, 2, 0) \mapsto (0, 1, 1, 2, 1, 1, 0) \mapsto \\
& \quad (0, 1, 0, 1, 2, 0, 2) \mapsto (0, 0, 2, 1, 0, 1, 2) \mapsto (0, 1, 1, 1, 3, 0, 0) \mapsto \\
& \quad (1, 0, 1, 1, 2, 1, 0) \\
n = 7, l = 6 : & \quad (1, 0, 1, 2, 0, 2, 1) \mapsto (0, 2, 0, 2, 1, 0, 2) \mapsto (1, 1, 1, 0, 0, 3, 1) \mapsto \\
& \quad (0, 1, 2, 1, 1, 0, 2) \mapsto (0, 1, 2, 0, 2, 1, 1) \mapsto (0, 1, 1, 2, 1, 1, 1) \mapsto \\
& \quad (1, 0, 1, 2, 0, 2, 1) \\
n = 8, l = 6 : & \quad (0, 1, 1, 1, 3, 0, 2) \mapsto (1, 0, 2, 0, 1, 2, 2) \mapsto (1, 1, 1, 2, 1, 1, 1) \mapsto \\
& \quad (0, 2, 1, 0, 2, 0, 3) \mapsto (0, 1, 2, 1, 0, 2, 2) \mapsto (0, 1, 2, 2, 2, 1, 0) \mapsto \\
& \quad (0, 1, 1, 1, 3, 0, 2) \\
n = 9, l = 11 : & \quad (1, 0, 1, 2, 0, 2, 3) \mapsto \dots \\
n = 10, l = 2 : & \quad (0, 1, 1, 2, 1, 1, 4) \mapsto (1, 0, 2, 4, 1, 2, 0) \mapsto (0, 1, 1, 2, 1, 1, 4) \\
n = 12, l = 5 : & \quad (1, 1, 2, 2, 1, 3, 2) \mapsto (1, 2, 1, 4, 1, 2, 1) \mapsto (0, 2, 2, 0, 3, 0, 5) \mapsto \\
& \quad (0, 2, 3, 0, 3, 2, 2) \mapsto (0, 2, 1, 4, 1, 2, 2) \mapsto (1, 1, 2, 2, 1, 3, 2) \\
n = 13, l = 6 : & \quad (1, 0, 4, 2, 3, 2, 1) \mapsto (1, 1, 2, 4, 2, 1, 2) \mapsto (1, 2, 1, 2, 1, 2, 4) \mapsto \\
& \quad (0, 4, 2, 0, 3, 2, 2) \mapsto (0, 2, 4, 0, 4, 2, 1) \mapsto (0, 1, 4, 2, 4, 1, 1) \mapsto \\
& \quad (1, 0, 4, 2, 3, 2, 1) \\
n = 14, l = 9 : & \quad (0, 1, 4, 1, 6, 0, 2) \mapsto \dots
\end{aligned}$$

$n = 16, l = 12 : (0, 1, 4, 2, 4, 1, 4) \mapsto \dots$
 $n = 19, l = 4 : (1, 1, 6, 0, 5, 3, 3) \mapsto (0, 3, 2, 6, 3, 1, 4) \mapsto (1, 4, 2, 2, 1, 6, 3) \mapsto$
 $(0, 4, 3, 4, 4, 2, 2) \mapsto (1, 1, 6, 0, 5, 3, 3)$
 $n = 20, l = 5 : (1, 1, 6, 2, 5, 3, 2) \mapsto (1, 3, 3, 6, 3, 3, 1) \mapsto (0, 2, 5, 0, 6, 0, 7) \mapsto$
 $(0, 1, 6, 1, 4, 2, 6) \mapsto (0, 1, 6, 2, 6, 1, 4) \mapsto (1, 1, 6, 2, 5, 3, 2)$
 $n = 20, l = 7 : (1, 2, 4, 4, 3, 4, 2) \mapsto \dots$
 $n = 22, l = 5 : (1, 1, 7, 2, 6, 3, 2) \mapsto (1, 3, 4, 6, 4, 3, 1) \mapsto (0, 2, 6, 0, 7, 0, 7) \mapsto$
 $(0, 2, 6, 0, 6, 2, 6) \mapsto (0, 1, 6, 3, 4, 2, 6) \mapsto (1, 1, 7, 2, 6, 3, 2)$
 $n = 24, l = 5 : (1, 3, 6, 2, 5, 5, 2) \mapsto (1, 3, 5, 6, 5, 3, 1) \mapsto (0, 2, 7, 0, 8, 0, 7) \mapsto$
 $(0, 2, 7, 0, 7, 2, 6) \mapsto (0, 2, 5, 4, 5, 2, 6) \mapsto (1, 3, 6, 2, 5, 5, 2)$
 $n = 24, l = 5 : (1, 3, 5, 2, 4, 5, 4) \mapsto (1, 5, 0, 10, 0, 5, 3) \mapsto (0, 4, 4, 0, 5, 2, 9) \mapsto$
 $(1, 4, 6, 0, 5, 6, 2) \mapsto (0, 3, 5, 6, 6, 1, 3) \mapsto (1, 3, 5, 2, 4, 5, 4)$
 $n = 26, l = 24 : (1, 1, 9, 2, 8, 3, 2) \mapsto \dots$
 $n = 28, l = 7 : (1, 1, 9, 2, 8, 3, 4) \mapsto \dots$
 $n = 30, l = 25 : (1, 1, 11, 2, 10, 3, 2) \mapsto \dots$
 $n = 32, l = 13 : (1, 1, 10, 2, 9, 3, 6) \mapsto \dots$
 $n = 34, l = 13 : (1, 1, 11, 2, 10, 3, 6) \mapsto \dots$
 $n = 37, l = 5 : (1, 5, 7, 6, 6, 7, 5) \mapsto (1, 9, 2, 12, 2, 9, 2) \mapsto (1, 2, 9, 2, 9, 2, 12) \mapsto$
 $(0, 4, 10, 0, 11, 2, 10) \mapsto (0, 4, 8, 4, 8, 4, 9) \mapsto (1, 5, 7, 6, 6, 7, 5)$
 $n = 42, l = 11 : (1, 3, 12, 2, 11, 5, 8) \mapsto \dots$
 $n = 43, l = 5 : (1, 5, 9, 6, 8, 7, 7) \mapsto (1, 11, 2, 14, 2, 11, 2) \mapsto (1, 2, 11, 2, 11, 2, 14) \mapsto$
 $(0, 4, 12, 0, 13, 2, 12) \mapsto (0, 4, 10, 4, 10, 4, 11) \mapsto (1, 5, 9, 6, 8, 7, 7)$
 $n = 46, l = 7 : (1, 4, 12, 4, 11, 6, 8) \mapsto \dots$
 $n = 51, l = 3 : (1, 8, 10, 4, 9, 10, 9) \mapsto (1, 12, 0, 20, 0, 12, 6) \mapsto (0, 7, 11, 0, 12, 5, 16) \mapsto$
 $(1, 8, 10, 4, 9, 10, 9)$

References

- [1] K. Downes-Ward, Cycles in the Kaprekar Routine, unpublished dissertation, Loughborough University, 2019.
- [2] D. R. Kaprekar, Another solitaire game, *Scripta Math.* **15** (1949), 244–245.
- [3] D. R. Kaprekar, An interesting property of the number 6174, *Scripta Math.* **21** (1955), 304.
- [4] A. Kay and K. Downes-Ward, Fixed points and cycles of the Kaprekar transformation: 2. Even bases, in preparation.
- [5] J. F. Lapenta, A. L. Ludington, and G. D. Prichett, An algorithm to determine self-producing r -digit g -adic integers, *J. Reine Angew. Math.* **310** (1979), 100–110.

- [6] J. Myers, Kaprekar map, in *Online Encyclopedia of Integer Sequences*, https://oeis.org/wiki/Index_to_OEIS:_Section_K#Kaprekar_map.
- [7] J. Myers, Re: Mini-max numbers, 2013, <http://list.seqfan.eu/pipermail/seqfan/2013-May/011146.html>.
- [8] G. D. Prichett, A. L. Ludington, and J. F. Lapenta, The determination of all decadic Kaprekar constants, *Fibonacci Quart.* **19** (1981), 45–52.
- [9] N. J. A. Sloane et al., *The On-Line Encyclopedia of Integer Sequences*, available at <https://oeis.org>, 2022.
- [10] A. Yamagami, On 2-adic Kaprekar constants and 2-digit Kaprekar distances, *J. Number Theory* **185** (2018), 257–280.
- [11] A. Yamagami and Y. Matsui, On 3-adic Kaprekar loops, *JP J. Algebra Number Theory Appl.* **40** (2018), 957–1028.

2020 *Mathematics Subject Classification*: Primary 11A99.

Keywords: Kaprekar transformation, cycle, fixed point.

(Concerned with sequences [A000010](#), [A003558](#), [A165002](#), [A165003](#), [A165041](#), [A165042](#), [A165080](#), [A165081](#), [A165119](#), and [A165120](#).)

Received April 21 2022; revised versions received July 17 2022; July 18 2022; July 21 2022.
Published in *Journal of Integer Sequences*, July 22 2022.

Return to [Journal of Integer Sequences home page](#).