

Generating Functions for Straight Polyomino Tilings of Narrow Rectangles

Mudit Aggarwal and Samrith Ram¹

Indraprastha Institute of Information Technology Delhi

New Delhi 110020

India

mudit19063@iiitd.ac.in

samrith@iiitd.ac.in

Abstract

Let m, k be fixed positive integers. Determining the generating function for the number of tilings of an $m \times n$ rectangle by $k \times 1$ rectangles is a longstanding open problem to which the answer is only known in certain special cases. We give an explicit formula for this generating function in the case where $m < 2k$. This result is used to obtain the generating function for the number of tilings of an $m \times n \times k$ box with $k \times k \times 1$ bricks.

1 Introduction and main results

We consider the problem of enumerating the number of tilings of an $m \times n$ rectangle with $k \times 1$ tiles. An example of such a tiling for $k = 3, m = 5$ and $n = 6$ is shown in Figure 1. By a theorem of Klarner [9, Thm. 5] such tilings exist if and only if k divides either m or n . Let $h_k(m, n)$ denote the number of such tilings. A beautiful result of Kasteleyn [7] and Temperley-Fisher [13] gives an explicit formula for $h_2(m, n)$:

$$h_2(m, n) = \prod_{j=1}^{\lceil \frac{m}{2} \rceil} \prod_{k=1}^{\lceil \frac{n}{2} \rceil} \left(4 \cos^2 \frac{j\pi}{m+1} + 4 \cos^2 \frac{k\pi}{n+1} \right). \quad (1)$$

¹The second author was partially supported by a MATRICS grant MTR/2017/000794 awarded by the Science and Engineering Research Board.




Figure 1: Tiling a 5×6 rectangle with 3×1 tiles.

Equation (1) is represented by the OEIS sequence [A099390](#). For arbitrary integers m, n , no such formula is known for $h_k(m, n)$ for any integer $k \geq 3$. However, for fixed values of m and k , the generating function $H_{k,m}(x) = \sum_{n \geq 0} h_k(m, n)x^n$ can be obtained by using the transfer-matrix method as described by Read [11], which expresses the number of tilings as the number of walks between two vertices in a suitably defined digraph. Klarner and Pollack [8] computed $H_{2,m}(x)$ for $m \leq 8$ and gave an algorithm to compute polynomials P_m and Q_m such that $H_{2,m}(x) = P_m(x)/Q_m(x)$. Hock and McQuistan [6] found recurrences for $h_2(m, n)$ in terms of n for each integer $m \leq 10$. Stanley [12] used the explicit formula (1) above to determine the degrees of the polynomials $P_m(x)$ and $Q_m(x)$ and prove several properties of the polynomials $P_m(x)$ and $Q_m(x)$.

By using the transfer-matrix approach, Mathar [10] derived several generating functions which enumerate tilings of the $m \times n$ rectangle with $a \times b$ tiles for fixed values of a, b . Tilings can also be enumerated by using integer linear programming; see, for instance, Garvie and Burkardt [5]. It is worth noting that these methods can be cumbersome for large values of the parameters.

For $k \geq 3$, there does not appear to have been substantial progress on the general problem of computing $H_{k,m}(x)$. In this paper, we compute $H_{k,m}(x)$ for arbitrary positive integers m and k satisfying $m < 2k$. The cases where $m \leq k$ are somewhat trivial but the case $k < m < 2k$ is more interesting and we prove (see Theorem 8) that the generating function is surprisingly simple:

$$\sum_{n \geq 0} h_k(m, n)x^n = \frac{(1 - x^k)^{k-1}}{(1 - x^k)^k - (m - k + 1)x^k}. \quad (2)$$

The generating function above allows for recursive computation of $h_k(m, n)$, and asymptotic estimates for large values of n can be obtained by considering the smallest positive root of the denominator (see Flajolet and Sedgewick [4, p. 255]). In Theorem 9 we obtain a refinement of Equation (2) by proving that if $b_k(m, n, r)$ denotes the number of tilings in which precisely r tiles are vertical, then

$$\sum_{n, r \geq 0} b_k(m, n, r)x^n y^r = \frac{(1 - x^k)^{k-1}}{(1 - x^k)^k - (m - k + 1)x^k y^k}.$$

Under the same constraints on m , our results can also be used to derive the generating function (see Theorem 12) for $\bar{h}_k(m, n)$, the number of tilings of an $m \times n \times k$ box with

$k \times k \times 1$ bricks:

$$\sum_{n \geq 0} \bar{h}_k(m, n) x^n = \frac{(1 - 2x^k)^{k-1}}{(1 - 2x^k)^{k-1} (1 - (m - k + 2)x^k) - (m - k + 1)x^k}.$$

In this case, we also obtain a refined multivariate generating function which accounts for tilings with a specific number of tiles in a given orientation. More precisely, we prove (see Corollary 13) that if $d_k(m, n, r, s)$ denotes the number of tilings of an $m \times n \times k$ box with $k \times k \times 1$ bricks which contain precisely r bricks parallel to the yz -plane and s bricks parallel to the xy -plane, then

$$\sum_{n, r, s \geq 0} d_k(m, n, r, s) x^n y^r z^s = \frac{(1 - x^k - x^k z^k)^{k-1}}{(1 - x^k - (m - k + 1)x^k z^k)(1 - x^k - x^k z^k)^{k-1} - (m - k + 1)x^k y^k}.$$

The key idea of our approach is to enumerate vertically fault-free tilings, which are explained in the next section.

2 Vertically fault-free tilings

Fix, throughout this paper, a positive integer $k \geq 2$. It will be convenient to introduce a coordinate system with the origin at the bottom left corner of the $m \times n$ rectangle such that a side of length m of the rectangle is along the y -axis. Each of the unit squares, or cells, of the $m \times n$ rectangle is then represented by a pair (r, s) corresponding to the coordinates of its top right corner. For instance the cells $(1, 1)$ and $(3, 4)$ are shaded in Figure 2.




Figure 2: The cells $(1, 1)$ and $(3, 4)$ of a 4×5 rectangle.

A tiling of the $m \times n$ rectangle is said to have a *vertical fault* at $x = a$ for some $1 \leq a \leq n - 1$ if the line $x = a$ does not intersect the interior of any tile. For instance, the tiling in Figure 3 has only one vertical fault at $x = 1$ while the tiling in Figure 4 has no vertical faults; such a tiling is called *vertically fault-free*. It is easy to see that if a tiling has l vertical faults then it can be decomposed uniquely into $l + 1$ vertically fault-free tilings. Let $a(m, n)$ denote the number of vertically fault-free tilings of an $m \times n$ rectangle with $k \times 1$ tiles. The following well-known result [4, Thm. I.1, p. 27] relates all tilings $h(m, n) = h_k(m, n)$ to vertically fault-free tilings.




Figure 3: A vertical fault at $x = 1$.




Figure 4: A vertically fault-free tiling.

Lemma 1. If $H(x) = \sum_{n \geq 0} h(m, n)x^n$ and $A(x) = \sum_{n \geq 0} a(m, n)x^n$, then

$$H(x) = \frac{1}{1 - A(x)}.$$

For $m < k$ it is clear that $h(m, n) = 1$ when n is a multiple of k and 0 otherwise. For $m = k$ we have the following proposition.

Proposition 2. For $k > 1$, we have

$$\sum_{n \geq 0} h(k, n)x^n = \frac{1}{1 - x - x^k}.$$

Proof. Here $a(k, 1) = 1 = a(k, k)$ while $a(k, n) = 0$ for $n \notin \{1, k\}$ (see Figure 5). Thus the generating function for vertically fault-free tilings is $A(x) = x + x^k$ and the proposition follows from Lemma 1. \square




Figure 5: All possible vertically fault-free tilings for $m = k = 3$.

Remark 3. It follows from Proposition 2 that tilings of a $k \times n$ rectangle with $k \times 1$ tiles are in bijection with compositions of n (tuples (n_1, \dots, n_r) of positive integers with $\sum n_i = n$) in which all parts are equal to 1 or k .

Proposition 2 corresponds to the OEIS sequences [A000045](#), [A000930](#), [A003269](#), [A003520](#), [A005708](#), [A005709](#), and [A005710](#). We now consider the case $k < m < 2k$. In this case, the following lemma is the key to computing vertically fault-free tilings.

Lemma 4. Suppose $k < m < 2k$ and $n > k$. In any vertically fault-free tiling of an $m \times n$ rectangle by $k \times 1$ tiles there exist k contiguous rows such that all tiles in the remaining $m - k$ rows are horizontal.

Proof. A cell (i, j) is said to be in row i and column j . Consider a vertically fault-free tiling of the $m \times n$ rectangle for $n > k$. Not all cells in column 1 are covered by horizontal tiles since this would create a vertical fault at $x = k$. It follows that there is precisely one vertical tile in the first column (see Figure 6); suppose the topmost cell of this tile is $(1, b)$.




Figure 6: Vertical tile in the first column.

We claim that the $m - k$ rows that do not intersect this vertical tile consist of only horizontal tiles. If not, then consider the leftmost vertical tile that intersects one of the aforementioned rows and suppose its top cell is (a', b') where $b' \neq b$. By reflecting through a horizontal line if necessary, we may assume without loss of generality that $b' > b$ (see Figure 7). By the minimality of a' , it follows that $a' = k\ell + 1$ for some positive integer ℓ .




Figure 7: The contradiction obtained.

Moreover, the first $k\ell$ cells in each row that does not intersect the vertical tile in the first column are covered by ℓ horizontal tiles. Now consider the tile covering the cell $(k\ell, b)$. If this tile were vertical, there would be a vertical fault at $x = k\ell$ and, therefore, this tile must be horizontal. By backtracking, the tile covering the cell $(k(\ell - 1), b)$ must be horizontal. Continuing this line of reasoning, it is clear that the tile covering the cell (k, b) must be horizontal which is impossible since this tile would then cover $(1, b)$ which is already covered by the vertical tile in the first column. This proves the claim and the lemma. \square

The idea used to prove Lemma 4 can also be used to show that for $n > k$, any vertically fault-free tiling of an $m \times n$ rectangle in which each tile has dimensions $k \times j$ for some $1 \leq j \leq k$ has $m - k$ rows consisting of only horizontal tiles but we do not require this stronger result here. In the following proposition we consider $m \times n$ rectangles where n is a multiple of ℓ since these cover all admissible values of n for which tilings exist by Klarner's divisibility criterion.

Proposition 5. *Let $k > 1$ be a positive integer and suppose $k < m < 2k$. The number of vertically fault-free tilings of an $m \times k\ell$ rectangle is given by*

$$a(m, k\ell) = \begin{cases} m - k + 2, & \text{if } \ell = 1; \\ (m - k + 1) \binom{k+\ell-3}{k-2}, & \text{if } \ell \geq 2. \end{cases}$$

Proof. For an $m \times k$ rectangle there is one vertically fault-free tiling in which all tiles are horizontal. Any other vertically fault-free tiling of this rectangle contains precisely $m - k$ horizontal tiles and k vertical tiles; there are $m - k + 1$ such tilings. Thus $a(m, k) = m - k + 2$.

Now consider a vertically fault-free tiling of an $m \times k\ell$ rectangle. Such a tiling has $m - k$ horizontal rows by Lemma 4. If these $m - k$ rows are removed, what remains is a tiling of a $k \times k\ell$ rectangle in which no vertical faults occur at $x = jk$ for $j \geq 1$; denote by N the number of tilings so obtained. By Remark 3, these tilings are in bijection with compositions (n_1, \dots, n_r) of $k\ell$ where each $n_i \in \{1, k\}$ and such that k does not divide $\sum_{i=1}^s n_i$ for each $s < r$. This condition on the partial sums implies $n_1 = 1 = n_r$. Further, the number of n_i 's equal to 1 must be a multiple of k but the condition on the partial sums ensures that no more than k of them can equal 1. It follows that precisely k of the n_i are 1 and hence $r = k + \ell - 1$. Since precisely $k - 2$ of the n_i ($2 \leq i \leq r - 1$) are equal to 1 and all these choices are possible, we obtain $N = \binom{k+\ell-3}{k-2}$. On the other hand, the number of vertically fault-free tilings of an $m \times k\ell$ rectangle that yield a given $k \times k\ell$ rectangle upon removing the $m - k$ horizontal rows is clearly $m - k + 1$. Therefore

$$a(m, k\ell) = (m - k + 1) \binom{k + \ell - 3}{\ell - 1}. \quad \square$$

Corollary 6. *Under the hypotheses of Proposition 5 every vertically fault-free tiling of an $m \times k\ell$ rectangle ($\ell > 1$) by $k \times 1$ tiles contains precisely k vertical tiles.*

Remark 7. It is clear from the proof of Proposition 5 that each vertically fault-free tiling of an $m \times k\ell$ rectangle ($\ell > 1; k < m < 2k$) contains precisely $\ell - 1$ 'blocks', where a block is defined as a collection of k contiguous horizontal $k \times 1$ tiles, one on top of the other.

Theorem 8. *Suppose $k < m < 2k$ and $h_k(m, n)$ denotes the number of tilings of an $m \times n$ rectangle with $k \times 1$ tiles. Then*

$$\sum_{n \geq 0} h_k(m, n) x^n = \frac{(1 - x^k)^{k-1}}{(1 - x^k)^k - (m - k + 1)x^k}.$$

Proof. By Proposition 5, it follows that the generating function for vertically fault-free tilings is given by

$$\begin{aligned}
A(x) &= \sum_{\ell \geq 1} a(m, k\ell)x^{k\ell} \\
&= (m - k + 2)x^k + (m - k + 1) \sum_{\ell \geq 2} \binom{k + \ell - 3}{\ell - 1} x^{k\ell} \\
&= x^k + \frac{(m - k + 1)x^k}{(1 - x^k)^{k-1}}. \tag{3}
\end{aligned}$$

The theorem now follows from Lemma 1. \square

Several OEIS sequences which correspond to the generating function in Theorem 8 are shown in Table 1.

| k | m | OEIS entry |
|-----|-----|-------------------------|
| 2 | 3 | A001835 |
| 3 | 4 | A049086 |
| 3 | 5 | A236576 |
| 4 | 5 | A236579 |
| 4 | 6 | A236580 |
| 4 | 7 | A236581 |

Table 1: OEIS entries

Theorem 8 may be viewed as a special case of the following more general result.

Theorem 9. *Suppose $k < m < 2k$ and let $b_k(m, n, r)$ denote the number of $k \times 1$ tilings of an $m \times n$ rectangle which contain precisely r vertical tiles. Then*

$$\sum_{n, r \geq 0} b_k(m, n, r)x^n y^r = \frac{(1 - x^k)^{k-1}}{(1 - x^k)^k - (m - k + 1)x^k y^k}. \tag{4}$$

Proof. Consider the number of vertical tiles in the vertically fault-free tilings of an $m \times k\ell$ rectangle by $k \times 1$ tiles. For $\ell = 1$, there is one such tiling with no vertical tiles and $m - k + 1$ tilings with precisely k vertical tiles. By Corollary 6, each such vertically fault-free tiling for $\ell > 1$ contains precisely k vertical tiles. Therefore the generating function for $b_k(m, n, r)$ is

$$\frac{1}{1 - A(x, y)},$$

where, $A(x, y)$ can be computed from the expression for $A(x)$ in Equation (3) as

$$A(x, y) = x^k + \frac{(m - k + 1)x^k y^k}{(1 - x^k)^{k-1}}. \tag{4}$$

3 Brick tilings of a box

A 3-dimensional analog of Klarner's divisibility criterion for filling boxes with bricks can be found in de Bruijn [3] (see also Boisen [1]) who proved that an $A \times B \times C$ box can be tiled with $a \times ab \times abc$ bricks if and only if the numbers A, B, C are (in some order) a multiple of a , a multiple of ab , and a multiple of abc . A higher dimensional analog appears in Bower and Michael [2]. The results of the previous section can be used to derive the generating function for the number of tilings of an $m \times n \times k$ box with $k \times k \times 1$ bricks. Such tilings exist if and only if k divides either m or n . In order to prove the result we will require the following theorem on tilings of an $m \times n$ rectangle with $k \times 1$ and $k \times k$ tiles. Vertically fault-free tilings for $m \leq k$ are easily enumerated, so we consider the case $m > k$.

Theorem 10. *Suppose $k < m < 2k$ and $h'(m, n)$ denotes the number of tilings of an $m \times n$ rectangle with $k \times 1$ tiles and $k \times k$ tiles. Then*

$$\sum_{n \geq 0} h'(m, n)x^n = \frac{(1 - 2x^k)^{k-1}}{(1 - 2x^k)^{k-1}(1 - (m - k + 2)x^k) - (m - k + 1)x^k}.$$

Proof. Let $a'(m, n)$ denote the number of vertically fault-free tilings of an $m \times n$ rectangle with $k \times 1$ and $k \times k$ tiles. For vertically fault-free tilings of an $m \times k$ rectangle, we have

1. 1 tiling with only horizontal $k \times 1$ tiles;
2. $m - k + 1$ tilings containing a $k \times k$ tile;
3. $m - k + 1$ tilings containing precisely k vertical $k \times 1$ tiles.

Thus $a'(m, k) = 2m - 2k + 3$. Now suppose $n = k\ell$ with $\ell > 1$. Let $T'_k(m, n)$ be the set of all vertically fault-free tilings of an $m \times n$ rectangle by $k \times 1$ and $k \times k$ tiles. Denote by $T_k(m, n)$ the subset of $T'_k(m, n)$ consisting of vertically fault-free tilings of an $m \times n$ rectangle with only $k \times 1$ tiles. To each tiling $T' \in T'_k(m, n)$, we can associate a new tiling by replacing each $k \times k$ tile in T' by k horizontal $k \times 1$ tiles (see Figure 8); it is easily seen that this new tiling is vertically fault-free, and therefore lies in $T_k(m, n)$. This correspondence gives




Figure 8: A vertically fault-free tiling and its image under Ψ .

a map $\Psi : T'_k(m, n) \rightarrow T_k(m, n)$, which is clearly surjective. In fact the fibers of Ψ are of

cardinality $2^{\ell-1}$. To see this, note that each $T \in T_k(m, n)$ has precisely $\ell - 1$ blocks as defined in Remark 7. The tilings in $\Psi^{-1}(T)$ are precisely those which can be obtained from T by optionally replacing each of the blocks in T by a $k \times k$ tile, for a total of $2^{\ell-1}$ choices. In summary, we have

$$a'(m, k\ell) = \begin{cases} 2m - 2k + 3, & \text{if } \ell = 1; \\ 2^{\ell-1}(m - k + 1) \binom{k+\ell-3}{\ell-1}, & \text{if } \ell \geq 2. \end{cases}$$

The generating function for vertically fault-free tilings is therefore

$$A(x) = (m - k + 2)x^k + (m - k + 1) \frac{x^k}{(1 - 2x^k)^{k-1}}.$$

The theorem now follows from Lemma 1. □

For $k = 2$ and $m = 3$, Theorem 10 corresponds to the OEIS sequence [A083066](#). The next result extends the above theorem by accounting for the number of tiles of a given type and orientation.

Theorem 11. *Suppose $k < m < 2k$ and let $c_k(m, n, r, s)$ denote the number of tilings of an $m \times n$ rectangle with $k \times 1$ and $k \times k$ tiles which contain precisely r vertically placed $k \times 1$ tiles and s square tiles. Then*

$$\sum_{n, r, s \geq 0} c_k(m, n, r, s) x^n y^r z^s = \frac{(1 - x^k - x^k z)^{k-1}}{(1 - x^k - (m - k + 1)x^k z)(1 - x^k - x^k z)^{k-1} - (m - k + 1)x^k y^k}. \quad (5)$$

Proof. We argue as in the proof of Theorem 9. If $C(x, y, z)$ denotes the generating function for $c_k(m, n, r, s)$, then

$$C(x, y, z) = \frac{1}{1 - A(x, y, z)},$$

where $A(x, y, z)$ is computed from the discussion in the proof of Theorem 10 as

$$\begin{aligned} A(x, y, z) &= x^k(1 + (m - k + 1)z + (m - k + 1)y^k) \\ &\quad + \sum_{\ell \geq 2} (m - k + 1) \binom{\ell + k - 3}{\ell - 1} x^{k\ell} y^k (1 + z)^{\ell-1} \\ &= x^k(1 + (m - k + 1)z) \\ &\quad + \sum_{\ell \geq 1} (m - k + 1) \binom{\ell + k - 3}{\ell - 1} y^k x^{k\ell} (1 + z)^{\ell-1} \\ &= x^k(1 + (m - k + 1)z) + \frac{(m - k + 1)x^k y^k}{(1 - x^k - x^k z)^{k-1}}. \end{aligned} \quad \square$$

Note that if we set $z = 0$ in Theorem 11 then we obtain Theorem 9 while the substitution $y = z = 1$ yields Theorem 10.

Theorem 12. *Suppose $k < m < 2k$ and let $\bar{h}(m, n)$ denote the number of tilings of an $m \times n \times k$ box with $k \times k \times 1$ bricks. Then*

$$\sum_{n \geq 0} \bar{h}(m, n) x^n = \frac{(1 - 2x^k)^{k-1}}{(1 - 2x^k)^{k-1} (1 - (m - k + 2)x^k) - (m - k + 1)x^k}.$$

Proof. Consider a box in the positive octant with a corner at the origin and its sides of length n, m and k along the x, y and z -axes, respectively. Let τ be a tiling of this box by $k \times k \times 1$ bricks. The bricks of the tiling which touch the xy -plane determine a tiling τ' of an $m \times n$ rectangle with $k \times 1$ and $k \times k$ tiles as shown in Figure 9. In fact, τ is uniquely




Figure 9: A tiling τ' for $m = 5, n = 12$ and $k = 3$.

determined by τ' as follows. The tiles in τ' can be seen to correspond to the projections of the bricks in τ onto the xy -plane: each $k \times 1$ tile of τ' is the projection of a single brick of τ while each $k \times k$ tile of τ' is the projection of precisely k bricks in τ . This gives a one-one correspondence between tilings of the box by $k \times k \times 1$ bricks and tilings of an $m \times n$ rectangle by tiles of size $k \times 1$ and $k \times k$. The result now follows from Theorem 10. \square

Replacing z by z^k in the generating function (5), the following result is obtained.

Corollary 13. *Suppose $k < m < 2k$ and let $d_k(m, n, r, s)$ denote the number of tilings of an $m \times n \times k$ box with $k \times k \times 1$ bricks which contain precisely r bricks parallel to the yz -plane and s bricks parallel to the xy -plane. Then*

$$\sum_{n, r, s \geq 0} d_k(m, n, r, s) x^n y^r z^s = \frac{(1 - x^k - x^k z^k)^{k-1}}{(1 - x^k - (m - k + 1)x^k z^k)(1 - x^k - x^k z^k)^{k-1} - (m - k + 1)x^k y^k}.$$

4 Acknowledgments

We thank an anonymous referee for several helpful comments and suggestions.

References

- [1] Paul Boisen, Polynomials and packings: a new proof of de Bruijn’s theorem, *Discrete Math.* **146** (1995), 285–287.
- [2] Richard J. Bower and T. S. Michael, When can you tile a box with translates of two given rectangular bricks?, *Electron. J. Combin.* **11** (2004), Note 7, 9.
- [3] N. G. de Bruijn, Filling boxes with bricks, *Amer. Math. Monthly* **76** (1969), 37–40.
- [4] Philippe Flajolet and Robert Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [5] Marcus R. Garvie and John Burkardt, A parallelizable integer linear programming approach for tiling finite regions of the plane with polyominoes, *Algorithms* **15** (2022), 164.
- [6] J. L. Hock and R. B. McQuistan, A note on the occupational degeneracy for dimers on a saturated two-dimensional lattice space, *Discrete Appl. Math.* **8** (1984), 101–104.
- [7] P. W. Kasteleyn, Dimer statistics and phase transitions, *J. Mathematical Phys.* **4** (1963), 287–293.
- [8] David Klarner and Jordan Pollack, Domino tilings of rectangles with fixed width, *Discrete Math.* **32** (1980), 45–52.
- [9] David A. Klarner, Packing a rectangle with congruent n -ominoes, *J. Combinatorial Theory* **7** (1969), 107–115.
- [10] R. J. Mathar, Tilings of rectangular regions by rectangular tiles: Counts derived from transfer matrices, arxiv preprint arXiv:1406.7788 [math.CO], 2014. Available at <https://arxiv.org/abs/1406.7788>.
- [11] Ronald C. Read, A note on tiling rectangles with dominoes, *Fibonacci Quart.* **18** (1980), 24–27.
- [12] Richard P. Stanley, On dimer coverings of rectangles of fixed width, *Discrete Appl. Math.* **12** (1985), 81–87.
- [13] H. N. V. Temperley and Michael E. Fisher, Dimer problem in statistical mechanics—an exact result, *Philos. Mag.* **6** (1961), 1061–1063.

2020 *Mathematics Subject Classification*: Primary 05B45; Secondary 05A15, 05A19, 05B50.
Keywords: tiling, generating function, polyomino tiling, brick tiling.

(Concerned with sequences [A000045](#), [A000930](#), [A001835](#), [A003269](#), [A003520](#), [A005708](#), [A005709](#), [A005710](#), [A049086](#), [A083066](#), [A099390](#), [A236576](#), [A236579](#), [A236580](#), and [A236581](#).)

Received November 15 2022; revised versions received December 9 2022; December 13 2022; December 20 2022. Published in *Journal of Integer Sequences*, January 1 2023.

Return to [Journal of Integer Sequences home page](#).