

The Akiyama-Tanigawa algorithm for Bernoulli numbers

Masanobu Kaneko

Graduate School of Mathematics
Kyushu University
Fukuoka 812-8581, Japan

Email address: mkaneko@math.kyushu-u.ac.jp

Abstract

A direct proof is given for Akiyama and Tanigawa's algorithm for computing Bernoulli numbers. The proof uses a closed formula for Bernoulli numbers expressed in terms of Stirling numbers. The outcome of the same algorithm with different initial values is also briefly discussed.

1 The Algorithm

In their study of values at non-positive integer arguments of multiple zeta functions, S. Akiyama and Y. Tanigawa [1] found as a special case an amusing algorithm for computing Bernoulli numbers in a manner similar to “Pascal’s triangle” for binomial coefficients.

Their algorithm reads as follows: Start with the 0-th row $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots$ and define the first row by $1 \cdot (1 - \frac{1}{2}), 2 \cdot (\frac{1}{2} - \frac{1}{3}), 3 \cdot (\frac{1}{3} - \frac{1}{4}), \dots$ which produces the sequence $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$. Then define the next row by $1 \cdot (\frac{1}{2} - \frac{1}{3}), 2 \cdot (\frac{1}{3} - \frac{1}{4}), 3 \cdot$

$(\frac{1}{4} - \frac{1}{5}), \dots$, thus giving $\frac{1}{6}, \frac{1}{6}, \frac{3}{20}, \dots$ as the second row. In general, denoting the m -th ($m = 0, 1, 2, \dots$) number in the n -th ($n = 0, 1, 2, \dots$) row by $a_{n,m}$, the m -th number in the $(n + 1)$ -st row $a_{n+1,m}$ is determined recursively by

$$a_{n+1,m} = (m + 1) \cdot (a_{n,m} - a_{n,m+1}).$$

Then the claim is that the 0-th component $a_{n,0}$ of each row (the “leading diagonal”) is just the n -th Bernoulli numbers B_n , where

$$\sum_{n=0}^{\infty} B_n \frac{x^n}{n!} = \frac{xe^x}{e^x - 1} \left(= \frac{x}{e^x - 1} + x \right).$$

Note that we are using the definition of the Bernoulli numbers in which $B_1 = \frac{1}{2}$. This is the definition used by Bernoulli (and independently Seki, published one year prior to Bernoulli). Incidentally, this is more appropriate for the Euler formula $\zeta(1 - k) = -B_k/k$ ($k = 1, 2, 3, \dots$) for the values of the Riemann zeta function.

2 Proof

The proof is based on the following identity for Bernoulli numbers, a variant of which goes as far back as Kronecker (see [4]). Here we denote by $\left\{ \begin{smallmatrix} n \\ m \end{smallmatrix} \right\}$ the Stirling number of the second kind:

$$x^n = \sum_{m=0}^n \left\{ \begin{smallmatrix} n \\ m \end{smallmatrix} \right\} x^m,$$

where $x^m = x(x - 1) \cdots (x - m + 1)$ for $m \geq 1$ and $x^0 = 1$. (We use Knuth’s notation [7]. For the Stirling number identities that we shall need, the reader is referred for example to [5].)

Theorem 1

$$B_n = \sum_{m=0}^n \frac{(-1)^m m! \left\{ \begin{smallmatrix} n+1 \\ m+1 \end{smallmatrix} \right\}}{m + 1}, \quad \forall n \geq 0.$$

We shall give later a proof of this identity for the sake of completeness. Once we have this, the next proposition ensures the validity of our algorithm.




Figure 1: Akiyama-Tanigawa triangle

Proposition 2 Given an initial sequence $a_{0,m}$ ($m = 0, 1, 2, \dots$), define the sequences $a_{n,m}$ ($n \geq 1$) recursively by

$$a_{n,m} = (m + 1) \cdot (a_{n-1,m} - a_{n-1,m+1}) \quad (n \geq 1, m \geq 0). \quad (1)$$

Then

$$a_{n,0} = \sum_{m=0}^n (-1)^m m! \left\{ \begin{matrix} n+1 \\ m+1 \end{matrix} \right\} a_{0,m}. \quad (2)$$

Proof. Put

$$g_n(t) = \sum_{m=0}^{\infty} a_{n,m} t^m.$$

By the recursion (1) we have for $n \geq 1$

$$\begin{aligned}
g_n(t) &= \sum_{m=0}^{\infty} (m+1)(a_{n-1,m} - a_{n-1,m+1})t^m \\
&= \frac{d}{dt} \left(\sum_{m=0}^{\infty} a_{n-1,m} t^{m+1} \right) - \frac{d}{dt} \left(\sum_{m=0}^{\infty} a_{n-1,m+1} t^{m+1} \right) \\
&= \frac{d}{dt} (t g_{n-1}(t)) - \frac{d}{dt} (g_{n-1}(t) - a_{n-1,0}) \\
&= g_{n-1}(t) + (t-1) \frac{d}{dt} (g_{n-1}(t)) \\
&= \frac{d}{dt} ((t-1) g_{n-1}(t)).
\end{aligned}$$

Hence, by putting $(t-1)g_n(t) = h_n(t)$ we obtain

$$h_n(t) = (t-1) \frac{d}{dt} (h_{n-1}(t)) \quad (n \geq 1),$$

and thus

$$h_n(t) = \left((t-1) \frac{d}{dt} \right)^n (h_0(t)).$$

Applying the formula (*cf.* [5, Ch. 6.7 Exer. 13])

$$\left(x \frac{d}{dx} \right)^n = \sum_{m=0}^n \left\{ \begin{matrix} n \\ m \end{matrix} \right\} x^m \left(\frac{d}{dx} \right)^m$$

for $x = t - 1$, we have

$$h_n(t) = \sum_{m=0}^n \left\{ \begin{matrix} n \\ m \end{matrix} \right\} (t-1)^m \left(\frac{d}{dt} \right)^m h_0(t).$$

Putting $t = 0$ we obtain

$$\begin{aligned}
-a_{n,0} &= \sum_{m=0}^n \left\{ \begin{matrix} n \\ m \end{matrix} \right\} (-1)^m m! (a_{0,m-1} - a_{0,m}) \\
&= \sum_{m=0}^{n-1} \left\{ \begin{matrix} n \\ m+1 \end{matrix} \right\} (-1)^{m+1} (m+1)! a_{0,m} - \sum_{m=0}^n \left\{ \begin{matrix} n \\ m \end{matrix} \right\} (-1)^m m! a_{0,m} \\
&= - \sum_{m=0}^n (-1)^m m! a_{0,m} \left((m+1) \left\{ \begin{matrix} n \\ m+1 \end{matrix} \right\} + \left\{ \begin{matrix} n \\ m \end{matrix} \right\} \right) \\
&= - \sum_{m=0}^n (-1)^m m! \left\{ \begin{matrix} n+1 \\ m+1 \end{matrix} \right\} a_{0,m}.
\end{aligned}$$

(We have used the recursion $\left\{ \begin{matrix} n+1 \\ m+1 \end{matrix} \right\} = (m+1) \left\{ \begin{matrix} n \\ m+1 \end{matrix} \right\} + \left\{ \begin{matrix} n \\ m \end{matrix} \right\}$.) This proves the proposition.

Proof of Theorem 1. We show the generating series of the right hand side coincide with that of B_n . To do this, we use the identity

$$\frac{e^x (e^x - 1)^m}{m!} = \sum_{n=m}^{\infty} \left\{ \begin{matrix} n+1 \\ m+1 \end{matrix} \right\} \frac{x^n}{n!} \quad (3)$$

which results from the well-known generating series for the Stirling numbers (cf. [5, (7.49)])

$$\frac{(e^x - 1)^m}{m!} = \sum_{n=m}^{\infty} \left\{ \begin{matrix} n \\ m \end{matrix} \right\} \frac{x^n}{n!}$$

by replacing m with $m+1$ and differentiating with respect to x . With this, we have

$$\begin{aligned}
&\sum_{n=0}^{\infty} \left(\sum_{m=0}^n \frac{(-1)^m m! \left\{ \begin{matrix} n+1 \\ m+1 \end{matrix} \right\}}{m+1} \right) \frac{x^n}{n!} \\
&= \sum_{m=0}^{\infty} \frac{(-1)^m m!}{m+1} \sum_{n=m}^{\infty} \left\{ \begin{matrix} n+1 \\ m+1 \end{matrix} \right\} \frac{x^n}{n!} = \sum_{m=0}^{\infty} \frac{(-1)^m m!}{m+1} \frac{e^x (e^x - 1)^m}{m!} \\
&= e^x \sum_{m=0}^{\infty} \frac{(1 - e^x)^m}{m+1} = \frac{e^x}{1 - e^x} \sum_{m=1}^{\infty} \frac{(1 - e^x)^m}{m} \\
&= \frac{e^x}{1 - e^x} (-\log(1 - (1 - e^x))) = \frac{x e^x}{e^x - 1}.
\end{aligned}$$

This proves Theorem 1.

Remark. A referee suggested the following interpretation of the algorithm using generating function:

Suppose the first row is a_0, a_1, a_2, \dots , with ordinary generating function

$$A(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Let the leading diagonal be $b_0 = a_0, b_1, b_2, \dots$, with exponential generating function

$$\mathbb{B}(x) = \sum_{n=0}^{\infty} b_n \frac{x^n}{n!}.$$

Then we have

$$\mathbb{B}(x) = e^x A(1 - e^x).$$

This follows from (2) and (3), the calculation being parallel to that of the proof of Theorem 1. To get the Bernoulli numbers we take $a_0 = 1, a_1 = \frac{1}{2}, a_2 = \frac{1}{3}, \dots$ with $A(x) = -\log(1 - x)/x$, and find $\mathbb{B}(x) = xe^x/(e^x - 1)$.

3 Poly-Bernoulli numbers

If we replace the initial sequence $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$ by $1, \frac{1}{2^k}, \frac{1}{3^k}, \frac{1}{4^k}, \dots$ and apply the same algorithm, the resulting sequence is $(-1)^n D_n^{(k)}$ ($n = 0, 1, 2, \dots$), where $D_n^{(k)}$ is a variant of “poly-Bernoulli numbers” ([6], [2], [3]): For any integer k , we define a sequence of numbers $D_n^{(k)}$ by

$$\frac{Li_k(1 - e^{-x})}{e^x - 1} = \sum_{n=0}^{\infty} D_n^{(k)} \frac{x^n}{n!},$$

where $Li_k(t) = \sum_{m=1}^{\infty} \frac{t^m}{m^k}$ (k -th polylogarithm when $k \geq 1$). The above assertion is then a consequence of the following (or, is just a special case of the preceding remark)

Proposition 3 *For any $k \in \mathbf{Z}$ and $n \geq 0$, we have*

$$D_n^{(k)} = (-1)^n \sum_{m=0}^n \frac{(-1)^m m! \{m+1\}}{(m+1)^k}.$$

Proof. The proof can be given completely in the same way as the proof of Theorem 1 using generating series, and hence will be omitted.

Acknowledgements

I should like to thank the referee for several comments and suggestions.

References

- [1] Akiyama, S. and Tanigawa, Y. : Multiple zeta values at non-positive integers, *preprint* (1999).
- [2] Arakawa, T. and Kaneko, M. : Multiple zeta values, poly-Bernoulli numbers, and related zeta functions, *Nagoya Math. J.* **153** (1999), 189–209.
- [3] Arakawa, T. and Kaneko, M. : On poly-Bernoulli numbers, *Comment. Math. Univ. Sanct. Pauli* **48-2** (1999), 159–167.
- [4] Gould, H. G. : Explicit formulas for Bernoulli numbers, *Amer. Math. Monthly* **79** (1972), 44–51.
- [5] Graham, R., Knuth, D. and Patashnik, O.: Concrete Mathematics, Addison-Wesley, (1989).
- [6] Kaneko, M. : Poly-Bernoulli numbers, *Jour. Th. Nombre Bordeaux* **9** (1997), 199–206.
- [7] Knuth, D. : Two notes on notation, *Amer. Math. Monthly* **99** (1992), 403–422.

(The Bernoulli numbers are [A027641](#) and [A027642](#). The table in Figure 1 yields sequences [A051714](#) and [A051715](#). Other sequences which mention this paper are [A000367](#), [A002445](#), [A026741](#), [A045896](#), [A051712](#), [A051713](#), [A051716](#), [A051717](#), [A051718](#), [A051719](#), [A051720](#), [A051721](#), [A051722](#), and [A051723](#).)

Received August 7, 2000; published in Journal of Integer Sequences December 12, 2000.

Return to [Journal of Integer Sequences home page](#).
