

## Lecture 15

**Instructor's Comments:** If you did the surveys, you could go over them at the beginning

Handout or Document Camera or Class Exercise

**Fibonacci Sequence Definition:** Define a sequence by  $f_1 = 1$ ,  $f_2 = 1$  and

$$f_n = f_{n-1} + f_{n-2} \quad \text{For all } n \geq 3$$

so  $f_3 = 2$ ,  $f_4 = 3$ ,  $f_5 = 5$ , and so on.

(i) Prove that  $\sum_{r=1}^n f_r^2 = f_n f_{n+1}$  for all  $n \in \mathbb{N}$ .

(ii) Prove that  $f_n < \left(\frac{7}{4}\right)^n$  for all  $n \in \mathbb{N}$ .

**Solution:** We prove only the first one. The second can be found on the Math 135 resources page

<http://www.cemc.uwaterloo.ca/~cbruni/Math135Resources.php>

(i) Base case:  $n = 1$

$$\begin{aligned} \text{LHS} &= \sum_{r=1}^n f_r^2 \\ &= \sum_{r=1}^1 f_r^2 \\ &= f_1^2 \\ &= 1^2 \\ &= 1 \end{aligned}$$

and

$$\text{RHS} = f_n f_{n+1} = f_1 f_2 = (1)(1) = 1 = \text{LHS}$$

(ii) Inductive Hypothesis. Assume that

$$\sum_{r=1}^k f_r^2 = f_k f_{k+1}$$

holds for some  $k \in \mathbb{N}$ .

(iii) Inductive Step. We want to show that

$$\sum_{r=1}^{k+1} f_r^2 = f_{k+1} f_{k+2}.$$

We begin with the left and proceed towards the right

$$\begin{aligned} \text{LHS} &= \sum_{r=1}^{k+1} f_r^2 \\ &= \sum_{r=1}^k f_r^2 + f_{k+1}^2 \\ &= f_k f_{k+1} + f_{k+1}^2 && \text{Induction Hypothesis} \\ &= f_{k+1}(f_k + f_{k+1}) \\ &= f_{k+1} f_{k+2} && \text{By definition of Fibonacci Sequence} \\ &= \text{RHS} \end{aligned}$$

Hence  $\sum_{r=1}^n f_r^2 = f_n f_{n+1}$  for all  $n \in \mathbb{N}$  by the Principle of Mathematical Induction. ■

**Instructor's Comments: This easily is the 20-30 minute mark. Students might struggle with the notation.**

**Definition:** Closed form: “Easy to put into a calculator” (This is not a formal definition!)

**Example:** Find a closed form expression for

$$P_n = \prod_{r=2}^n \left(1 - \frac{1}{r^2}\right)$$

where  $n \geq 2$  and prove it is correct by induction.

**Proof:** We begin with some guessing and napkin (discovery) work.

$$P_2 = \prod_{r=2}^2 \left(1 - \frac{1}{r^2}\right) = \left(1 - \frac{1}{2^2}\right) = 1 - \frac{1}{4} = \frac{3}{4}$$

$$P_3 = \prod_{r=2}^3 \left(1 - \frac{1}{r^2}\right) = \left(1 - \frac{1}{2^2}\right) \left(1 - \frac{1}{3^2}\right) = \left(1 - \frac{1}{4}\right) \left(1 - \frac{1}{9}\right) = \frac{3}{4} \cdot \frac{8}{9} = \frac{2}{3} = \frac{4}{6}$$

$$P_4 = \prod_{r=2}^4 \left(1 - \frac{1}{r^2}\right) = \left(1 - \frac{1}{2^2}\right) \left(1 - \frac{1}{3^2}\right) \left(1 - \frac{1}{4^2}\right) = \left(1 - \frac{1}{4}\right) \left(1 - \frac{1}{9}\right) \left(1 - \frac{1}{16}\right) = \frac{3}{4} \cdot \frac{8}{9} \cdot \frac{15}{16} = \frac{5}{8}$$

Claim:  $P_5 = \frac{6}{10}$  and in general  $P_n = \frac{n+1}{2n}$  for all  $n \geq 2$ . We prove this by induction.

(i) Base case:  $n = 2$

$$P_2 = \prod_{r=2}^2 \left(1 - \frac{1}{r^2}\right) = \left(1 - \frac{1}{2^2}\right) = 1 - \frac{1}{4} = \frac{3}{4} = \frac{n+1}{2n}$$

(ii) Inductive Hypothesis. Assume that  $P(k)$  is true for some  $k \geq 2$  and  $k \in \mathbb{N}$ , that is, assume

$$\prod_{r=2}^k \left(1 - \frac{1}{r^2}\right) = \frac{k+1}{2k}$$

(iii) Inductive Step. We want to show that

$$\prod_{r=2}^{k+1} \left(1 - \frac{1}{r^2}\right) = \frac{(k+1)+1}{2(k+1)} = \frac{k+2}{2k+2}$$

We proceed starting from the left.

$$\begin{aligned} \text{LHS} &= \prod_{r=2}^{k+1} \left(1 - \frac{1}{r^2}\right) \\ &= \prod_{r=2}^k \left(1 - \frac{1}{r^2}\right) \cdot \left(1 - \frac{1}{(k+1)^2}\right) \\ &= \frac{k+1}{2k} \cdot \frac{(k+1)^2 - 1}{(k+1)^2} && \text{Inductive Hypothesis} \\ &= \frac{k+1}{2k} \cdot \frac{k^2 + 2k}{(k+1)^2} \\ &= \frac{k+1}{2k} \cdot \frac{k(k+2)}{(k+1)^2} \\ &= \frac{k+2}{2(k+1)} \\ &= \text{RHS} \end{aligned}$$

Therefore, by the Principle of Mathematical Induction, we have that

$$P_n = \frac{n+1}{2n}$$

for all  $n \geq 2$ . ■

**Instructor's Comments: This is the 50 minute mark.**