

Lecture 16 : Spectral Sparsification

We study a generalization of the cut sparsification problem, called spectral sparsification, and see how this problem can be solved nicely using random sampling and a matrix Chernoff bound.

Spectral sparsification

A graph H is called an ε -cut approximator of G if $(1-\varepsilon)w_G(\delta(S)) \leq w_H(\delta(S)) \leq (1+\varepsilon)w_G(\delta(S))$ for all $S \subseteq V$, where $w_G(\delta(S))$ is the total weight of the edges crossing S .

Benczur and Karger proved that for any G , there exists a $(1+\varepsilon)$ -cut approximator with $O(\frac{n \log n}{\varepsilon^2})$ edges.

Today we will prove a spectral generalization of this result.

Spectral approximator

We say a graph H is an ε -spectral approximator of G if $(1-\varepsilon)L_G \preceq L_H \preceq (1+\varepsilon)L_G$, or

equivalently $(1-\varepsilon)x^T L_G x \leq x^T L_H x \leq (1+\varepsilon)x^T L_G x \quad \forall x \in \mathbb{R}^n$ where n is the number of vertices.

Claim An ε -spectral approximator is an ε -cut approximator.

Proof For $S \subseteq V$, let $x_S \in \mathbb{R}^n$ be the vector with $x_S(i) = 1$ if $i \in S$ and zero otherwise.

Then, $x_S^T L_G x_S = \sum_{ij \in E} w_G(ij) (x_S(i) - x_S(j))^2 = w_G(\delta(S))$ and similarly $x_S^T L_H x_S = w_H(\delta(S))$.

Since H is a $(1+\varepsilon)$ -spectral approximator of G , we have

$$(1-\varepsilon)x_S^T L_G x_S \leq x_S^T L_H x_S \leq (1+\varepsilon)x_S^T L_G x_S \quad \forall S \subseteq V \text{ and thus } (1-\varepsilon)w_G(\delta(S)) \leq w_H(\delta(S)) \leq (1+\varepsilon)w_G(\delta(S)) \quad \forall S \subseteq V. \square$$

The following theorem by Spielman and Srivastava thus generalizes the result of Benczur and Karger.

Theorem Any graph has an ε -spectral approximator with $O(n \log n / \varepsilon^2)$ edges.

Reduction

The spectral sparsification result can be reduced to the following purely linear algebraic result.

Theorem Suppose $v_1, \dots, v_m \in \mathbb{R}^n$ are given with $\sum_{i=1}^m v_i v_i^T = I_n$.

There exist scalars s_1, \dots, s_m with at most $O(n \log n / \varepsilon^2)$ non-zeros such that

$$(1-\varepsilon)I_n \leq \sum_{i=1}^m s_i v_i v_i^T \leq (1+\varepsilon)I_n.$$

We sketch the proof of the reduction of the spectral sparsification result to the above result.

The idea is to apply a linear transformation so that the Laplacian matrix becomes the identity matrix.

Let M be a positive semidefinite matrix with eigen-decomposition $M = \sum_{i=1}^n \lambda_i u_i u_i^T$.

Let M be a positive semidefinite matrix with eigen-decomposition $M = \sum_{i=1}^n \lambda_i u_i u_i^T$.

The pseudo-inverse of M is defined as $M^+ = \sum_{i:\lambda_i > 0} \frac{1}{\lambda_i} u_i u_i^T$, and $M^{1/2} = \sum_{i:\lambda_i > 0} \frac{1}{\sqrt{\lambda_i}} u_i u_i^T$.

Given $L_G = \sum_{e \in E} L_e = \sum_{e \in E} b_e b_e^T$, we consider $I = L_G^{1/2} L_G L_G^{1/2} = \sum_{e \in E} (L_G^{1/2} b_e)(b_e^T L_G^{1/2}) = \sum_{e \in E} s_e v_e v_e^T$, where we define $v_e = L_G^{1/2} b_e \quad \forall e \in E$.

Apply the above theorem gives us s_e with at most $O(n \log n / \varepsilon^2)$ non-zeros so that

$$(1-\varepsilon)I \leq \sum_{e \in E} s_e v_e v_e^T \leq (1+\varepsilon)I.$$

Now, multiplying $L_G^{1/2}$ on the left and right gives us $(1-\varepsilon)L_G \leq \sum_{e \in E} s_e b_e b_e^T \leq (1+\varepsilon)L_G$, so by scaling the weight of each edge by a factor of s_e , we get our spectral sparsifier.

The above "proof" is not precise as we are dealing with the pseudo-inverse (but not the inverse), but the missing details are rather routine and is not the important part of the proof, and so omitted.

Sampling algorithm

Now, our focus is to prove the linear algebraic result, by random sampling.

First, we get some intuition about the condition $\sum_{i=1}^m v_i v_i^T = I_n$, the "isotropy" condition.

When $m=n$, then v_1, \dots, v_n must be an orthonormal basis.

When $m > n$, we can also think of it as an "overcomplete" basis, as we can write any $x \in \mathbb{R}^n$ as

$$x = I_n x = \left(\sum_{i=1}^m v_i v_i^T \right) x = \sum_{i=1}^m \langle x, v_i \rangle v_i.$$

Similarly, for any unit vector $y \in \mathbb{R}^n$, we have $1 = y^T y = y^T I y = y^T \left(\sum_{i=1}^m v_i v_i^T \right) y = \sum_{i=1}^m y^T v_i v_i^T y = \sum_{i=1}^m \langle v_i, y \rangle^2$.

Intuitively, the vectors are "evenly spread out", so that the projection of any direction y to these vectors are the same.

Idea: Given $\sum_{i=1}^m v_i v_i^T = I_n$, we would like to find a small subset of vectors $S \subseteq \{1, \dots, m\}$ and some scaling factors so that $\sum_{i \in S} s_i v_i v_i^T \approx I_n$.

So, the subsets should still be "evenly spread out", with contributions in each direction about the same.

As in the graph sparsification case, uniform sampling won't work. For example, if some v_j has $\|v_j\|=1$, then we must include v_j in the solution, as otherwise that direction will not be covered in the solution and so it won't be a spectral sparsifier. The analogy in the graph sparsification result is that a cut edge must be included in any sparsifier.

So, as in the graph sparsification case, we need to do non-uniform sampling (if we do random sampling).

The idea is similar: for longer vectors, the sampling probability is higher; for shorter vectors, we can be more aggressive in setting the sampling probability to be smaller, and when we

choose them, we reweight the vector so that it has the correct expected value.

More concretely, we sample each vector v_i with probability $\|v_i\|_2^2$, and if it is chosen, we set the scalar $s_i = \frac{1}{\|v_i\|_2^2}$, so that $E[s_i v_i v_i^\top] = \frac{v_i v_i^\top}{\|v_i\|_2^2} \cdot \Pr(v_i \text{ is chosen}) = \frac{v_i v_i^\top}{\|v_i\|_2^2} \cdot \|v_i\|_2^2 = v_i v_i^\top$.

Algorithm

The actual algorithm is basically the same as described above, but we need to repeat this experiment $C = \Theta(\log n)$ times and take the average, so that we can prove concentration.

- Initially, $F \leftarrow \emptyset$, $s \leftarrow 0$, $C = \frac{6 \log n}{\epsilon^2}$

- For $1 \leq t \leq C$ do

For each $e \in E$, with probability $p_e = \|v_e\|_2^2$, update $F \leftarrow F \cup \{e\}$ and $s_e \leftarrow s_e + \frac{1}{C p_e}$.

- Return $\sum_{i \in F} s_i v_i v_i^\top$ as our spectral approximator.

Analysis

There are two steps in the analysis.

One is to show that there are $O(n \log n / \epsilon^2)$ non-zeros scalars, i.e. $|F| = O(n \log n / \epsilon^2)$.

Another is to show that the returned solution is an ϵ -spectral sparsifier.

We first bound the number of nonzero scalars.

Claim With probability at least 0.9, $|F| = O(n \log n / \epsilon^2)$.

Proof The expected value is $E[|F|] = \sum_{i=1}^m \Pr(\text{vector } i \text{ is in } F) = \sum_{i=1}^m (1 - (1-p_i)^C) \leq \sum_{i=1}^m (1 - (1 - Cp_i)) = C \cdot \sum_{i=1}^m p_i$, which can also be seen by a union bound.

Note that $\sum_{i=1}^m p_i = \sum_{i=1}^m \|v_i\|_2^2 = \sum_{i=1}^m v_i^\top v_i = \sum_{i=1}^m \text{tr}(v_i v_i^\top) = \sum_{i=1}^m \text{tr}(v_i v_i^\top) = \text{tr}\left(\sum_{i=1}^m v_i v_i^\top\right) = \text{tr}(I_n) = n$, where

$\text{tr}(A) = \sum_j A_{jj}$ and we use the fact that $\text{tr}(AB) = \text{tr}(BA)$ (or directly check that $v^\top v = \text{tr}(vv^\top)$).

Therefore, $E[|F|] \leq C \sum_{i=1}^m p_i = Cn = 6n \log n / \epsilon^2$. The result follows from Markov's inequality. \square

Matrix Chernoff bound

There is an elegant generalization of the Chernoff-Hoeffding bound to the matrix setting.

Theorem (Tropp) Let X_1, \dots, X_k be independent, $n \times n$ symmetric matrices with $0 \preceq X_i \preceq R I$.

Let $\mu_{\min} I \preceq \sum_{i=1}^k E[X_i] \preceq \mu_{\max} I$. For any $\epsilon \in [0, 1]$,

- $\Pr(\lambda_{\max}(\sum_{i=1}^k X_i) \geq (1+\epsilon)\mu_{\max}) \leq n e^{-\frac{\epsilon^2 - \mu_{\max}}{2R}}$
- $\Pr(\lambda_{\min}(\sum_{i=1}^k X_i) \leq (1-\epsilon)\mu_{\min}) \leq n e^{-\frac{\epsilon^2 - \mu_{\min}}{2R}}$.

Note that it is almost an exact analog of the Chernoff-Hoeffding bound in the scalar case, by using the maximum eigenvalue and minimum eigenvalue to measure the "size" of a matrix.

It says that if we consider the sum of independent random matrices, where each matrix is not too "big/influential", then the sum is concentrated around the expectation in terms of the eigenvalues.

Concentration

The proof that our solution is an ε -spectral sparsifier is a direct application of the matrix Chernoff bound.

The random variables are $X_{i,t} = \begin{cases} \frac{v_i v_i^\top}{C p_i} & \text{with probability } p_i = \|v_i\|_2^2 \\ 0 & \text{otherwise} \end{cases}$, for vector i in iteration t .

Note that the output of the algorithm is $S := \sum_{t=1}^c \sum_{i=1}^m X_{i,t}$.

$$\text{As discussed before, } E[S] = \sum_{t=1}^c \sum_{i=1}^m E[X_{i,t}] = \sum_{t=1}^c \sum_{i=1}^m \frac{v_i v_i^\top}{C p_i} \cdot p_i = \sum_{t=1}^c \sum_{i=1}^m \frac{v_i v_i^\top}{C} = \sum_{i=1}^m v_i v_i^\top = I.$$

So, the expected value is correct, with $\mu_{\max} = \mu_{\min} = 1$ in this problem.

To apply the matrix Chernoff bound, we just need to find a bound for R so that $X_{i,t} \leq R I$.

Note that $X_{i,t} = \frac{v_i v_i^\top}{C p_i} = \frac{v_i v_i^\top}{C \|v_i\|_2^2} = \frac{1}{C} \left(\frac{v_i}{\|v_i\|_2} \right) \left(\frac{v_i}{\|v_i\|_2} \right)^\top$. This is a rank one matrix of a unit vector.

and so the maximum eigenvalue is just $\frac{1}{C}$ (with the only eigenvector being $\frac{v_i}{\|v_i\|_2}$). So, $R = \frac{1}{C}$.

By Tropp's theorem, we get $\Pr(\lambda_{\max}(S) \geq 1 + \varepsilon) \leq n e^{-\varepsilon^2 C / 3} = n e^{-2 \log n} = \frac{1}{n}$, as $C = 6 \log n / \varepsilon^2$.

The lower tail follows similarly.

So, with probability at least $1 - \frac{2}{n}$, we have $\lambda_{\max}(S) \leq 1 + \varepsilon$ and $\lambda_{\min}(S) \geq 1 - \varepsilon$, and so

$(1 - \varepsilon)I \leq S \leq (1 + \varepsilon)I$, proving that our solution S is an ε -spectral sparsifier of I_n .

By a union bound, we know that an ε -spectral sparsifier with $O(n \log n / \varepsilon^2)$ edges exist, and

indeed the random sampling algorithm will succeed with high probability, proving the theorem.

Discussions

There are a few things to discuss about.

- ① By considering this linear algebraic generalization of cut sparsification, we have a clean and arguably simple proof of the result of Benczur and Karger.

A subsequent amazing result by Batson, Spielman and Srivastava proves that every graph has an ε -spectral sparsifier with $O(n/\varepsilon^2)$ edges, which is best possible.

We don't know of an alternative (combinatorial) proof to achieve the same bound even for cut approximator (a special case).

This linear algebraic perspective seems to be the correct way to look at the problem.

② The sampling probability p_e is directly proportional to the effective resistance of the edge e .

Recall that $p_e = \|\mathbf{v}_e\|_2^2 = \|\mathbf{L}_G^+ \mathbf{b}_e\|_2^2 = \mathbf{b}_e^T \mathbf{L}_G^+ \mathbf{b}_e$. Let $\mathbf{e} = \mathbf{u}\mathbf{v}$.

Note that $\mathbf{L}_G^+ \mathbf{b}_e$ is a solution \mathbf{x} to $\mathbf{L}_G \mathbf{x} = \mathbf{b}_e$, which is the potential vector $\vec{\phi}$ of the electrical flow problem when one unit of electrical flow is sent from u to v .

Then, $\mathbf{b}_e^T \mathbf{L}_G^+ \mathbf{b}_e = \mathbf{b}_e^T \vec{\phi} = \phi(u) - \phi(v)$ is just the definition of $R_{\text{eff}}(u, v)$.

So, the sampling algorithm works by sampling each edge with probability proportional to its effective resistance, a somewhat surprising application of this concept.

③ There is a nearly linear time algorithm to estimate the effective resistances of all edges.

The main tools are a near linear time algorithm to solve a Laplacian system of equations (another breakthrough result by Spielman and Teng), and also dimension reduction.

So, we have a near linear time (randomized) algorithm for constructing spectral sparsifiers.

④ The analysis of the random sampling algorithm is tight.

In a complete graph, the effective resistance of every edge is the same, as the graph is symmetric.

So, the random sampling algorithm on a complete graph is just the uniform sampling algorithm.

And by a "coupon collector" argument that it won't work with $O(n \log n / \epsilon^2)$ edges (see HW2).

References

- Spielman and Srivastava, Graph sparsification by effective resistance, 2008.
- Batson, Spielman, and Srivastava, Twice-Ramanujan sparsifiers, 2009.
- Lecture notes by Nick Harvey on Tropp's inequality and spectral sparsification.