

# Improved Analysis of Higher Order Random Walks and Applications\*

Vedat Levi Alev  
University of Waterloo  
Canada

Lap Chi Lau  
University of Waterloo  
Canada

## ABSTRACT

The motivation of this work is to extend the techniques of higher order random walks on simplicial complexes to analyze mixing times of Markov chains for combinatorial problems. Our main result is a sharp upper bound on the second eigenvalue of the down-up walk on a pure simplicial complex, in terms of the second eigenvalues of its links. We show some applications of this result in analyzing mixing times of Markov chains, including sampling independent sets of a graph and sampling common independent sets of two partition matroids.

## CCS CONCEPTS

- **Theory of computation** → **Random walks and Markov chains**;
- **Mathematics of computing** → *Spectra of graphs*.

## KEYWORDS

spectral gap, Markov chains, random sampling, high dimensional expansion

## ACM Reference Format:

Vedat Levi Alev and Lap Chi Lau. 2020. Improved Analysis of Higher Order Random Walks and Applications. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing (STOC '20)*, June 22–26, 2020, Chicago, IL, USA. ACM, New York, NY, USA, 14 pages. <https://doi.org/10.1145/3357713.3384317>

## 1 INTRODUCTION

Consider the following random walks [KM17, DK17, KO18, DDFH18] defined<sup>1</sup> on a simplicial complex  $X$ . Initially, the random walk starts from an arbitrary face  $\alpha_1$  of dimension  $k$  in  $X$ .

- **DOWN-UP WALK:** In each step  $t \geq 1$ , we choose a uniform random element  $i \in \alpha_t$  and delete  $i$  from  $\alpha_t$ , and set  $\alpha_{t+1}$  to be a uniform random face of dimension  $k$  in  $X$  that contains  $\alpha_t \setminus \{i\}$ . This is called the  $k$ -th down-up walk of  $X$ , and its transition matrix is denoted by  $P_k^\nabla$ .

\*The submitted full paper is available at arXiv:2001.02827.

<sup>1</sup>All the definitions in the introduction will be formally defined again in a more general setting in Section 2.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than the author(s) must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from [permissions@acm.org](mailto:permissions@acm.org).  
STOC '20, June 22–26, 2020, Chicago, IL, USA

© 2020 Copyright held by the owner/author(s). Publication rights licensed to ACM.  
ACM ISBN 978-1-4503-6979-4/20/06...\$15.00  
<https://doi.org/10.1145/3357713.3384317>

- **UP-DOWN WALK:** In each step  $t \geq 1$ , we choose a uniform random face  $\beta$  of dimension  $k + 1$  in  $X$  that contains  $\alpha_t$ , and choose a uniform random element  $i \in \beta$  and set  $\alpha_{t+1} = \beta \setminus \{i\}$ . This is called the  $k$ -th up-down walk of  $X$ , and its transition matrix is denoted by  $P_k^\Delta$ .

The stationary distribution of these random walks is the uniform distribution on the faces of dimension  $k$  in the simplicial complex  $X$ . The question of interest is the mixing time of these random walks, i.e. the number of steps  $t$  required for the distribution of  $\alpha_t$  to be close to the uniform distribution.

A graph is a simplicial complex of dimension 1. The transition matrix of the lazy random walk on a graph is  $P_0^\Delta$ . Fundamental results in spectral graph theory state that (i) the mixing time of the lazy random walk is small, if and only if (ii) the second eigenvalue of  $P_0^\Delta$  is small, if and only if (iii) the graph is an expander graph. See [HLW06, WLP09] for surveys on this topic.

Since the theory of expander graphs has many applications, there are various motivations in generalizing these results for graphs to simplicial complexes. Several definitions of high-dimensional expanders have been studied in the literature (e.g. [LM06, Gro10, PRT16, DKW16, KM17, Opp18]), and these results have found interesting applications in discrete geometry, complexity theory, coding theory, and property testing (e.g. [LM06, MW09, FGL<sup>+</sup>11, KL14, EK16, KM16, KKL16, DK17, DHK<sup>+</sup>19]).

**Local Spectral Expanders.** In this paper, we consider the definition of  $\gamma$ -local-spectral expanders developed in [KM17, DK17, KO18, Opp18, DDFH18] for the study of random walks on simplicial complexes. The local structures of a simplicial complex are described by its links. The link  $X_\alpha$  of a face  $\alpha \in X$  is defined as the simplicial complex  $X_\alpha = \{\beta \setminus \alpha : \beta \in X, \beta \supset \alpha\}$ . The graph  $G_\alpha = (V_\alpha, E_\alpha)$  of the link  $X_\alpha$  is defined as follows: (i) each vertex  $i$  in  $V_\alpha$  corresponds to a singleton  $\{i\}$  in  $X_\alpha$ , (ii) two vertices  $i, j \in V_\alpha$  have an edge in  $E_\alpha$  if and only if  $\{i, j\}$  is contained in some face of  $X_\alpha$ , (iii) the weight  $w_{ij}$  of an edge  $ij \in E_\alpha$  is proportional to the number of maximal faces in  $X_\alpha$  that contains  $\{i, j\}$ .

Informally, a simplicial complex  $X$  is a  $\gamma$ -local-spectral expander if  $G_\alpha$  is an expander graph for every  $\alpha \in X$ . In the following, we say  $X$  is a pure simplicial complex if every maximal face of  $X$  is of the same dimension, and we call this the dimension of  $X$ .

**Definition 1.1** ( $\gamma$ -local-spectral expanders [Opp18, KO18]). A  $d$ -dimensional pure simplicial complex  $X$  is a  $\gamma$ -local-spectral expander if  $\lambda_2(G_\alpha) \leq \gamma$  for every face  $\alpha \in X$  of dimension up to  $d - 2$ , where  $\lambda_2(G_\alpha)$  denotes the second largest eigenvalue of the random walk matrix of  $G_\alpha$  (where the transition probabilities are proportional to the edge weights).

**Kaufman-Oppenheim Theorem.** Kaufman and Oppenheim [KO18] proved that the  $k$ -th down-up walk and the  $(k - 1)$ -th up-down walk

have a non-trivial spectral gap as long as the simplicial complex is a  $\gamma$ -local-spectral expander for  $\gamma < 2/k^2$ .

**Theorem 1.2** ([KO18]). *Let  $X$  be a pure  $d$ -dimensional simplicial complex. Suppose  $X$  is a  $\gamma$ -local-spectral expander. Then, for every  $0 \leq k \leq d$ ,*

$$\lambda_2(P_k^\nabla) = \lambda_2(P_{k-1}^\Delta) \leq 1 - \frac{1}{k+1} + \frac{k\gamma}{2},$$

**Theorem 1.2** states that the spectral gap of  $P_k^\nabla$  is at least  $g := 1 - \lambda_2(P_k^\nabla) \geq \frac{1}{k+1} - \frac{k\gamma}{2}$ , which implies by a standard argument (see **Theorem 2.8**) that the mixing time of these walks is at most  $O(\frac{(k+1)\log(n)}{g})$  where  $n$  is the size of the ground set of  $X$ . For example, if  $\gamma \leq 0$ , then the mixing time of  $P_k^\nabla$  is at most  $O(k^2 \log(n))$ .

**Theorem 1.2** can also be used to bound the spectral gap of certain “longer” random walks on simplicial complexes (see **Corollary 1.11** and **Section 1.2.5**). Dinur and Kaufman [DK17] use these results with the Ramanujan complexes of [LSV05] to construct efficient agreement testers, which have applications to PCP constructions. Recently, these ideas have also found applications in coding theory [DHK<sup>+</sup>19].

*Oppenheim’s Trickling Down Theorem.* Kaufman-Oppenheim **Theorem 1.2** provides a way to bound the mixing time of the down-up walks and up-down walks. To apply the theorem, however, one needs to check that  $\lambda_2(G_\alpha) \leq \gamma$  for every face  $\alpha \in X$  of dimension at most  $d - 2$ . This is not an easy task. There are exponentially many graphs  $G_\alpha$  to check, and these graphs are defined implicitly where computing the edge weights involve non-trivial counting problems. A very useful result by Oppenheim [Opp18] makes this task easier, by relating the second eigenvalue of the graph of a lower-dimensional link to that of a higher-dimensional link.

**Theorem 1.3** ([Opp18]). *Let  $X$  be a pure  $d$ -dimensional simplicial complex. Suppose  $\lambda_2(G_\beta) \leq \gamma \leq \frac{1}{2}$  for every face  $\beta$  of dimension  $k$ , and  $G_\alpha$  is connected for every face  $\alpha$  of dimension  $k - 1$ . Then, for every face  $\alpha$  of dimension  $k - 1$ , it holds that*

$$\lambda_2(G_\alpha) \leq \frac{\gamma}{1 - \gamma}.$$

Applying this theorem inductively, we can reduce the problem of bounding  $\lambda_2(G_\alpha)$  for every  $\alpha$  to bounding  $\lambda_2(G_\beta)$  for only those faces  $\beta$  of highest dimension.

**Corollary 1.4** ([Opp18]). *Let  $X$  be a pure  $d$ -dimensional simplicial complex. Suppose  $\lambda_2(G_\beta) \leq \gamma \leq \frac{1}{d}$  for every face  $\beta$  of dimension  $d - 2$ , and  $G_\alpha$  is connected for every face  $\alpha$ . Then, for every  $k \leq d - 2$ , and for every face  $\alpha$  of dimension  $k$ , it holds that*

$$\lambda_2(G_\alpha) \leq \frac{\gamma}{1 - (d - 2 - k)\gamma}.$$

**Corollary 1.4** is useful for two reasons: First, note that the weight of every edge in  $G_\beta$  for face  $\beta$  of dimension  $d - 2$  is either zero or one, which makes the task of bounding its second eigenvalue more tractable. Second, if one can prove that  $\lambda_2(G_\beta) = O(\frac{1}{d^2})$  for every face  $\beta$  of dimension  $d - 2$  and  $G_\alpha$  is connected for every face  $\alpha$ , then one can conclude that  $\lambda_2(G_\alpha) = O(\frac{1}{d^2})$  for every face  $\alpha$  and hence the simplicial complex is a  $O(\frac{1}{d^2})$ -local-spectral expander. So, the reduction of Oppenheim is basically lossless in the regime where Kaufman-Oppenheim’s **Theorem 1.2** applies.

*Analyzing Mixing Times of Markov Chains.* Recently, Anari, Liu, Oveis Gharan, and Vinzant [ALOV19] found a striking application of **Theorem 1.2** and **Corollary 1.4** in proving the matroid expansion conjecture of Mihail and Vazirani [MV87], answering a long standing open question in Markov chain Monte Carlo methods.

To illustrate their result, consider the special case of sampling a random spanning tree from a graph  $G = (V, E)$ . Let  $X$  be the simplicial complex where the ground set is  $E$  and each acyclic subgraph of  $G$  is a face of  $X$ . Then  $X$  is a pure  $d$ -dimensional simplicial complex, where  $d = |V| - 2$  and the spanning trees of  $G$  are the maximal faces of  $X$ . Note that  $P_d^\nabla$  in  $X$  is exactly the natural Markov chain on the spanning trees of  $G$ , where in each step we delete a uniformly random edge  $e$  from the current spanning tree  $T$  and add a uniformly random edge  $f$  so that  $T - e + f$  is a spanning tree. So, the problem of proving the Markov chain on spanning trees is fast mixing is equivalent to upper bounding  $\lambda_2(P_d^\nabla)$  of the simplicial complex  $X$ .

Using the nice structures of matroids, Anari, Liu, Oveis Gharan, and Vinzant [ALOV19] showed that the graph  $G_\beta$  is a complete multi-partite graph for every face  $\beta$  of dimension  $d - 2$ , and this implies that  $\lambda_2(G_\alpha) \leq 0$  for every face  $\beta$  of dimension  $d - 2$ . Thus, it follows from Oppenheim’s **Corollary 1.4** that  $\lambda_2(G_\alpha) \leq 0$  for every face  $\alpha$ .<sup>2</sup> Then Kaufman-Oppenheim’s **Theorem 1.2** implies that  $\lambda_2(P_d^\nabla) \leq 1 - \frac{1}{d+1}$ , and thus the mixing time of the Markov chain of sampling matroid bases is at most  $O(d^2 \log n)$ . This provides the first FPRAS for counting the number of matroid bases, and also proves that the basis exchange graph of a matroid is an expander graph.

The proof of the matroid expansion conjecture shows that the techniques developed in higher order random walks provide a new simplicial complex approach to analyze mixing times of Markov chains. It is thus natural to investigate whether this approach can be extended to other problems. Here we would like to discuss some limitations of the current techniques. It can be shown that  $\lambda_2(G_\beta) \leq 0$  only if  $G_\beta$  is a complete multi-partite graph [God] and more generally a 0-local-expander is a weighted matroid complex [BH19], and so the same analysis as in [ALOV19] only works for matroids. Note that Kaufman-Oppenheim **Theorem 1.2** only applies when  $\lambda_2(G_\alpha) \leq O(\frac{1}{d^2})$  for every face  $\alpha$  up to dimension  $d - 2$ . For many problems that we have considered, it does not hold that  $\lambda_2(G_\beta) \leq O(\frac{1}{d^2})$  even when restricted to faces  $\beta$  of dimension  $d - 2$ .

## 1.1 Main Result

The main motivation of this work is to extend this simplicial complex approach to analyze mixing times of more general Markov chains. Our main result is the following improved eigenvalue bound for higher order random walks.

**Theorem 1.5.** *Let  $X$  be a pure  $d$ -dimensional simplicial complex. Define*

$$\gamma_j := \max_{\alpha} \{\lambda_2(G_\alpha) : \alpha \in X \text{ and } \alpha \text{ is of dimension } j\},$$

<sup>2</sup>The result that every matroid complex is a 0-local-spectral expander was also proved by Huh and Wang [HW17], using techniques from Hodge theory for matroids [AHK19] instead of Oppenheim’s theorem.

For any  $0 \leq k \leq d$ ,

$$\lambda_2(\mathbb{P}_k^\nabla) = \lambda_2(\mathbb{P}_{k-1}^\Delta) \leq 1 - \frac{1}{k+1} \prod_{j=-1}^{k-2} (1 - \gamma_j).$$

The following are some remarks about [Theorem 1.5](#).

- (1) A basic result is that a simplicial complex  $X$  is gallery connected (i.e.  $\lambda_2(\mathbb{P}_d^\nabla) < 1$ ) if  $G_\alpha$  is connected (i.e.  $\lambda_2(G_\alpha) < 1$ ) for every face  $\alpha$  of dimension up to  $d - 2$ . [Theorem 1.5](#) provides a quantitative generalization of this result.
- (2) A corollary of [Theorem 1.5](#) is that the spectral gap  $1 - \lambda_2(\mathbb{P}_k^\nabla)$  of the  $k$ -th down-up walk is at least  $\Omega(1/k)$  if  $X$  is a  $O(\frac{1}{k})$ -local-spectral expander. This is an improvement of [Theorem 1.2](#) where it requires the simplicial complex  $X$  to be a  $O(\frac{1}{k^2})$ -local-spectral expander to conclude that  $\mathbb{P}_k^\nabla$  has a non-zero spectral gap.
- (3) It can be shown that the spectral gap  $1 - \lambda_2(\mathbb{P}_k^\nabla)$  of the  $k$ -th down-up walk is at most  $O(\frac{1}{k})$  for any simplicial complex (see [Proposition 3.3](#)), so [Theorem 1.5](#) shows that any  $O(\frac{1}{k})$ -local-spectral expander has the optimal spectral gap for the  $k$ -th down-up walk up to a constant factor.
- (4) The refinement of having a different bound  $\gamma_j$  for links of different dimension is very useful for analyzing Markov chains. We will see some applications in [Section 4](#).
- (5) [Theorem 1.5](#) can be used to provide a tighter bound on the spectral gap of certain “longer” random walks (see [Corollary 1.12](#)) which were known to be useful in coding theory and agreement testing (see [Section 1.2.5](#)).

Combined with [Oppenheim’s Theorem 1.3](#), [Theorem 1.5](#) provides the following bound for the second eigenvalue of higher order random walks in a black box fashion. See [Section 3](#) for the proof.

**Corollary 1.6.** *Let  $X$  be a pure  $d$ -dimensional simplicial complex. For any  $0 \leq k \leq d$ , suppose  $\gamma_{k-2} \leq \frac{1}{k+1}$  and  $G_\alpha$  is connected for every face  $\alpha$  up to dimension  $k - 2$ , then*

$$\lambda_2(\mathbb{P}_k^\nabla) = \lambda_2(\mathbb{P}_{k-1}^\Delta) \leq 1 - \frac{1}{(k+1)^2}.$$

This provides a convenient way to bound the mixing time of Markov chains. Recall that the edge weights in  $G_\beta$  for face  $\beta$  of dimension  $d - 2$  are either zero or one, and so it is easier to bound their second eigenvalue. [Corollary 1.6](#) states that as long as we can prove  $\lambda_2(G_\beta) \leq 1/(d+1)$  for these unweighted graphs in the highest dimension, then we can conclude that  $\mathbb{P}_d^\nabla$  is fast mixing.

## 1.2 Applications

We present several applications of [Theorem 1.5](#) and [Corollary 1.6](#), in analyzing mixing times of Markov chains ([Section 1.2.1](#), [Section 1.2.2](#), [Section 1.2.3](#)), in analyzing constructions of high-dimensional expanders ([Section 1.2.4](#)), and in analyzing longer random walks ([Section 1.2.5](#)).

**1.2.1 Sampling Independent Sets of Fixed Size.** One of the most natural simplicial complexes to consider is the independent set complex of a graph [[Mes01](#), [AB06](#)]. Let  $G = (V, E)$  be a graph. The independent set complex  $I_{G,k}$  has the vertex set  $V$  as the ground set, and a subset  $S \subset V$  is a face in  $X$  if and only if  $S$  is an independent set in  $G$  with  $|S| \leq k$ .

We are interested in bounding  $\lambda_2(\mathbb{P}_{k-1}^\nabla)$  for this simplicial complex  $X$ . The  $(k - 1)$ -th down-up walk corresponds to a natural Markov chain on sampling independent sets of size  $k$ . Initially, the random walk starts from an arbitrary independent set  $S_1$  of size  $k$ . In each step  $t \geq 1$ , we choose a uniform random vertex  $u \in S_t$  and delete it from  $S_t$ , and we choose a uniform random vertex  $v$  so that  $S_t - u + v$  is still an independent set of size  $k$  and set  $S_{t+1} := S_t - u + v$ . This Markov chain is known to mix in polynomial time for  $k \leq \frac{|V|}{2\Delta+1}$  where  $\Delta$  is the maximum degree of  $G$ , by using the path coupling technique [[BD97](#), [MU05](#)]. We prove a more refined result using the simplicial complex approach.

**Theorem 1.7.** *Let  $G = (V, E)$  be a graph with maximum degree  $\Delta$ . Let  $\mathbb{P}_{k-1}^\nabla$  be the  $(k - 1)$ -th down-up walk on the simplicial complex  $I_{G,k}$ . Let  $A_G$  be the adjacency matrix of  $G$ .*

$$\text{If } k \leq \frac{|V|}{\Delta + |\lambda_{\min}(A_G)|}, \text{ then } \lambda_2(\mathbb{P}_{k-1}^\nabla) \leq 1 - \frac{1}{k^2}.$$

It is well-known that  $|\lambda_{\min}(A_G)| \leq \Delta$  for a graph with maximum degree  $\Delta$ , and so [Theorem 1.7](#) recovers the previous result that the Markov chain is fast mixing if  $k \leq \frac{|V|}{2\Delta}$ . There are various graph classes with  $|\lambda_{\min}(A_G)|$  smaller than  $\Delta$ , and [Theorem 1.7](#) allows us to sample larger independent sets. For example, it is known that  $|\lambda_{\min}(A_G)| \leq O(\sqrt{\Delta})$  for planar graphs and more generally for graphs with bounded arboricity [[Hay06](#)], and also for random graphs and more generally for two-sided expander graphs [[HLW06](#)].

**1.2.2 Sampling Common Independent Sets in Two Partition Matroids.** A matroid  $M = (E, \mathcal{I})$  on the ground set  $E$  with the set of independent sets  $\mathcal{I} \subset 2^E$  is a combinatorial object satisfying the following properties:

- (containment property) if  $S \in \mathcal{I}$  and  $T \subset S$ , then  $T \in \mathcal{I}$ ,
- (extension property) if  $S, T \in \mathcal{I}$  such that  $|S| > |T|$  then there is some  $x \in S \setminus T$  such that  $\{x\} \cup T \in \mathcal{I}$ .

A partition matroid is the special case where the ground set  $E$  is partitioned into disjoint blocks  $B_1, \dots, B_l \subseteq E$  with parameters  $0 \leq d_i \leq |B_i|$  for  $1 \leq i \leq l$ , and a subset  $S$  is in  $\mathcal{I}$  if and only if  $|S \cap B_i| \leq d_i$  for  $1 \leq i \leq l$ .

The intersection of two matroids  $M_1 = (E, \mathcal{I}_1)$  and  $M_2 = (E, \mathcal{I}_2)$  over the same ground set  $E$  can be used to formulate various interesting combinatorial optimization problems [[Sch03](#)]. We are interested in the problem of sampling a uniform random common independent set of size  $k$ , i.e. a random subset  $F \in \mathcal{I}_1 \cap \mathcal{I}_2$  with  $|F| = k$ .

Matroids naturally correspond to simplicial complexes. We let  $C_{M_1, M_2, k}$  be the matroid intersection complex with ground set  $E$ , where a subset  $F \subset E$  is a face in  $C_{M_1, M_2, k}$  if and only if  $F \in \mathcal{I}_1 \cap \mathcal{I}_2$  and  $|F| \leq k$ . The  $(k - 1)$ -th down-up walk of this complex corresponds to a natural Markov chain on sampling common independent sets of  $M_1$  and  $M_2$  of size  $k$ . We show that this Markov chain is fast mixing for  $k$  up to one third the size of a maximum common independent set, when  $M_1$  and  $M_2$  are partition matroids and there are no two elements belonging to the same block in both matroids (i.e. there are no two elements  $x, y$  such that  $x$  and  $y$  are in the same block in  $M_1$  and also in the same block in  $M_2$ ).

**Theorem 1.8.** *Let  $M_1 = (E, \mathcal{I}_1)$  and  $M_2 = (E, \mathcal{I}_2)$  be two given partition matroids with a common independent set of size  $r$  and no two elements belonging to the same block in both matroids. If  $k \leq r/3$ , then*

$$\lambda_2(P_{k-1}^\nabla) \leq 1 - \frac{1}{k^2},$$

where  $P_{k-1}^\nabla$  is the  $(k-1)$ -th down-up walk on the matroid intersection complex  $C_{M_1, M_2, k}$ .

Sampling a common independent set of two partition matroids can be reduced to sampling a random matching in a bipartite graph. In this regime, stronger results are known to exist [JS89, JSV04] whose analysis is based on the canonical paths method while our analysis provides an alternative approach using a spectral method.

The proof of [Theorem 1.8](#) reveals an interesting property of the links of the simplicial complex  $C_{M_1, M_2, k}$ . For any face  $\beta$  of dimension  $k-3$ , we show that the graph  $G_\beta$  is the complement of the line graph of a bipartite graph. We note that this holds for any two matroids, not just for partition matroids. By the additional assumptions that the two matroids are partition matroids and there are no two elements in the same block in both matroids, the graph  $G_\beta$  is the line graph of a *simple* bipartite graph. Using the fact that the adjacency matrix of the line graph of a simple graph has minimum eigenvalue at least  $-2$ , we prove that  $\lambda_2(G_\beta) \leq \frac{1}{k}$  as long as  $k \leq \frac{r}{3}$ . We can then use [Corollary 1.6](#) to conclude [Theorem 1.8](#).

**1.2.3 Sampling Independent Sets from Hardcore Distributions.** Very recently, Anari, Liu, and Oveis Gharan [ALO] use [Theorem 1.5](#) to prove a strong result about sampling independent sets from the hardcore distribution. Given a graph  $G = (V, E)$  and a parameter  $\lambda > 0$ , the problem is to sample an independent set  $S$  with probability  $\frac{\lambda^{|S|}}{Z_G(\lambda)}$  where  $Z_G(\lambda) := \sum_{S \subset V: S \text{ independent}} \lambda^{|S|}$  is the partition function. An important work of Weitz [Wei06] gave a deterministic fully polynomial time approximation scheme to estimate  $Z_G(\lambda)$  for  $\lambda$  up to the “uniqueness threshold”, but the exponent of the runtime depends on the maximum degree  $\Delta$  of  $G$ . It is conjectured that the natural Markov chain for sampling independent sets mixes in polynomial time up to the uniqueness threshold. Anari, Liu, and Oveis Gharan prove this conjecture and obtain a polynomial time algorithm to estimate  $Z_G(\lambda)$  up to the uniqueness threshold for any graph (even with unbounded maximum degree). They consider a pure  $n$ -dimensional simplicial complex for sampling independent sets, and prove that  $\gamma_j = \Theta(\frac{1}{n-j})$  for  $0 \leq j \leq n-2$  by using the techniques from correlation decay. Then it follows from [Theorem 1.5](#) that the Markov chain is fast mixing. Note that it is crucial to have a different bound  $\gamma_j$  for links of different dimension in [Theorem 1.5](#), so even when  $\gamma_{n-2} = \Theta(1)$  it is still possible to conclude fast mixing.

**1.2.4 Combinatorial Constructions of High Dimensional Expanders.** Recently, Liu, Mohanty, and Yang [LMY19] presented an interesting combinatorial construction of a sparse simplicial complex where all higher order random walks have a constant spectral gap. Their construction is by taking a certain tensor product of a graph  $G$  on  $n$  vertices and a small  $H$ -dimensional complete simplicial complex  $\mathcal{B}$  on  $s$  vertices.

**Theorem 1.9** ([LMY19]). *Let  $G$  be a  $T$ -regular triangle free graph on  $n$  vertices. There is an explicit family  $(X^{(s, H, G)})_{H \geq 1, s \geq H+1}$  of simplicial complexes, satisfying the following properties:*

- (1)  $X^{(s, H, G)}$  is a pure  $H$ -dimensional simplicial complex with  $\Theta(n)$  maximal faces.
- (2) The spectral gap of the graphs of  $j$  dimensional links of the complex  $X^{(s, H, G)}$  satisfies

$$1 - \gamma_j \geq \begin{cases} \frac{1}{2} & \text{if } j \in [0, H-2], \\ \left(\frac{1}{2} - \frac{1}{2(T^2 H+1)}\right)(1 - \sigma_2(G)) & \text{if } j = -1, \end{cases}$$

where  $\sigma_2(G)$  is the second largest eigenvalue of the normalized adjacency matrix of  $G$ .

- (3) For any  $-1 \leq j \leq H-2$ ,

$$\lambda_2(P_{j+1}^\nabla) = \lambda_2(P_j^\Delta) \leq 1 - \Omega\left(\frac{1 - \sigma_2(G)}{T^2 \cdot j^2 \cdot (s-j) \cdot 2^j}\right),$$

The main technical part of their proof is in establishing Item (3) in [Theorem 1.9](#). They use the special structures of their construction and the decomposition technique from [JST<sup>+</sup>04] to bound the spectral gap of the higher order random walks. The authors ask the question whether the spectral property in Item (2) alone is enough to prove the fast mixing result in Item (3). Note that Kaufman-Oppenheim’s [Theorem 1.2](#) does not apply in this regime.

Using [Theorem 1.5](#), we answer their question affirmatively, by deriving Item (3) from Item (2) in a black box fashion. This slightly improves their bound and considerably simplifies their analysis.

**Corollary 1.10.** *Let  $X := X^{(s, H, G)}$  be a complex from [Theorem 1.9](#) satisfying Item (2). For any  $-1 \leq j \leq H-2$ ,*

$$\lambda_2(P_{j+1}^\nabla) = \lambda_2(P_j^\Delta) \leq 1 - \Omega\left(\frac{1 - \sigma_2(G)}{j \cdot 2^j}\right).$$

**1.2.5 Longer Random Walks and Other Applications.** Consider the following generalization of the up-down walk where we take “longer” steps. Initially, the random walk starts from an arbitrary  $\alpha_1$  face of dimension  $a$  in  $X$ . In each step  $t \geq 1$ , we sample a uniformly random face  $\beta$  of dimension  $b > a$  that contains  $\alpha_t$ , and set  $\alpha_{t+1}$  to be a uniformly random subset of  $\beta$  of dimension  $a$ . We call this the  $a$ -th up-down walk through the  $b$ -th dimension, and denote its transition matrix by  $P_{a,b}^\Delta$ . The  $k$ -th up-down walk defined before is the special case  $P_{k, k+1}^\Delta$ . Dinur and Kaufman [DK17] derived the following result about  $P_{a,b}^\Delta$  from the result about the ordinary up-down walks.

**Corollary 1.11** ([DK17]). *Let  $X$  be a  $d$ -dimensional pure simplicial complex. If  $X$  is a  $\gamma$ -local-spectral expander, then for any  $0 \leq a < b \leq d-1$ ,*

$$\lambda_2(P_{a,b}^\Delta) \leq \frac{a+1}{b+1} + O(a(b-a)\gamma).$$

Using [Theorem 1.5](#), we obtain the following improved bound. See [Section 3](#) for the proof.

**Corollary 1.12.** *Let  $X$  be a  $d$ -dimensional pure simplicial complex. If  $X$  is a  $\gamma$ -local-spectral expander, then for any  $0 \leq a < b \leq d-1$ ,*

$$\lambda_2(P_{a,b}^\Delta) \leq (1 + \gamma)^{b-a} \cdot \frac{a+1}{b+1}.$$

In particular, if  $\gamma \leq \frac{\epsilon}{b-a}$  for some  $0 \leq \epsilon \leq 1$ , then  $\lambda_2(P_{a,b}^\Delta) \leq e^\epsilon \cdot \frac{a+1}{b+1}$ .

Whereas the bound from [Corollary 1.11](#) requires  $\gamma = O(\frac{1}{b \cdot (b-a)})$  to give a nontrivial upper bound on the second eigenvalue of  $P_{a,b}^\Delta$ ,



**Corollary 1.12** only requires  $\gamma \leq O(\frac{1}{b-a})$  to give a comparable bound.

**Corollary 1.11** has found applications in agreement testing and coding theory [DK17, DHK<sup>+</sup>19, AJQ<sup>+</sup>20]. We believe that **Corollary 1.12** can be of independent interest because of those applications. One potential application would be in constructing double samplers from Ramanujan complexes under a weaker expansion assumption [DK17].

### 1.3 Related Work

*Higher Order Random Walks and Applications.* Our work follows a sequence of works [KM17, DK17, Opp18, KO18, DDFH18] which use the spectral properties of the links of simplicial complexes to analyze higher order random walks. Higher order random walks on simplicial complexes were first introduced by Kaufman and Mass [KM17]. They formulated related but more combinatorial notions of skeleton expansion and colorful expansion to establish fast mixing of higher order random walks. Dinur and Kaufman [DK17] introduced the definition of two-sided  $\gamma$ -local-spectral expanders, which is similar to **Definition 1.1** but requires all but the first eigenvalue to have absolute value at most  $\gamma$  (i.e. it also controls the negative eigenvalues). They used this stronger assumption to prove a similar theorem as in **Theorem 1.2**, and applied it to construct efficient agreement tester with applications to PCP constructions. The one-sided  $\gamma$ -local-expander in **Definition 1.1** was first studied by Oppenheim [Opp18], where he proved **Theorem 1.3**. Then, Kaufman and Oppenheim [KO18] strengthened the result in [DK17] and prove **Theorem 1.2**.

Dikstein, Dinur, Filmus and Harsha [DDFH18] studied an alternative definition of high dimensional expanders, based on the operator norm of the difference between the (non-lazy) up-down and down-up operators. Using this definition, they show that it is possible to approximately characterize all the eigenvalues and eigenvectors of higher order random walks. Their techniques were used in [AJT19] to analyze the “swap walks” on high dimensional expanders, with applications in designing good approximation algorithms for solving constraint satisfaction problems on high-dimensional expanders. Independently, the same “swap walks” were also studied by [DD19] under the name “complement walks”, where applications in agreement testing were given.

The results in higher order random walks have also found applications in coding theory. The double samplers in [DK17] are used in [DHK<sup>+</sup>19] to design an efficient algorithm to decode direct product codes over high dimensional expanders. The swap walks in [AJT19] are used in [AJQ<sup>+</sup>20] to recover the same result and also to design an efficient algorithm to decode direct sum codes over high dimensional expanders.

*Analyzing Mixing Times of Markov Chains.* Mixing time of Markov chains is an extensively studied topic with various applications (see e.g. [WLP09, MT05]). There are several well-developed approaches to bound the mixing time of a Markov chain. Perhaps the most widely used approach is the coupling method (e.g. [Ald83, BD97]), which has applications in sampling graph colorings (e.g. [Jer95, Vig00]) and many other problems (see [WLP09]). The canonical path (or more generally multicommodity flow) method developed

in [JS89, Sin92, Sin93] was used in the important problem of sampling perfect matchings in bipartite graphs [JS89, JSV04] and other problems including sampling matroid bases [FM92]. Geometric methods are used in the important problem of sampling a random point in a convex body [DFK91, LV06]. Analytical methods such as (modified) log-Sobolev inequalities and Nash inequalities [DSC<sup>+</sup>96, BT06] are useful in proving sharp bounds on mixing time, e.g. a recent paper [CGM19] used a modified log-Sobolev inequality to prove optimal mixing time of the natural Markov chain on sampling matroid bases.

The simplicial complex approach studied in this paper is quite different from the above approaches. It is linear algebraic and designed to bound the second eigenvalue directly using ideas from simplicial complexes. On the other hand, the coupling method is probabilistic and designed to compare two random processes, while the canonical path method and the geometric method are designed to bound the underlying expansion of the graph or the geometric object. The analytical methods are more difficult to apply and are not as widely applicable, but when they work they could be used to prove very sharp results.

## 2 PRELIMINARIES

### 2.1 Linear Algebra

*Vectors and Inner-Products.* Bold faces will be used for scalar functions, i.e.  $\mathbf{f} \in \mathbb{R}^V$ . The notation  $\mathbf{1}_V \in \mathbb{R}^V$  will be reserved for the all-one vector in  $\mathbb{R}^V$ ; the subscript may be omitted when the vector space  $\mathbb{R}^V$  is clear from the context.

Throughout this text, we use  $\Pi \in \mathbb{R}^V$  to denote various probability distributions, i.e.  $\sum_{x \in V} \Pi(x) = 1$  and  $\Pi(x) \geq 0$  for  $x \in V$ . Given  $\mathbf{f}, \mathbf{g} \in \mathbb{R}^V$ , we use the notations  $\langle \mathbf{f}, \mathbf{g} \rangle_\Pi$  and  $\|\mathbf{f}\|_\Pi$  to denote the inner-product and the norm with respect to the distribution  $\Pi$ , i.e.

$$\langle \mathbf{f}, \mathbf{g} \rangle_\Pi = \sum_x \Pi(x) \mathbf{f}(x) \mathbf{g}(x) \quad \text{and} \quad \|\mathbf{f}\|_\Pi^2 = \langle \mathbf{f}, \mathbf{f} \rangle_\Pi.$$

We reserve  $\langle \mathbf{f}, \mathbf{g} \rangle = \sum_x \mathbf{f}(x) \mathbf{g}(x)$  for the standard inner-product. Given  $\mathbf{f} \in \mathbb{R}^V$ , we write  $\|\mathbf{f}\|_{\ell_1} = \sum_{x \in V} |\mathbf{f}(x)|$  for its  $\ell_1$ -norm, and  $\|\mathbf{f}\|_{\ell_2} = (\sum_{x \in V} \mathbf{f}(x)^2)^{\frac{1}{2}}$  for its  $\ell_2$ -norm.

*Matrices and Eigenvalues.* Serif faces will be used for matrices, i.e.  $A \in \mathbb{R}^{V \times V}$ . Let  $G = (V, E)$  be an edge-weighted undirected graph with a weight  $w_e > 0$  on each edge  $e \in E$ . The adjacency matrix of  $G$  is denoted by  $A_G \in \mathbb{R}^{V \times V}$  with  $A_G(u, v) = w_{uv}$  for  $uv \in E$  and  $A_G(u, v) = 0$  for  $uv \notin E$ . The diagonal degree matrix of  $G$  is denoted by  $D_G$  with  $D_G(v, v) = \deg(v) = \sum_{u: uv \in E} w_{uv}$  for  $v \in V$ . The random walk matrix of  $G$  is denoted by  $M_G := D_G^{-1} A_G$ . Note that  $M_G$  is a row-stochastic matrix where every row sums to one. Throughout this text, we will use  $M \in \mathbb{R}^{U \times V}$  to denote row-stochastic operators, where  $M \mathbf{1}_V = \mathbf{1}_U$ .

The adjoint of the operator  $B \in \mathbb{R}^{V \times U}$ , with respect to the inner-products defined by  $\Pi_U$  and  $\Pi_V$  on  $U$  and  $V$ , is the unique operator  $B^* \in \mathbb{R}^{U \times V}$  such that

$$\langle \mathbf{f}, B\mathbf{g} \rangle_{\Pi_U} = \langle B^* \mathbf{f}, \mathbf{g} \rangle_{\Pi_V} \quad \text{for all } \mathbf{f} \in \mathbb{R}^U, \mathbf{g} \in \mathbb{R}^V.$$

If  $U = V$  and  $\Pi_U = \Pi_V$ , the operator  $B$  is called self-adjoint when  $B^* = B$ . Note that a real symmetric matrix is self-adjoint with respect to the standard inner-product.

If  $M$  is a row-stochastic self-adjoint operator (with respect to the stationary distribution  $\Pi$ ), then the Markov chain described by  $M$  is called reversible. The random walk operator of an edge-weighted undirected graph is described by the self-adjoint row-stochastic operator  $M_G$  (with respect to the stationary distribution  $\Pi = D_G^{-1} \sum_v \deg(v)$ ) and is a reversible Markov chain.

Let  $W \in \mathbb{R}^{V \times V}$  be a self-adjoint operator with respect to the inner-product defined by  $\Pi$ . It is a fundamental result in linear algebra that  $W$  has only real eigenvalues, and an orthonormal set of eigenvectors with respect to the inner-product defined by  $\Pi$ , i.e.  $\langle f, g \rangle_\Pi = 0$  for eigenvectors  $f \neq g$ . We write  $\lambda_i(W)$  for the  $i$ -th largest eigenvalue of  $W$  so that  $\lambda_1(W) \geq \dots \geq \lambda_{|V|}(W)$ , and write  $\lambda_{\min}(W)$  for the smallest eigenvalue of  $W$ , i.e.  $\lambda_{\min}(W) = \lambda_{|V|}(W)$ . The largest eigenvalue  $\lambda_1(W)$  of a self-adjoint matrix  $W$  with respect to the measure  $\Pi$  obeys the variational formula

$$\lambda_1(W) = \max \left\{ \langle f, Wf \rangle_\Pi : f \in \mathbb{R}^V, \|f\|_\Pi = 1 \right\}. \quad (2.1)$$

It is well-known that the maximizers of the Eq. (2.1) are precisely the unit eigenvectors of  $W$  corresponding to  $\lambda_1(W)$ , i.e.  $Wf = \lambda_1(W) \cdot f$  if and only if  $f$  maximizes the RHS in the Eq. (2.1).

Given an arbitrary operator  $B \in \mathbb{R}^{V \times U}$  we will write  $\sigma_i(B)$  for the  $i$ -th largest singular value of  $B$  so that  $\sigma_1(B) \geq \dots \geq \sigma_{\min\{|U|, |V|\}}(B)$ . It is well known that the singular values of a real operator  $B$  coincide with the eigenvalues of the self-adjoint operator  $BB^*$ .

A self-adjoint operator  $A \in \mathbb{R}^{V \times V}$  with respect to inner-product defined by  $\Pi$  is called positive semi-definite, denoted by  $A \succeq_\Pi 0$ , if it satisfies  $\langle f, Af \rangle_\Pi \geq 0$  for all  $f \in \mathbb{R}^V$ . The condition is equivalent to the condition that  $\lambda_{\min}(A) \geq 0$ . For self-adjoint operators  $A \in \mathbb{R}^{V \times V}$  and  $B \in \mathbb{R}^{V \times V}$  with respect to the same inner-product defined by  $\Pi$ , we will write  $A \preceq_\Pi B$  if

$$\langle f, Af \rangle_\Pi \leq \langle f, Bf \rangle_\Pi \text{ for all } f \in \mathbb{R}^V.$$

This is equivalent to  $A - B$  being positive-semidefinite, i.e.  $A - B \succeq_\Pi 0$ . If  $\Pi$  is just the standard inner-product, we will drop the subscript  $\Pi$ .

We will use the following results about eigenvalues in Section 3 and Section 4; see e.g. [Bha13].

**Fact 2.1.** Let  $A \in \mathbb{R}^{U \times V}$  and  $B \in \mathbb{R}^{V \times U}$ . Then, the non-zero spectrum of  $AB$  coincides with that of  $BA$  with the same multiplicity.

**Fact 2.2.** Let  $A, B \in \mathbb{R}^{V \times V}$  be two self-adjoint matrices with respect to the inner-product defined by  $\Pi$  satisfying  $A \preceq_\Pi B$ . Then,  $\lambda_i(A) \leq \lambda_i(B)$  for all  $1 \leq i \leq |V|$ .

**Theorem 2.3** (Cauchy Interlacing Theorem). Let  $A \in \mathbb{R}^{V \times V}$  be a symmetric matrix and  $B \in \mathbb{R}^{U \times U}$  be a principal submatrix of  $A$ . Let  $n = |V|$  and  $m = |U|$ . For any  $0 \leq j \leq m$ ,

$$\lambda_j(A) \geq \lambda_j(B) \geq \lambda_{n-m+j}(A).$$

**Theorem 2.4** (Weyl Interlacing Theorem). Let  $A, B \in \mathbb{R}^{V \times V}$  be two symmetric matrices. For any  $i, j$ ,

$$\lambda_{i+j-1}(A+B) \leq \lambda_i(A) + \lambda_j(B).$$

## 2.2 Simplicial Complexes

A simplicial complex  $X$  is a collection of subsets that is downward closed, i.e. if  $\beta \in X$  and  $\alpha \subset \beta$  then  $\alpha \in X$ . The elements  $\alpha, \beta$

in  $X$  are called faces/simplices of  $X$ . The dimension of a face  $\alpha$  is defined as  $|\alpha| - 1$ , e.g. an edge is of dimension 1, a vertex/singleton is of dimension 0, the empty set is of dimension  $-1$ . The collection of faces of dimension  $j$  is denoted by  $X(j)$ . The dimension of a simplicial complex is defined as the maximum dimension of its faces. A  $d$ -dimensional simplicial complex is called pure if every maximal face is of dimension  $d$ . All simplicial complexes considered in this paper are pure.

*Weighted Simplicial Complexes.* A simplicial complex  $X$  can be equipped with a weighted function which assigns a positive weight to each face of  $X$ . We follow the formalism of [DDFH18] where the weight function is a probability distribution  $\Pi$  on the faces of the same dimension. Let  $X$  be a  $d$ -dimensional simplicial complex. Given a probability distribution  $\Pi := \Pi_d$  on  $X(d)$ , we can inductively obtain probability distributions  $\Pi_j$  on all  $X(j)$  by considering the marginal distributions, i.e.

$$\Pi_j(\alpha) = \frac{1}{j+2} \sum_{\substack{\beta \in X(j+1), \\ \beta \supset \alpha}} \Pi_{j+1}(\beta). \quad (2.2)$$

Equivalently, we can understand  $\Pi_j$  as the probability distribution of the following random process: Sample a random face  $\beta \in X(d)$  using the probability distribution  $\Pi_d$ , and then sample a uniform random subset of  $\beta$  in  $X(j)$ . The pair  $(X, \Pi)$  will be referred as a weighted simplicial complex. We write  $(X, \Pi)$  simply as  $X$  when  $\Pi$  is the uniform distribution.

*Links and Graphs.* Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex. The link  $X_\alpha$  of a face  $\alpha$  is the simplicial complex defined as

$$X_\alpha := \{\beta \setminus \alpha \mid \beta \in X, \beta \supset \alpha\}.$$

The probability distributions  $\Pi_0, \dots, \Pi_d$  on  $X$  can naturally be used to define the probability distributions  $\Pi_0^\alpha, \dots, \Pi_{d-|\alpha|}^\alpha$  on  $X_\alpha$  using conditional probability. Suppose  $\alpha \in X(j)$ . The probability distribution  $\Pi_l^\alpha$  for  $X_\alpha(l)$  is defined as

$$\Pi_l^\alpha(\tau) = \frac{\Pr_{\beta \sim \Pi_{j+1+l}} [\beta = \alpha \cup \tau \mid \beta \supset \alpha]}{\binom{|\alpha \cup \tau|}{|\alpha|} \cdot \Pi_j(\alpha)} \quad (2.3)$$

for all  $\tau \in X_\alpha(l)$ , where the latter equality is obtained by applying Eq. (2.2) repeatedly.

Given a link  $X_\alpha$ , the graph  $G_\alpha = (X_\alpha(0), X_\alpha(1), \Pi_1^\alpha)$  is defined as the 1-skeleton of  $X_\alpha$ . More explicitly, each singleton  $\{v\}$  in  $X_\alpha$  is a vertex  $v$  in  $G_\alpha$ , each pair  $\{u, v\}$  in  $X_\alpha$  is an edge  $uv$  in  $G_\alpha$ , and the weight of  $uv$  in  $G_\alpha$  is equal to  $\Pi_1^\alpha(\{u, v\})$ . A simple observation is that if  $X$  is a pure  $d$ -dimensional simplicial complex and  $\Pi$  is the uniform distribution on  $X(d)$ , then for any  $\alpha \in X(d-2)$  the weighting  $\Pi_1^\alpha$  on the edges of  $G_\alpha$  is uniform. We will use this observation in Section 4.

## 2.3 Local Spectral Expanders

*Random Walk Matrices.* The definition of local spectral expanders will be based on the random walk matrix of  $G_\alpha$ . Let  $A_\alpha$  be the adjacency matrix of  $G_\alpha$ . Let  $D_\alpha$  be the diagonal degree matrix where  $D_\alpha(x, x) = \sum_y A_\alpha(x, y) = 2\Pi_0^\alpha(x)$  where the last equality is by Eq. (2.2). The random walk matrix  $M_\alpha$  of  $G_\alpha$  is defined as

$M_\alpha := D_\alpha^{-1}A_\alpha$ , with

$$M_\alpha(x, y) = \frac{\Pi_1^\alpha(x, y)}{2\Pi_0^\alpha(x)} \quad \text{for all } \{x, y\} \in X_\alpha(1).$$

The distribution  $\Pi_0^\alpha$  is the stationary distribution of  $M_\alpha$ , as

$$(\Pi_0^\alpha)^\top M_\alpha = (\Pi_0^\alpha)^\top D_\alpha^{-1}A_\alpha = \mathbf{1}^\top A_\alpha = (\Pi_0^\alpha)^\top.$$

The matrix  $M_\alpha$  can be seen to be self-adjoint with respect to the inner-product defined by  $\Pi_0^\alpha$ . So,  $M_\alpha$  have only real eigenvalues, and an orthonormal basis of eigenvectors with respect to the inner-product defined by  $\Pi_0^\alpha$ . The largest eigenvalue of  $M_\alpha$  is 1, as  $M_\alpha \mathbf{1} = \mathbf{1}$  and  $M_\alpha$  is row-stochastic.

Given a vector  $f$ , we will be interested in writing it as  $f = f^1 + f^{\perp 1}$ , so that  $f^1 = c\mathbf{1}$  for some scalar  $c$  and  $\langle f^1, f^{\perp 1} \rangle_{\Pi_0^\alpha} = 0$ .

It follows that  $c = \frac{\langle f, \mathbf{1} \rangle_{\Pi_0^\alpha}}{\langle \mathbf{1}, \mathbf{1} \rangle_{\Pi_0^\alpha}} = \langle f, \mathbf{1} \rangle_{\Pi_0^\alpha} = \mathbb{E}_{x \sim \Pi_0^\alpha} [f(x)]$ . We write

$J_\alpha = (\Pi_0^\alpha)^\top$  as the operator to map  $f$  to  $f^1$ , so that

$$J_\alpha f = (\Pi_0^\alpha)^\top f = \langle f, \Pi_0^\alpha \rangle \cdot \mathbf{1} = \mathbb{E}_{x \sim \Pi_0^\alpha} [f(x)] \cdot \mathbf{1} = f^1 \quad (2.4)$$

*Local Spectral Expanders and Oppenheim's Theorem.* Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex. Define

$$\gamma_j := \gamma_j(X, \Pi) = \max_{\alpha \in X(j)} \lambda_2(M_\alpha) \quad \text{for all } j = -1, \dots, d-2,$$

where  $\lambda_2(M_\alpha)$  is the second largest eigenvalue of the operator  $M_\alpha$ . We say  $X$  is a  $\gamma$ -local-spectral expander if  $\gamma_i \leq \gamma$  for  $-1 \leq i \leq d-2$ .

Oppenheim's Theorem relates the second eigenvalue of the graph of a lower-dimensional link to that of a higher-dimensional link. It works for any weighted simplicial complex with a "balanced" weight function  $w$ , where for any  $\alpha \in X(k)$  and any  $-1 \leq k \leq d-1$  it holds that

$$w(\alpha) = c_k \sum_{\substack{\beta \in X(k+1), \\ \beta > \alpha}} w(\beta)$$

for some constant  $c_k$  that only depends on  $k$ . Note that the weight function in Eq. (2.2) satisfies this condition with  $c_k = 1/(k+2)$ .

**Theorem 2.5** (Oppenheim's Theorem). *Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex where  $\Pi$  satisfies Eq. (2.2). For any  $0 \leq j \leq d-2$ , if  $G_\alpha$  is connected for every  $\alpha \in X(j-1)$ , then*

$$\gamma_{j-1} \leq \frac{\gamma_j}{1 - \gamma_j}.$$

An inductive argument proves the following corollary.

**Corollary 2.6** (Oppenheim's Corollary). *Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex where  $\Pi$  satisfies Eq. (2.2). If  $G_\alpha$  is connected for every  $\alpha \in X(k)$  and every  $k \leq d-2$ , then*

$$\gamma_j \leq \frac{\gamma_{d-2}}{1 - (d-2-j) \cdot \gamma_{d-2}}.$$

## 2.4 Higher Order Random Walks

*Up and Down Operators.* Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex. In the following definitions,  $\alpha \in X(k)$ ,  $\beta \in X(k+1)$ ,  $f \in \mathbb{R}^{X(j)}$ ,  $g \in \mathbb{R}^{X(k+1)}$ , and  $j \in \{-1, 0, \dots, d-1\}$ .

The  $j$ -th up operator  $U_j : \mathbb{R}^{X(j)} \rightarrow \mathbb{R}^{X(j+1)}$  is defined as

$$[U_j f](\beta) = \frac{1}{j+2} \sum_{x \in \beta} f(\beta \setminus x) = \sum_{\substack{\alpha \subset \beta, \\ \alpha \in X(j)}} \frac{f(\alpha)}{j+2}.$$

The  $(j+1)$ -st down operator  $D_{j+1} : \mathbb{R}^{X(j+1)} \rightarrow \mathbb{R}^{X(j)}$  is defined as

$$\begin{aligned} [D_{j+1} g](\alpha) &= \sum_{x \in X_\alpha(0)} \frac{\Pi_{j+1}(\alpha \cup x)}{(j+2)\Pi_j(\alpha)} \cdot g(\alpha \cup x) \\ &= \sum_{\substack{\beta > \alpha, \\ \beta \in X(j+1)}} \frac{\Pi_{j+1}(\beta) \cdot g(\beta)}{(j+2)\Pi_j(\alpha)}. \end{aligned}$$

It can be checked [KO18, DDFH18] that the adjoint of the operator  $U_j$  with respect to the inner-products defined by  $\Pi_{j+1}$  and  $\Pi_j$  is the operator  $D_{j+1}$ , i.e.

$$\langle g, U_j f \rangle_{\Pi_{j+1}} = \langle D_{j+1} g, f \rangle_{\Pi_j}, \quad \forall g \in \mathbb{R}^{X(j+1)}, f \in \mathbb{R}^{X(j)}. \quad (2.5)$$

And it follows that the adjoint of  $D_{j+1}$  with respect to the inner-products defined by  $\Pi_j$  and  $\Pi_{j+1}$  is  $U_j$ , i.e.  $\langle f, D_{j+1} g \rangle_{\Pi_j} = \langle U_j f, g \rangle_{\Pi_{j+1}}$  for all  $g \in \mathbb{R}^{X(j+1)}$ ,  $f \in \mathbb{R}^{X(j)}$ .

*Remark 2.7.* We have stayed consistent with the notations introduced in [DDFH18], and named  $U_j$  and  $D_{j+1}$  up and down operators with their right-action on functions (or vectors) in mind. However, in terms of random walks,  $U_j$  describes a random down-movement from  $X(j+1)$  to  $X(j)$ , whereas  $D_{j+1}$  describes a random up-movement from  $X(j)$  to  $X(j+1)$ , since the action of the probability distribution is from the left.

*Down-Up Walk, Up-Down Walk, and Non-Lazy Up-Down Walk.* We use the up and down operators to define three random walk operators on  $X(j)$ . The  $j$ -th down-up walk  $P_j^\Delta$  and the  $j$ -th up-down walk  $P_j^\nabla$  are defined as

$$P_j^\nabla = U_{j-1} D_j \quad \text{and} \quad P_j^\Delta = D_{j+1} U_j.$$

As  $U_i^* = D_{i+1}$ , it is easy to observe that these operators are positive semi-definite. One useful property of  $P_j^\Delta$  and  $P_j^\nabla$  is that they have the same non-zero spectrum with the same multiplicity by Fact 2.1, and in particular  $\lambda_2(P_j^\Delta) = \lambda_2(P_j^\nabla)$ . Also, we define the  $j$ -th non-lazy up-down walk as

$$P_j^\wedge = \frac{j+2}{j+1} \left( P_j^\Delta - \frac{1}{j+2} \mathbf{1} \right), \quad (2.6)$$

which is the up-down walk conditioned on not looping. It follows from Eq. (2.5), i.e. the adjointness of  $U_j$  and  $D_{j+1}$  that all  $P_j^\nabla$ ,  $P_j^\Delta$ , and  $P_j^\wedge$  are self-adjoint with respect to the inner-product defined by  $\Pi_j$ , e.g. given any  $f_1, f_2 \in \mathbb{R}^{X(j)}$ . This implies that  $\Pi_j$  is the stationary distribution for all these random walks  $P_j^\nabla$ ,  $P_j^\Delta$ , and  $P_j^\wedge$ .

We can understand the higher order random walks as a random walk on a bipartite graph between  $X(j)$  and  $X(j+1)$  as explained in [ALOV19, DK17]. Consider the bipartite graph  $H = (X(j), X(j+1), E)$  in which a face  $\alpha \in X(j)$  and a face  $\beta \in X(j+1)$  are connected if and only if  $\alpha \subset \beta$ . The edge  $\{\alpha, \beta\} \in H$  is assigned the weight  $\frac{1}{j+2} \cdot \Pi_{j+1}(\beta)$ . Using Eq. (2.2), it can be seen that the weighted degree of any  $\alpha \in X(j)$  is  $\Pi_j(\alpha)$ . And the weighted degree of any  $\beta \in X(j+1)$  is exactly  $\Pi_{j+1}(\beta)$ . Thus, the graph  $H$  has the (weighted) random walk matrix

$$M_H = \begin{pmatrix} 0 & U_j \\ D_{j+1} & 0 \end{pmatrix}.$$

One step of the down-up walk  $P_{j+1}^\nabla$  can be thought as a two step random walk in  $M_H$ : starting from some  $\beta \in X(j+1)$ , the random

walk will go down from  $\beta \in X(j+1)$  to  $\alpha \in X(j)$  by dropping an element of  $\beta$ , which is chosen uniformly at random as prescribed by  $U_j$ , and then the random walk will go up from  $\alpha \in X(j)$  to a random face  $\beta' \in X(j+1)$  which contains  $\alpha$  as prescribed by  $D_{j+1}$ . Similarly, one step of the up-down walk  $P_j^\Delta$  can be thought as a two step random walk in  $M_H$  starting from some  $\alpha \in X(j)$ . More precisely,

$$M_H^2 = \begin{pmatrix} U_j D_{j+1} & 0 \\ 0 & D_{j+1} U_j \end{pmatrix} = \begin{pmatrix} P_{j+1}^\nabla & 0 \\ 0 & P_j^\Delta \end{pmatrix}.$$

It is instructive to check that when the distribution  $\Pi$  of the simplicial complex is the uniform distribution, then the down-up walks and the up-down walks are as described as in the introduction.

*Longer Random Walks.* Suppose now  $-1 \leq a < b \leq d$ . We define the up-down walk on  $X(a)$  through  $X(b)$  to be

$$P_{a,b}^\Delta = D_{a+1} \cdots D_b \cdot U_{b-1} \cdots U_a.$$

Similar to the intuition that was presented about the up-down and the down-up walks, we can think of  $P_{a,b}^\Delta$  as simulating two-steps of the random walk starting from some face  $\alpha \in X(a)$  on the weighted bipartite graph  $H = (X(a), X(b), E)$  where  $\{\alpha, \beta\}$  is an edge of this graph with weight proportional to  $\Pi_b(\beta)$  whenever  $\alpha < \beta$ .

## 2.5 Mixing Times of Markov Chains

Recall that two distributions  $\Pi$  and  $\Pi'$  are said to be  $\varepsilon$ -close if

$$\|\Pi - \Pi'\|_{\ell_1} = \sum_{x \in V} |\Pi(x) - \Pi'(x)| \leq \varepsilon. \quad (\varepsilon\text{-close})$$

The mixing time  $T(\varepsilon, P)$  of the random walk operator  $P$  is defined to be the least time step where the distribution of the random walk is  $\varepsilon$ -close to the stationary distribution  $\Pi$  of  $P$  in the  $\ell_1$  distance, i.e.

$$T(\varepsilon, P) = \min\{t \in \mathbb{N}_{\geq 0} : \|P^t(x, \bullet) - \Pi\|_{\ell_1} \leq \varepsilon \text{ for all } x \in V\}.$$

For our applications in sampling in Section 4, we will use the following well known relation between the mixing time of the random walk and the spectral gap of its transition matrix (see e.g. [MT05, Proposition 2.12]).

**Theorem 2.8** (Spectral Mixing Time Bound). *Let  $P \in \mathbb{R}^{V \times V}$  be a random walk matrix with stationary distribution  $\Pi$ . One has,*

$$T(\varepsilon, P) \leq \frac{1}{1 - \sigma_2(P)} \cdot \log \frac{1}{\varepsilon \cdot \min_{x \in V} \Pi(x)},$$

where  $\sigma_2(P)$  is the second largest singular value of  $P$ .

The operator of importance for us will be  $P = P_j^\nabla$ . As this operator is positive semi-definite as explained in Section 2.4, we have  $\sigma_2(P_j^\nabla) = \lambda_2(P_j^\nabla)$ . Also, recall from Section 2.4 that the stationary distribution of  $P_j^\Delta$  is  $\Pi_j$ , we obtain

$$T(\varepsilon, P_j^\nabla) \leq \frac{1}{1 - \lambda_2(P_j^\nabla)} \cdot \log \frac{1}{\varepsilon \cdot \min_{\alpha \in X(j)} \Pi_j(\alpha)}.$$

*Approximate Sampling and Approximate Counting.* There is a well-known equivalence between approximate sampling and approximate counting for self-reducible problems. Let  $\Omega := \{\Omega_s\}_s$  be a collection of sets parametrized by some strings  $s$ , e.g.  $s$  can be describing a graph and  $\Omega_s$  the set of perfect matchings in  $G$ . Suppose a randomized algorithm  $\mathcal{A}$  is given whose output distribution

is described by  $\mu_{\mathcal{A}(s)}$ . Then  $\mathcal{A}$  is called a fully polynomial time randomized approximate uniform sampler (FPRAUS) for  $\Omega_s$ , if for every input string  $s$  we have

$$\|\mu_{\mathcal{A}(s)} - \Pi_{\Omega_s}\|_{\ell_1} \leq \delta,$$

where  $\Pi_{\Omega_s}$  describes the uniform distribution over  $\Omega_s$  and the algorithm  $\mathcal{A}$  runs in time  $\text{poly}(\langle s \rangle, \log(1/\delta))$ , where  $\langle s \rangle$  denotes the size of the input.

Similarly, an algorithm  $\mathcal{A}'$  is called a fully polynomial time randomized approximation scheme (FPRAS) for  $\Omega$ , if we for every input  $s$  we have

$$\Pr[(1 - \delta) \cdot |\Omega_s| \leq \mathcal{A}'(s) \leq (1 + \delta) \cdot |\Omega_s|] \geq 1 - \varepsilon,$$

and the algorithm  $\mathcal{A}'$  runs in time  $\text{poly}(\langle s \rangle, 1/\varepsilon, \log(1/\delta))$ .

A well-known result proven in [JVV86] asserts that approximate counting and approximate sampling are equivalent for self-reducible problems.

**Theorem 2.9** (Informal). *For self-reducible sets  $\Omega$  in NP, the existence of an FPRAS for  $\Omega$  is equivalent to the existence of an FPRAUS for  $\Omega$ .*

In Section 4, we will give approximate samplers for independent sets a graph, and it follows from this equivalence we can also approximately count the number of independent sets in the graph.

## 3 EIGENVALUE BOUNDS FOR HIGHER ORDER RANDOM WALKS

Our main result is a quantitative generalization of the basic fact that a pure  $d$ -dimensional simplicial complex  $X$  is gallery connected (i.e.  $\lambda_2(P_d^\nabla) < 1$ ) if and only if the graph  $G_\alpha$  is connected for every  $\alpha \in X$  up to dimension  $d-2$  (i.e.  $\gamma_j < 1$  for  $-1 \leq j \leq d-2$ ). The statement is essentially the same as in Theorem 1.5 but for more general weighted simplicial complexes.

**Theorem 3.1.** *Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex. For any  $0 \leq k \leq d$ ,*

$$\lambda_2(P_k^\nabla) = \lambda_2(P_{k-1}^\Delta) \leq 1 - \frac{1}{k+1} \prod_{j=-1}^{k-2} (1 - \gamma_j).$$

Using an inductive argument as in [ALOV19, Theorem 3.3], we can prove a more general statement about the entire range of eigenvalues.

**Theorem 3.2.** *Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex. Then, for any  $0 \leq k \leq d-1$  and for any  $-2 \leq r \leq k$ , the matrix  $P_k^\Delta$  has at most  $|X(r)|$  eigenvalues with value strictly greater than*

$$1 - \frac{r+2}{k+2} \prod_{j=r}^{k-1} (1 - \gamma_j),$$

where we adopt the conventions that  $X(-2) = \emptyset$  and  $\prod_{j=r}^{-1} (1 - \gamma_j) = 1$ .

Note that Theorem 3.1 is a special case of Theorem 3.2 where  $r = -1$  (recall that  $X(-1) = \{\emptyset\}$  and so  $|X(-1)| = 1$ ). Further, Theorem 3.1 can only prove that  $\lambda_2(P_d^\nabla) \leq 1 - \frac{1}{d+1}$ . We observe that this bound is almost tight. We prove the following in the full version of this paper.



**Proposition 3.3.** *Let  $X$  be a  $d$ -dimensional simplicial complex. Let  $n = |X(0)|$ . Suppose  $2(d+1) \leq n$ . Then  $\lambda_2(P_d^\nabla) = \lambda_2(P_{d-1}^\Delta) \geq 1 - \frac{2}{d+1}$ .*

Before we prove [Theorem 3.1](#) and [Theorem 3.2](#), we present two corollaries of [Theorem 3.1](#).

Combining with Oppenheim's [Corollary 2.6](#), [Theorem 3.1](#) provides a bound on the second eigenvalue of the  $d$ -th down-up walk based only on the maximum second eigenvalue of the graphs in dimension  $d - 2$ . This will be useful in [Section 4](#).

**Corollary 3.4.** *Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex. For any  $0 \leq k \leq d$ , suppose  $\gamma_k \leq \frac{1}{k+1}$  and  $\gamma_j < 1$  for  $-1 \leq j \leq k - 2$ , then*

$$\lambda_2(P_k^\nabla) = \lambda_2(P_{k-1}^\Delta) \leq 1 - \frac{1}{(k+1)^2}.$$

PROOF. Since  $\gamma_{k-2} \leq \frac{1}{k+1}$  and  $\gamma_j < 1$  for  $-1 \leq j \leq k - 2$ , it follows from Oppenheim's [Corollary 2.6](#) that for any  $-1 \leq j \leq k - 3$ ,

$$\gamma_j \leq \frac{\gamma_{k-2}}{1 - (k-2-j) \cdot \gamma_{k-2}} \leq \frac{\frac{1}{k+1}}{1 - \frac{k-2-j}{k+1}} = \frac{1}{j+3}.$$

Therefore, by [Theorem 3.1](#),

$$\lambda_2(P_k^\nabla) \leq 1 - \frac{1}{k+1} \prod_{j=-1}^{k-2} \frac{j+2}{j+3} = 1 - \frac{1}{(k+1)^2}. \quad \square$$

**Corollary 3.5.** *Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex. Let  $0 \leq a < b \leq d - 1$ . If  $X$  is a  $\gamma$ -local-spectral expander, then*

$$\lambda_2(P_{a,b}^\Delta) \leq (1 + \gamma)^{b-a} \cdot \frac{a+1}{b+1}.$$

We will first prove [Theorem 3.1](#) in [Section 3.1](#), then [Theorem 3.2](#) in [Section 3.3](#), and finally [Corollary 3.5](#) in [Section 3.4](#).

### 3.1 Proof of [Theorem 3.1](#)

The key lemma in proving [Theorem 3.1](#) is the following result that quantifies a spectral bound on the difference of the  $k$ -th non-lazy up-down walk and the  $k$ -th down-up walk in terms of the second eigenvalue of the links at dimension  $k - 1$ .

**Lemma 3.6.** *Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex. For any  $0 \leq k \leq d - 1$ ,*

$$P_k^\Delta - P_k^\nabla \leq \Pi_k \gamma_{k-1} \cdot (I - P_k^\nabla).$$

The proof of [Lemma 3.6](#), will closely follow the proof of [[DDFH18](#), [Theorem 5.5](#)], where they prove the weaker inequality

$$P_k^\Delta - P_k^\nabla \leq \Pi_k \gamma_{k-1} \cdot I. \quad (3.1)$$

We remark that a similar statement was also used in [[KO18](#)] for proving [Theorem 1.2](#).

We will first show how [Lemma 3.6](#) implies [Theorem 3.1](#) by an inductive argument.

PROOF OF [THEOREM 3.1](#) FROM [LEMMA 3.6](#). We prove [Theorem 3.1](#) by induction on  $k$ . The base case is when  $k = 0$ , where  $P_0^\nabla = \mathbf{1}\mathbf{1}_0^\top$  is a rank one matrix and so  $\lambda_2(P_0^\nabla) \leq 0$ , and hence [Theorem 3.1](#) trivially holds.

For the induction step, suppose we have

$$\lambda_2(P_{j+1}^\nabla) = \lambda_2(P_j^\Delta) \leq 1 - \frac{1}{j+2} \prod_{i=-1}^{j-1} (1 - \gamma_i). \quad (3.2)$$

Since  $P_{j+1}^\nabla = U_j D_{j+1}$  and  $P_j^\Delta = D_{j+1} U_j$  have the same non-zero eigenvalues with the same multiplicity by [Fact 2.1](#), we only need to prove the statement for  $P_{j+1}^\Delta$ . By [Lemma 3.6](#),

$$P_{j+1}^\Delta \leq \Pi_{j+1} \gamma_j \cdot I + (1 - \gamma_j) P_{j+1}^\nabla$$

It follows from [Fact 2.2](#) that

$$\lambda_2(P_{j+1}^\Delta) \leq \gamma_j + (1 - \gamma_j) \cdot \lambda_2(P_{j+1}^\nabla) \leq 1 - \frac{1}{j+2} \prod_{i=-1}^j (1 - \gamma_i),$$

where the last equality is by plugging in the induction hypothesis, [Eq. \(3.2\)](#). The theorem now follows from the definition of the non-lazy up-down walk, [Eq. \(2.6\)](#), i.e.

$$P_{j+1}^\Delta = \frac{j+3}{j+2} \left( P_{j+1}^\Delta - \frac{1}{j+3} I \right) \iff P_{j+1}^\Delta = \frac{j+2}{j+3} \cdot P_{j+1}^\Delta + \frac{1}{j+3} I.$$

Therefore,

$$\lambda_2(P_{j+1}^\Delta) = \frac{j+2}{j+3} \cdot \lambda_2(P_{j+1}^\Delta) + \frac{1}{j+3} \leq 1 - \frac{1}{j+3} \prod_{i=-1}^j (1 - \gamma_i),$$

and this proves the induction step.  $\square$

### 3.2 Proof of [Lemma 3.6](#)

The proof of [Lemma 3.6](#) will rest on few useful identities established in [[KO18](#), [DDFH18](#)], which can be obtained through the ‘‘Garland Method’’, which decomposes the higher order random walk matrices into the random walk matrices of the links. We provide proofs for these identities in the full version of the paper.

In the following, given  $f \in \mathbb{R}^{X(k)}$  and  $\alpha \in X(k-1)$ , we use  $f_\alpha$  to denote the restriction of  $f$  to the entries in  $\{\alpha \cup \{x\} \mid x \in X_\alpha(0)\}$ .

**Lemma 3.7.** *Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex. For all  $f \in \mathbb{R}^{X(j)}$  the following hold,*

- (1)  $\langle f, If \rangle_{\Pi_j} = \mathbb{E}_{\alpha \sim \Pi_{j-1}} \|f_\alpha\|_{\Pi_0^\alpha}^2 = \mathbb{E}_{\alpha \sim \Pi_{j-1}} \langle f_\alpha, f_\alpha \rangle_{\Pi_0^\alpha}$ ,
- (2)  $\langle f, P_j^\nabla f \rangle_{\Pi_j} = \mathbb{E}_{\alpha \sim \Pi_{j-1}} \|J_\alpha f_\alpha\|_{\Pi_0^\alpha}^2 = \mathbb{E}_{\alpha \sim \Pi_{j-1}} \langle f_\alpha, J_\alpha f_\alpha \rangle_{\Pi_0^\alpha}$ ,
- (3)  $\langle f, P_j^\Delta f \rangle_{\Pi_j} = \mathbb{E}_{\alpha \sim \Pi_{j-1}} \langle f_\alpha, M_\alpha f_\alpha \rangle_{\Pi_0^\alpha}$ .

PROOF OF [LEMMA 3.6](#). Let  $f \in \mathbb{R}^{X(j)}$  be arbitrary. By Items (2) and (3) in [Lemma 3.7](#), we write

$$\langle f, (P_j^\Delta - P_j^\nabla) f \rangle_{\Pi_j} = \mathbb{E}_{\alpha \sim \Pi_{j-1}} \left[ \langle f_\alpha, (M_\alpha - J_\alpha) f_\alpha \rangle_{\Pi_0^\alpha} \right].$$

Notice that since  $M_\alpha$  is a row-stochastic matrix (with top eigenvector  $\mathbf{1}$ ) and the matrix  $J_\alpha$  is the projector to its top eigenspace. Since both  $M_\alpha$  and  $J_\alpha$  are self-adjoint with respect to the inner-product defined by  $\Pi_0^\alpha$  (see [Section 2.3](#)), it follows that

$$M_\alpha - J_\alpha \leq \Pi_0^\alpha \lambda_2(M_\alpha) \cdot I.$$

Moreover, since the matrix  $M_\alpha - J_\alpha$  is only supported on the subspace perpendicular to  $\mathbf{1}$ , writing  $f_\alpha^{\perp 1}$  for the component of  $f_\alpha$  that is perpendicular to  $\mathbf{1}$ , we have

$$\langle f_\alpha, (M_\alpha - J_\alpha) f_\alpha \rangle_{\Pi_0^\alpha} = \langle f_\alpha^{\perp 1}, (M_\alpha - J_\alpha) f_\alpha^{\perp 1} \rangle_{\Pi_0^\alpha}.$$

As  $J_\alpha$  is the projector to the constant functions (2.4) we have,  $f_\alpha^{\perp 1} = (1 - J_\alpha)f_\alpha$  and thus

$$\begin{aligned} \langle f_\alpha, (M_\alpha - J_\alpha)f_\alpha \rangle &\leq \lambda_1(M_\alpha - J_\alpha) \cdot \|f_\alpha^{\perp 1}\|_{\Pi_0^\alpha}^2, \\ &\leq \lambda_2(M_\alpha) \cdot \|(1 - J_\alpha)f_\alpha\|_{\Pi_0^\alpha}^2, \end{aligned} \quad (3.3)$$

where the first inequality is by Eq. (2.1). Therefore,

$$\langle f, (P_j^\wedge - P_j^\vee)f \rangle_{\Pi_j} = \mathbb{E}_{\alpha \sim \Pi_{j-1}} \left[ \langle f_\alpha, (M_\alpha - J_\alpha)f_\alpha \rangle_{\Pi_0^\alpha} \right], \quad (3.4)$$

$$\leq \mathbb{E}_{\alpha \sim \Pi_{j-1}} \left[ \lambda_2(M_\alpha) \cdot \|(1 - J_\alpha)f_\alpha\|_{\Pi_0^\alpha}^2 \right], \quad (3.5)$$

$$\begin{aligned} &\leq \gamma_{j-1} \cdot \mathbb{E}_{\alpha \sim \Pi_{j-1}} \left[ \|(1 - J_\alpha)f_\alpha\|_{\Pi_0^\alpha}^2 \right], \\ &= \gamma_{j-1} \cdot \mathbb{E}_{\alpha \sim \Pi_{j-1}} \left[ \langle f_\alpha, (1 - J_\alpha)f_\alpha \rangle_{\Pi_0^\alpha} \right], \end{aligned} \quad (3.6)$$

$$= \gamma_{j-1} \cdot \langle f, (1 - P_j^\vee)f \rangle_{\Pi_j} \quad (3.7)$$

where we have obtained Eq. (3.4) by Items (2) and (3) in Lemma 3.7, Eq. (3.5) by Eq. (3.3), Eq. (3.6) by  $\langle J_\alpha f_\alpha, J_\alpha f_\alpha \rangle_{\Pi_0^\alpha} = \langle f_\alpha, J_\alpha f_\alpha \rangle_{\Pi_0^\alpha}$ , and Eq. (3.7) by Items (1) and (2) in Lemma 3.7. This proves the lemma.  $\square$

### 3.3 Proof of Theorem 3.2

We will prove Theorem 3.2 about the entire spectrum of the higher order random walks.

**Theorem 3.2.** *Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex. Then, for any  $0 \leq k \leq d-1$  and for any  $-2 \leq r \leq k$ , the matrix  $P_k^\Delta$  has at most  $|X(r)|$  eigenvalues with value strictly greater than*

$$1 - \frac{r+2}{k+2} \prod_{j=r}^{k-1} (1 - \gamma_j),$$

where we adopt the conventions that  $X(-2) = \emptyset$  and  $\prod_{j=r}^{r-1} (1 - \gamma_j) = 1$ .

**PROOF.** We prove by induction on  $k$ . The base case is when  $k = 0$ , where  $P_0^\Delta = \frac{1}{2}M_\emptyset + \frac{1}{2}I$ . For  $r = -2$ , the claim states that we have at most  $|X(-2)| = 0$  eigenvalues that are strictly greater than 1, which is true since  $P_0^\Delta$  is a stochastic matrix. For  $r = -1$ , the claim tells us that there are at most  $|X(-1)| = 1$  eigenvalues which are strictly larger than  $\frac{1}{2} + \frac{\gamma_{-1}}{2}$ , which is true by the definition of  $\gamma_{-1} = \lambda_2(M_\emptyset)$  and that since  $P_0^\Delta$  is of rank  $|X(0)|$ . For  $r = 0$ , the claim tells us that there are at most  $|X(0)|$  eigenvalues that are greater than 0, which is true since this is the rank of  $|X(0)|$ .

For the induction step, suppose that there exists some  $j \geq 1$  such that the claim of the theorem is true for all  $-1 \leq r \leq j$ . By Fact 2.1,  $P_{j+1}^\vee$  and  $P_j^\Delta$  have the same non-zero eigenvalues. By Lemma 3.6,  $P_{j+1}^\Delta \leq \Pi_{j+1} \gamma_j I + (1 - \gamma_j)P_j^\vee$ , and thus for  $-1 \leq r \leq j$  the matrix  $P_{j+1}^\Delta$  has at most  $|X(r)|$  eigenvalues with value greater than

$$\gamma_j + (1 - \gamma_j) \cdot \left( 1 - \frac{r+2}{j+2} \prod_{i=r}^{j-1} (1 - \gamma_i) \right) = 1 - \frac{r+2}{j+2} \prod_{i=r}^j (1 - \gamma_i).$$

Using the definition of the non-lazy up-down walk, Eq. (2.6), we have that for  $-1 \leq r \leq j$ ,  $P_{j+1}^\Delta$  has at most  $|X(r)|$  eigenvalues with

value greater than

$$\frac{j+2}{j+3} \left( 1 - \frac{r+2}{j+2} \prod_{i=r}^j (1 - \gamma_i) \right) + \frac{1}{j+3} = 1 - \frac{r+2}{j+3} \prod_{i=r}^j (1 - \gamma_i).$$

For  $r = j+1$ , the claim is trivial since the  $P_{j+1}^\Delta$  is of rank  $|X(j+1)|$ .  $\square$

### 3.4 Proof of Corollary 3.5

**Corollary 3.5.** *Let  $(X, \Pi)$  be a pure  $d$ -dimensional weighted simplicial complex. Let  $0 \leq a < b \leq d-1$ . If  $X$  is a  $\gamma$ -local-spectral expander, then*

$$\lambda_2(P_{a,b}^\Delta) \leq (1 + \gamma)^{b-a} \cdot \frac{a+1}{b+1}.$$

We will use two basic facts in the proof.

**Fact 3.8.** *Let  $M_1 \in \mathbb{R}^{V \times U}$  and  $M_2 \in \mathbb{R}^{U \times W}$  be two row-stochastic matrices. Then, we have  $\sigma_2(M_1 \cdot M_2) \leq \sigma_2(M_1) \cdot \sigma_2(M_2)$ .*

**Fact 3.9** (Bernoulli's Inequality). *Let  $x \geq -1$  and  $r \geq 1$  be real numbers. Then,  $(1+x)^r \geq 1+r \cdot x$ .*

**PROOF.** Recall that  $P_{a,b}^\Delta = D_{a+1} \cdots D_b \cdot U_{b-1} \cdots U_a$ . As  $U_i^* = D_{i+1}$ , it can be observed that  $P_{a,b}^\Delta$  is positive semi-definite and therefore,  $\sigma_2(P_{a,b}^\Delta) = \lambda_2(P_{a,b}^\Delta)$ . By Fact 3.8,

$$\lambda_2(P_{a,b}^\Delta) = \sigma_2(P_{a,b}^\Delta) \leq \sigma_2(D_{a+1}) \cdots \sigma_2(D_b) \cdot \sigma_2(U_{b-1}) \cdots \sigma_2(U_a).$$

Notice that as  $P_{a,j}^\Delta = D_{j+1}U_j$  and  $D_{j+1}^* = U_j$ , we have  $\lambda_2(P_{a,j}^\Delta) = \sigma_2(U_j) \cdot \sigma_2(D_{j+1})$ . Thus, by rearranging we obtain,

$$\begin{aligned} \lambda_2(P_{a,b}^\Delta) &\leq \prod_{j=0}^{b-a-1} \lambda_2(P_{a+j}^\Delta), \\ &\leq \prod_{j=0}^{b-a-1} \left( 1 - \frac{(1-\gamma)^{a+j+1}}{a+j+2} \right), \quad (\text{by Theorem 3.1}) \\ &\leq \prod_{j=0}^{b-a-1} \left( 1 - \frac{1 - (a+j+1) \cdot \gamma}{a+j+2} \right), \quad (\text{by Fact 3.9}) \\ &= \prod_{j=0}^{b-a-1} \left( (1+\gamma) \frac{a+j+1}{a+j+2} \right). \end{aligned}$$

By cancellations in the telescoping product, we have

$$\lambda_2(P_{b,a}^\Delta) \leq (1 + \gamma)^{b-a} \cdot \frac{a+1}{b+1}. \quad \square$$

## 4 ANALYZING MIXING TIMES OF MARKOV CHAINS

In this section, we will use Corollary 3.4 to analyze Markov chains for sampling independent sets of a graph of fixed size and sampling common independent sets of two partition matroids.

## 4.1 Independent Sets

Let  $G = (V, E)$  be a graph. A subset of vertices  $S \subset V$  is called an independent set if  $uv \notin E$  for every pair  $u, v \in S$ . We are interested in the problem of sampling a uniformly random independent set of size  $k$ . We will analyze a natural Markov chain for the problem by analyzing the down-up walk of a corresponding simplicial complex.

Define the  $(k-1)$ -dimensional simplicial complex  $I_{G,k}$  of  $G = (V, E)$  as

$$I_{G,k} = \{S \subset V : |S| \leq k \text{ and } S \text{ is independent}\},$$

the complex consisting of all independent sets in  $G$  of cardinality at most  $k$ . We endow  $I_{G,k}$  with the uniform distribution  $\Pi_{k-1}$  on  $I_{G,k}(k-1)$ , i.e. the set of independent sets of size  $k$ . We simply write  $I_{G,k}$  for the weighted simplicial complex  $(I_{G,k}, \Pi_{k-1})$ .

The  $(k-1)$ -th down-up walk  $P_{k-1}^\nabla$  on  $I_{G,k}$  corresponds to a natural Markov chain to sample independent sets of size  $k$ . It is known that this Markov chain is fast mixing when  $k \leq \frac{|V|}{2\Delta+1}$  using coupling techniques [BD97, MU05]. The main result in this subsection is the following improved bound using higher order random walks on simplicial complexes.

**Theorem 1.7.** *Let  $G = (V, E)$  be a graph with maximum degree  $\Delta$ . Let  $P_{k-1}^\nabla$  be the  $(k-1)$ -th down-up walk on the simplicial complex  $I_{G,k}$ . Let  $A_G$  be the adjacency matrix of  $G$ .*

$$\text{If } k \leq \frac{|V|}{\Delta + |\lambda_{\min}(A_G)|}, \text{ then } \lambda_2(P_{k-1}^\nabla) \leq 1 - \frac{1}{k^2}.$$

It is well-known that  $|\lambda_{\min}(A_G)| \leq \Delta$  for a graph with maximum degree  $\Delta$ , and so Theorem 1.7 recovers the previous result that the Markov chain is fast mixing if  $k \leq \frac{|V|}{2\Delta}$ . There are various graph classes with  $|\lambda_{\min}(A_G)|$  smaller than  $\Delta$ , and Theorem 1.7 allows us to sample larger independent sets. For example, it is known that  $|\lambda_{\min}(A_G)| \leq O(\sqrt{\Delta})$  for planar graphs and more generally for graphs with bounded arboricity [Hay06], and also for random graphs and more generally for two-sided expander graphs [HLW06].

Using the simple bound  $\min_{S \in I_{G,k}(k-1)} \Pi_{k-1}(S) \geq n^{-k}$  as  $\Pi_{k-1}$  is the uniform distribution, the following mixing time result follows from Theorem 2.8.

**Corollary 4.1.** *Let  $G = (V, E)$  be a graph with maximum degree  $\Delta$  and let  $A_G$  be the adjacency matrix of  $G$ . For any  $k \leq n/(\Delta + |\lambda_{\min}(A_G)|)$ , the down-up walk  $P_{k-1}^\nabla$  on the simplicial complex  $I_{G,k}$  samples a random independent set of  $G$  of size  $k$  whose distribution is  $\epsilon$ -close to the uniform distribution on all independent sets of size  $k$  in*

$$T(\epsilon, P_{k-1}^\nabla) \leq k^2 \cdot \left( \log\left(\frac{1}{\epsilon}\right) + k \cdot \log n \right)$$

many time steps.

This implies a polynomial time algorithm to approximately sample a uniform random independent set and also a FPRAS for approximately counting the number of independent set of size  $k$  for  $k \leq \frac{n}{\Delta + |\lambda_{\min}(A_G)|}$ .

**4.1.1 Proof of Theorem 1.7.** The plan is to use Corollary 3.4 to prove Theorem 1.7. To apply Corollary 3.4, we need to prove that:

- (1)  $I_{G,k}$  is a pure simplicial complex. It is a simple exercise that this complex is pure when  $k \leq \frac{n}{\Delta+1}$ .

- (2) For each  $S \in I_{G,k}$  with  $|S| \leq k-2$ , the random walk matrix  $M_S$  of the graph  $G_S$  of the link  $(I_{G,k})_S$  satisfies  $\lambda_2(M_S) < 1$ . This is proved in Lemma 4.2.

- (3) For each  $S \in I_{G,k}$  with  $|S| = k-2$ , the random walk matrix  $M_S$  of the graph  $G_S$  satisfies  $\lambda_2(M_S) \leq 1/k$ . This is proved in Lemma 4.3.

Assuming the three items are proven, Theorem 1.7 follows immediately from Corollary 3.4. We will prove the second item in Section 4.1.2 and the third item in Section 4.1.3.

**4.1.2 Proof of Lemma 4.2.** Let  $H_S = (V_S, E_S)$  be the underlying support graph of  $G_S$  of the link  $(I_{G,k})_S$ , i.e.  $G_S$  without edge weights. Let  $M_S$  be the random walk matrix of  $G_S$  as defined in Section 2.3. Note that  $\lambda_2(M_S) < 1$  if and only if  $H_S$  is connected.

We introduce some notation to describe  $H_S$ . We write  $N_G[S]$  as the union of  $S$  and the set of vertices which are connected to a vertex in  $S$  in  $G$ , i.e.

$$N_G[S] = S \cup \{v : \text{there exists some } uv \in E(G) \text{ such that } u \in S\}.$$

For a subset of vertices  $S \subset V(G)$ , we write  $\bar{S} = V(G) \setminus S$  for the complement of  $S$  in  $G$ , and  $G[S]$  for the induced subgraph of  $G$  on  $S$ . For a graph  $H$ , we write  $\bar{H}$  for the complement graph of  $H$ .

Recall that a vertex  $v$  is in  $V_S$  if and only if  $S \cup \{v\}$  is an independent set in  $G$  of size  $|S|+1$ , and so  $V_S$  is exactly  $V - N_G[S] = \bar{N}_G[S]$ . Two vertices  $u, v \in V_S$  have an edge in  $H_S$  if and only if  $S \cup \{u, v\}$  is an independent set in  $G$  of size  $|S|+2$ , and so  $uv \in E_S$  if and only if  $uv \notin E(G)$ . Therefore, we see that  $H_S = \overline{G[V_S]} = \overline{G[N[S]]}$ . With the description of  $H_S$ , we are ready to prove the second item in Section 4.1.1.

**Lemma 4.2.** *Let  $G = (V, E)$  be a graph with maximum degree  $\Delta$ . Suppose  $k \leq \frac{|V|}{\Delta+1}$ . For any  $S \in I_{G,k}$  with  $|S| \leq k-2$ , the random walk matrix  $M_S$  of the graph  $G_S$  of the link  $(I_{G,k})_S$  satisfies  $\lambda_2(M_S) < 1$ .*

**PROOF.** Note that  $\lambda_2(M_S) < 1$  if and only if the underlying support graph  $H_S$  of  $G_S$  is connected, so we focus on proving the latter. To prove that  $H_S$  is connected, we prove the stronger claim that every two vertices  $u, v \in H_S$  has a path of length at most two. If  $uv$  is an edge in  $H_S$ , then there is a path of length one. Suppose  $uv$  is not an edge in  $H_S$ . Then  $uv$  is an edge in  $G$ . Since  $G$  is of maximum degree  $\Delta$ , it implies that  $|N_G[\{u, v\}]| \leq (\deg_G(u) + 1) + (\deg_G(v) + 1) - 2 \leq 2\Delta$ , and also

$$|V_S| = |V| - |N_G[S]| \geq |V| - |S| \cdot (\Delta + 1) \geq 2\Delta + 2,$$

where we use the assumptions that  $|S| \leq k-2 \leq \frac{|V|}{\Delta+1} - 2$  in the last inequality. So, there must be some vertex  $w$  such that  $w \in V_S \setminus N_G[\{u, v\}]$ . This implies that  $wu \notin E(G)$  and  $wv \notin E(G)$ , and thus  $wu \in E(H_S)$  and  $wv \in E(H_S)$  and so there is a path of length two connecting  $u$  and  $v$  in  $H_S$ .  $\square$

**4.1.3 Proof of Lemma 4.3.** We observe that  $G_S$  is an unweighted graph for  $S$  with  $|S| = k-2$  when the distribution on  $I_{G,k}(k-1)$  is the uniform distribution. Therefore,  $G_S$  is simply a scaled version of  $H_S$ , and the random walk matrix  $M_S$  of  $G_S$  is the same as the random walk matrix of  $H_S$ . To bound the second eigenvalue, we will use some simple interlacing arguments. We need the stronger assumption that  $k \leq \frac{|V(G)|}{\Delta + |\lambda_{\min}(A_G)|}$  in the proof of the following lemma. (Note that for any unweighted graph  $G$ , we have  $|\lambda_{\min}(A_G)| \geq 1$ .)

**Lemma 4.3.** *Let  $G = (V, E)$  be a graph with maximum degree  $\Delta$ . Suppose  $k \leq |V|/(\Delta + |\lambda_{\min}(A_G)|)$ . For any  $S \in I_{G,k}$  with  $|S| = k - 2$ , the random walk matrix  $M_S$  of the graph  $G_S$  of the link  $(I_{G,k})_S$  satisfies  $\lambda_2(M_S) \leq \frac{1}{k}$ .*

**PROOF OF LEMMA 4.3.** Recall that for  $S$  with  $|S| = k - 2$ , the random walk matrix  $M_S$  of  $G_S$  is the same as the random walk matrix of  $H_S$ , and so we will focus on the latter. Let  $D_H$  be diagonal degree matrix of  $H_S$ . As argued above, the random walk matrix  $M_S$  of  $G_S$  is equal to  $M_S = D_H^{-1}A_H$ . We can write the adjacency matrix  $A_H$  of  $H_S$  as  $A_H = \mathbf{1}\mathbf{1}^\top - I - A_{G[\overline{N[S]}]}$ , where  $A_{G[\overline{N[S]}]}$  is the adjacency matrix of  $G[\overline{N[S]}]$ . By Weyl's Interlacing [Theorem 2.4](#),

$$\lambda_2(M_S) \leq \lambda_2(D_H^{-1/2}\mathbf{1}\mathbf{1}^\top D_H^{-1/2}) + \lambda_1(D_H^{-1/2}(-A_{G[\overline{N[S]}]})D_H^{-1/2}),$$

$$= \lambda_1(D_H^{-1/2}(-A_{G[\overline{N[S]}]})D_H^{-1/2}), \quad (4.1)$$

$$\leq \|D_H^{-1}\| \cdot \lambda_1(-A_{G[\overline{N[S]}]}) - 1, \quad (4.2)$$

$$\leq \|D_H^{-1}\| \cdot (|\lambda_{\min}(A_{G[\overline{N[S]}]})| - 1),$$

$$\leq \|D_H^{-1}\| \cdot (|\lambda_{\min}(A_G)| - 1),$$

where for [Eq. \(4.1\)](#) we have used that  $D_H^{-1/2}\mathbf{1}\mathbf{1}^\top D_H^{-1/2}$  is of rank 1, for [Eq. \(4.2\)](#) we have used [Eq. \(2.1\)](#) in the following way: Let  $W = -A_L - I$  and  $\mathbf{g}$  be an unit top-eigenvector of  $D_H^{-1/2}WD_H^{-1/2}$ , i.e.  $\|\mathbf{g}\| = 1$  and  $\langle \mathbf{g}, D_H^{-1/2}WD_H^{-1/2}\mathbf{g} \rangle = \lambda_1(D_H^{-1/2}WD_H^{-1/2})$ . Then,

$$\lambda_1(W) \geq \left\langle \frac{D_H^{-1/2}\mathbf{g}}{\|D_H^{-1/2}\mathbf{g}\|}, W \frac{D_H^{-1/2}\mathbf{g}}{\|D_H^{-1/2}\mathbf{g}\|} \right\rangle = \frac{\lambda_1(D_H^{-1/2}WD_H^{-1/2})}{\|D_H^{-1/2}\|^2}$$

$$\geq \frac{\lambda_1(D_H^{-1/2}WD_H^{-1/2})}{\|D_H^{-1}\|},$$

and for the last inequality we have used the Cauchy Interlacing [Theorem 2.3](#).

It remains to bound  $\|D_H^{-1}\| = (\min_v \deg_{H_S}(v))^{-1}$ . As  $H_S = G[\overline{N[S]}] = G[V - N[S]]$ ,

$$\deg_{H_S}(v) = |V| - |N[S]| - (\deg_{G[\overline{N[S]}]}(v) + 1) \geq |V| - (\Delta + 1)(|S| + 1),$$

where the last inequality uses that  $|N[S]| \leq |S| \cdot (\Delta + 1)$  and  $\deg_{G[\overline{N[S]}]}(v) \leq \deg_G(v) \leq \Delta$ . Therefore, using our bound  $\lambda_2(M_S) \leq \|D_H^{-1}\| \cdot (|\lambda_{\min}(A_G)| - 1)$ , we obtain

$$\lambda_2(M_S) \leq \frac{|\lambda_{\min}(A_G)| - 1}{|V| - (\Delta + 1) \cdot (|S| + 1)} = \frac{|\lambda_{\min}(A_G)| - 1}{|V| - (\Delta + 1) \cdot (k - 1)},$$

where we use  $|S| = k - 2$ . Finally, plugging in the assumption  $k \leq \frac{|V|}{\Delta + |\lambda_{\min}(A_G)|}$  implies the lemma.  $\square$

## 4.2 Matroid Intersection

A matroid  $M = (E, \mathcal{I})$  on the ground set  $E$  with the set of independent sets  $\mathcal{I} \subset 2^E$  is a combinatorial object satisfying the following properties:

- (containment property) if  $S \in \mathcal{I}$  and  $T \subset S$ , then  $T \in \mathcal{I}$ ,
- (extension property) if  $S, T \in \mathcal{I}$  such that  $|S| > |T|$  then there is some  $x \in S \setminus T$  such that  $\{x\} \cup T \in \mathcal{I}$ .

A partition matroid is the special case where the ground set  $E$  is partitioned into disjoint blocks  $B_1, \dots, B_l \subseteq E$  with parameters  $0 \leq d_i \leq |B_i|$  for  $1 \leq i \leq l$ , and a subset  $S$  is in  $\mathcal{I}$  if and only if  $|S \cap B_i| \leq d_i$  for  $1 \leq i \leq l$ .

The intersection of two matroids  $M_1 = (E, \mathcal{I}_1)$  and  $M_2 = (E, \mathcal{I}_2)$  over the same ground set  $E$  can be used to formulate various interesting combinatorial optimization problems [[Sch03](#)]. We are interested in the problem of sampling a uniform random common independent set of size  $k$ , i.e. a random subset  $F \in \mathcal{I}_1 \cap \mathcal{I}_2$  with  $|F| = k$ . We will analyze a natural Markov chain for the problem by analyzing the down-up walk of a corresponding simplicial complex.

Define the  $(k - 1)$ -dimensional matroid intersection complex  $C_{M_1, M_2, k}$  of  $M_1 = (E, \mathcal{I}_1)$  and  $M_2 = (E, \mathcal{I}_2)$  as

$$C_{M_1, M_2, k} = \{S \in \mathcal{I}_1 \cap \mathcal{I}_2 : |S| \leq k\},$$

the complex consisting of all common independent sets of both matroids containing at most  $k$  elements. We endow  $C_{M_1, M_2, k}(k - 1)$  with the uniform distribution  $\Pi_{k-1}$  on the common independent sets  $S \in \mathcal{I}_1 \cap \mathcal{I}_2$  with  $|S| = k$ . We write  $C_{M_1, M_2, k}$  for the weighted simplicial complex  $(C_{M_1, M_2, k}, \Pi_{k-1})$ .

The  $(k - 1)$ -th down-up walk  $P_{k-1}^\nabla$  on  $C_{M_1, M_2, k}$  corresponds to the following natural Markov chain to sample common independent sets of size  $k$ . Initially, the random walk starts from a common independent set  $S_1$  of size  $k$ . In each step  $t \geq 1$ , we choose a uniform random element  $i \in S_t$  and delete  $i$  from  $S_t$ , and set  $S_{t+1}$  to be a uniform random common independent set of size  $k$  that contains  $S_t \setminus \{i\}$ . The stationary distribution of  $P_{k-1}^\nabla$  is the uniform distribution  $\Pi_{k-1}$ ; see [Section 2](#).

The main result in this subsection is the following upper bound on the second eigenvalue of  $P_{k-1}^\nabla$ .

**Theorem 1.8.** *Let  $M_1 = (E, \mathcal{I}_1)$  and  $M_2 = (E, \mathcal{I}_2)$  be two given partition matroids with a common independent set of size  $r$  and no two elements belonging to the same block in both matroids. If  $k \leq r/3$ , then*

$$\lambda_2(P_{k-1}^\nabla) \leq 1 - \frac{1}{k^2},$$

where  $P_{k-1}^\nabla$  is the  $(k - 1)$ -th down-up walk on the matroid intersection complex  $C_{M_1, M_2, k}$ .

We note again that stronger results are known to exist [[JS89](#), [JSV04](#)], which are established by canonical paths arguments. Our result is going to follow from simple eigenvalue interlacing arguments alongside with [Corollary 3.4](#).

**4.2.1 Proof of Theorem 1.8.** The plan is to use [Corollary 3.4](#) to prove [Theorem 1.8](#). To apply [Corollary 3.4](#), we need to prove that:

- (1)  $C_{M_1, M_2, k}$  is a pure simplicial complex. This is a simple proof in [Claim 4.4](#).
- (2) For each  $S \in C_{M_1, M_2, k}$  with  $|S| \leq k - 2$ , the random walk matrix  $M_S$  of the graph  $G_S$  of the link  $(C_{M_1, M_2, k})_S$  satisfies  $\lambda_2(M_S) < 1$ . This is proved in [Lemma 4.5](#), showing that the underlying graph of  $G_S$  is the complement of the line graph of a bipartite graph.
- (3) For each  $S \in C_{M_1, M_2, k}$  with  $|S| = k - 2$ , the random walk matrix  $M_S$  of the graph  $G_S$  satisfies  $\lambda_2(M_S) \leq 1/k$ . This is ensured by [Lemma 4.8](#), which is obtained by the fact that the minimum eigenvalue of the adjacency matrix of the line graph of a simple graph is at least  $-2$ .



Assuming the three items are proven, [Theorem 1.8](#) follows immediately from [Corollary 3.4](#).

It remains to prove the three items. We will prove the second item in [Section 4.2.2](#) and the third item in [Section 4.2.3](#). We note that the first two items hold for any two matroids, and we only use the additional assumptions for the third item. The following is a simple proof for the first item.

**Claim 4.4.** *Let  $M_1 = (E, \mathcal{I}_1)$  and  $M_2 = (E, \mathcal{I}_2)$  be two matroids with a common independent set  $T \in \mathcal{I}_1 \cap \mathcal{I}_2$  of size  $|T| = r$ . Any common independent set  $S \in \mathcal{I}_1 \cap \mathcal{I}_2$  with  $|S| < r/2$  is contained in a larger common independent set. In particular, this implies that the simplicial complex  $C_{M_1, M_2, k}$  is a pure simplicial complex as long as  $k \leq r/2$ .*

**PROOF.** By the extension property of matroids, there is a subset  $T_1 \subset T$  with  $|T_1| \geq r - |S|$  such that  $S \cup \{x\}$  is an independent set in  $\mathcal{I}_1$  for any  $x \in T_1$ . Similarly, there is a subset  $T_2 \subset T$  with  $|T_2| \geq r - |S|$  such that  $S \cup \{y\}$  is an independent set in  $\mathcal{I}_2$  for any  $y \in T_2$ . As  $|S| < r/2$ , this implies that  $T_1 \cap T_2 \neq \emptyset$ , and  $S \cup \{z\}$  is a larger independent set that contains  $S$  for any  $z \in T_1 \cap T_2$ .  $\square$

**4.2.2 Proof of [Lemma 4.5](#).** Let  $H_S = (E_S, F_S)$  be the underlying support graph of  $G_S$  of the link  $(C_{M_1, M_2, k})_S$ , that is,  $H_S$  is  $G_S$  without edge weights. The vertex set of  $H_S$  is  $E_S = \{x \in E \mid S \cup \{x\} \in \mathcal{I}_1 \cap \mathcal{I}_2\}$  and the edge set of  $H_S$  is  $F_S = \{\{x, y\} \mid x, y \in E \text{ and } S \cup \{x, y\} \in \mathcal{I}_1 \cap \mathcal{I}_2\}$ . Let  $M_S$  be the random walk matrix of  $G_S$  as defined in [Section 2.3](#). It is a basic fact in spectral graph theory that  $\lambda_2(M_S) < 1$  if and only if  $H_S$  is connected.

We will see that  $H_S$  is the complement of the line graph of a bipartite graph  $B$ . To define the bipartite graph  $B$ , we first introduce the matroid partition property (see e.g. [\[ALOV19\]](#)). The matroid partition property says that there is a partition  $\mathcal{P} := \{P_1, \dots, P_p\}$  of the vertex set  $E_S$  (i.e.  $\bigcup_{i=1}^p P_i = E_S$  and  $P_i \cap P_j = \emptyset$  for  $i \neq j$ ) with the property that for any  $x, y \in E_S$ ,

$$S \cup \{x, y\} \notin \mathcal{I}_1 \iff x, y \in P_i \text{ for some } 1 \leq i \leq p.$$

In words, there is a partition  $\mathcal{P}$  of the vertex set  $E_S$  such that two elements  $x, y$  in  $E_S$  can be added to  $S$  to form an independent set in the first matroid  $M_1$  if and only if  $x, y$  do not belong to the same class of the partition  $\mathcal{P}$ . Similarly, there is a partition  $\mathcal{Q} := \{Q_1, \dots, Q_q\}$  of the vertex set  $E_S$  such that for any two elements  $x, y \in E_S$ , we have  $S \cup \{x, y\} \notin \mathcal{I}_2$  if and only if  $x, y \in Q_i$  for some  $1 \leq i \leq q$ .

We use the partitions  $\mathcal{P}$  and  $\mathcal{Q}$  to define the bipartite graph  $B$  as follows. The vertex set of  $B$  is  $P \sqcup Q$ , where we create a vertex  $i \in P$  in  $B$  for each  $P_i$  in  $\mathcal{P}$ , and we create a vertex  $j \in Q$  in  $B$  for each  $Q_j$  in  $\mathcal{Q}$ . Each edge in  $B$  corresponds to an element in  $E_S$ . For each element  $x \in E_S$ , we create the edge  $e_x = ij$  in  $B$  if and only if  $x \in P_i$  and  $x \in Q_j$ . Note that the edge  $e_x$  for  $x \in E_S$  is well-defined by the matroid partition property. By construction, it should be clear that the bipartite graph  $B$  satisfies the following important property:

$$\begin{aligned} e_x \text{ and } e_y \text{ do not share a vertex in } B &\iff S \cup \{x, y\} \in \mathcal{I}_1 \cap \mathcal{I}_2, \\ &\iff \{x, y\} \in F_S. \end{aligned} \quad (4.3)$$

Recall that the line graph  $L(B)$  of a graph  $B$  is defined as follows: the vertex set of  $L(B)$  is the edge set of  $B$ , and two vertices in  $L(B)$  have an edge if and only if the corresponding edges in  $B$  share an endpoint. Let  $\overline{L(B)}$  be the complement of  $L(B)$  where  $\overline{L(B)}$  and  $L(B)$

have the same vertex set and two vertices in  $\overline{L(B)}$  have an edge if and only if the corresponding vertices in  $L(B)$  do not have an edge. Then, we see from [Eq. \(4.3\)](#) that

$$H_S = \overline{L(B)}. \quad (4.4)$$

Using the bipartite graph  $B$ , it is easy to show the second item in [Section 4.2.1](#).

**Lemma 4.5.** *Let  $M_1 = (E, \mathcal{I}_1)$  and  $M_2 = (E, \mathcal{I}_2)$  be two matroids with a common independent set  $T \in \mathcal{I}_1 \cap \mathcal{I}_2$  of size  $|T| = r$ . Suppose  $k < r/2 - 1$ . For any  $S \in C_{M_1, M_2, k}$  with  $|S| \leq k - 2$ , the random walk matrix  $M_S$  of the graph  $G_S$  of the link  $(C_{M_1, M_2, k})_S$  satisfies  $\lambda_2(M_S) < 1$ .*

**PROOF.** It is well known that  $\lambda_2(M_S) < 1$  if and only if the underlying support graph  $H_S$  of  $G_S$  is connected, so we focus on proving the latter. Since  $|S| \leq k - 2 < r/2 - 3$ , it follows from [Claim 4.4](#) that there are four elements  $a, b, c, d \in E$  such that  $S \cup \{a, b, c, d\} \in \mathcal{I}_1 \cap \mathcal{I}_2$ . In the bipartite graph  $B$  in [Eq. \(4.4\)](#), the four elements  $a, b, c, d$  correspond to four vertex-disjoint edges  $e_a, e_b, e_c, e_d$  in  $B$  by [Eq. \(4.3\)](#). To prove that  $H_S$  is connected, we prove the stronger claim that every two vertices  $u, v \in H_S$  has a path of length at most two. If  $uv$  is an edge in  $H_S$ , then there is a path of length one. Suppose  $uv$  is not an edge in  $H_S$ . Then  $e_u$  and  $e_v$  shares a vertex in  $B$  and so they span at most three vertices in  $B$ . This implies that  $e_u \cup e_v$  cannot intersect all four (vertex-disjoint) edges  $e_a, e_b, e_c, e_d$ . So there must be an edge, say  $e_a$ , which is vertex-disjoint from both  $e_u$  and  $e_v$ . Then  $u-a-v$  is path of length two in  $H_S$  by [Eq. \(4.3\)](#), which completes the proof.  $\square$

**4.2.3 Proof of [Lemma 4.8](#).** For the third term, we need to prove that for each  $S \in C_{M_1, M_2, k}$  with  $|S| = k - 2$ , the random walk matrix  $M_S$  of the graph  $G_S$  satisfies  $\lambda_2(M_S) \leq \frac{1}{k}$ . We use the additional assumptions for the following property.

**Claim 4.6.** *If  $M_1$  and  $M_2$  are two partition matroids and there are no two elements  $x, y$  such that  $x, y$  belongs to the same block in  $M_1$  and also the same block in  $M_2$ , then [Eq. \(4.4\)](#) holds with the property that the bipartite graph  $B$  is a simple graph.*

Observe that  $G_S$  is an unweighted graph for  $S$  with  $|S| = k - 2$  when the distribution on  $C_{M_1, M_2, k}(k-1)$  is the uniform distribution (i.e. the distribution on the common independent sets of size  $k$  is the uniform distribution). This is because when  $|S| = k - 2$ , for any  $x, y \in E$ , either  $S \cup \{x, y\}$  is contained in exactly one or zero set of size  $k$  in  $C_{M_1, M_2, k}$ , and each set of size  $k$  is assigned the same weight in the uniform distribution (more formally see [Eq. \(2.3\)](#) for the definition of the weight). Therefore,  $G_S$  is simply a scaled version of  $H_S$ , and the random walk matrix  $M_S$  of  $G_S$  is the same as the random walk matrix of  $H_S$ .

**Fact 4.7.** *Let  $G = (V, E)$  be any simple graph and  $A_{L(G)}$  be the adjacency matrix of the line graph of  $G$ . One has,  $\lambda_{\min}(A_{L(G)}) \geq -2$ .*

**PROOF.** Define  $B \in \mathbb{R}^{E \times V}$  to be the edge-vertex incidence matrix of  $G = (V, E)$ , i.e.  $B(e, v) = 1[v \in e]$ . Observe that  $2I + A_{L(G)} = BB^T$  and as  $BB^T$  is a positive semidefinite matrix,  $\lambda_{\min}(A_{L(G)}) \geq -2$ .  $\square$

We need the stronger assumption that  $k \leq \frac{r}{3}$  for establishing the following lemma, whose proof omit it here, due to the many

similarities the proof bears with Lemma 4.3. We refer the reader to the full version of this paper for the proof.

**Lemma 4.8.** *Let  $M_1 = (E, \mathcal{I}_1)$  and  $M_2 = (E, \mathcal{I}_2)$  be two partition matroids with a common independent set  $T \in \mathcal{I}_1 \cap \mathcal{I}_2$  of size  $|T| = r$  and there are no two elements belonging to the same block in both matroids. Suppose  $k \leq r/3$ . For any  $S \in C_{M_1, M_2, k}$  with  $|S| = k - 2$ , the random walk matrix  $M_S$  of the graph  $G_S$  of the link  $(C_{M_1, M_2, k})_S$  satisfies  $\lambda_2(M_S) \leq 1/k$ .*

## ACKNOWLEDGEMENTS

We thank Kuikui Liu, Shayan Oveis-Gharan, Akshay Ramachandran, and Hong Zhou for many useful discussions.

## REFERENCES

- [AB06] Ron Aharoni and Eli Berger. The intersection of a matroid and a simplicial complex. *Transactions of the American Mathematical Society*, 358(11):4895–4917, 2006.
- [AHK18] Karim Adiprasito, June Huh, and Eric Katz. Hodge theory for combinatorial geometries. *Annals of Mathematics*, 188(2):381–452, 2018.
- [AJQ<sup>+</sup>20] Vedat Levi Alev, Fernando Granha Jeronimo, Dylan Quintana, Shashank Srivastava, and Madhur Tulsiani. List decoding of direct sum codes. In *SODA*, 2020.
- [AJT19] Vedat Levi Alev, Fernando Granha Jeronimo, and Madhur Tulsiani. Approximating constraint satisfaction problems on high dimensional expanders. In *FOCS*, 2019.
- [Ald83] David Aldous. Random walks on finite groups and rapidly mixing markov chains. In *Séminaire de Probabilités XVII 1981/82*, pages 243–297. Springer, 1983.
- [ALO] Nima Anari, Kuikui Liu, and Shayan Oveis Gharan. Spectral Independence in High-Dimensional Expanders and Applications to the Hardcore Model (Manuscript).
- [ALOV19] Nima Anari, Kuikui Liu, Shayan Oveis Gharan, and Cynthia Vinzant. Log-concave polynomials ii: high-dimensional walks and an fpras for counting bases of a matroid. In *STOC*, pages 1–12. ACM, 2019.
- [BD97] Russ Bubley and Martin E. Dyer. Path coupling: A technique for proving rapid mixing in markov chains. In *FOCS*, pages 223–231, 1997.
- [BH19] Petter Brändén and June Huh. Lorentzian polynomials. *arXiv preprint arXiv:1902.03719*, 2019.
- [Bha13] Rajendra Bhatia. *Matrix analysis*, volume 169. Springer Science & Business Media, 2013.
- [BT06] Sergey G Bobkov and Prasad Tetali. Modified logarithmic sobolev inequalities in discrete settings. *Journal of Theoretical Probability*, 19(2):289–336, 2006.
- [CGM19] Mary Cryan, Heng Guo, and Giorgos Mousa. Modified log-sobolev inequalities for strongly log-concave distributions. *CoRR*, abs/1903.06081, 2019.
- [DD19] Yotam Dikstein and Irit Dinur. Agreement testing theorems on layered set systems. 2019.
- [DDFH18] Yotam Dikstein, Irit Dinur, Yuval Filmus, and Praladh Harsha. Boolean function analysis on high-dimensional expanders. In *APPROX/RANDOM*, pages 38:1–38:20, 2018.
- [DFK91] Martin Dyer, Alan Frieze, and Ravi Kannan. A random polynomial-time algorithm for approximating the volume of convex bodies. *Journal of the ACM (JACM)*, 38(1):1–17, 1991.
- [DHK<sup>+</sup>19] Irit Dinur, Praladh Harsha, Tali Kaufman, Inbal Livni Navon, and Amnon Ta-Shma. List decoding with double samplers. In *SODA*, pages 2134–2153, 2019.
- [DK17] Irit Dinur and Tali Kaufman. High dimensional expanders imply agreement expanders. In *FOCS*, pages 974–985, 2017.
- [DKW16] Dominic Dottererr, Tali Kaufman, and Uli Wagner. On expansion and topological overlap. In *SOCC*, pages 35:1–35:10, 2016.
- [DSC<sup>+</sup>96] Persi Diaconis, Laurent Saloff-Coste, et al. Logarithmic sobolev inequalities for finite markov chains. *The Annals of Applied Probability*, 6(3):695–750, 1996.
- [EK16] Shai Evra and Tali Kaufman. Bounded degree cosystolic expanders of every dimension. In *STOC*, pages 36–48, 2016.
- [FGL<sup>+</sup>11] Jacob Fox, Mikhail Gromov, Vincent Lafforgue, Assaf Naor, and János Pach. Overlap properties of geometric expanders. In *SODA*, pages 1188–1197, 2011.
- [FM92] Tom Feder and Milena Mihail. Balanced matroids. In *Proceedings of the Twenty Fourth Annual ACM Symposium on the Theory of Computing*, pages 26–38. Citeseer, 1992.
- [God] Chris Godsil. Mathematics stackexchange url: <https://math.stackexchange.com/questions/2527245/what-graph-have-exactly-one-positive-eigenvalue>.
- [Gro10] Mikhail Gromov. Singularities, expanders and topology of maps. part 2: From combinatorics to topology via algebraic isoperimetry. *Geometric and Functional Analysis*, 20(2):416–526, 2010.
- [Hay06] Thomas P Hayes. A simple condition implying rapid mixing of single-site dynamics on spin systems. In *FOCS*, pages 39–46. IEEE, 2006.
- [HLW06] Shlomo Hoory, Nathan Linial, and Avi Wigderson. Expander graphs and their applications. *Bulletin of the American Mathematical Society*, 43(4):439–561, 2006.
- [HW17] June Huh and Botong Wang. Enumeration of points, lines, planes, etc. *Acta Mathematica*, 218(2):297–317, 2017.
- [Jer95] Mark Jerrum. A very simple algorithm for estimating the number of  $k$ -colorings of a low-degree graph. *Random Struct. Algorithms*, 7(2):157–166, 1995.
- [JS89] Mark Jerrum and Alistair Sinclair. Approximating the permanent. *SIAM J. Comput.*, 18(6):1149–1178, 1989.
- [JST<sup>+</sup>04] Mark Jerrum, Jung-Bae Son, Prasad Tetali, Eric Vigoda, et al. Elementary bounds on poincaré and log-sobolev constants for decomposable markov chains. *The Annals of Applied Probability*, 14(4):1741–1765, 2004.
- [JSV04] Mark Jerrum, Alistair Sinclair, and Eric Vigoda. A polynomial-time approximation algorithm for the permanent of a matrix with nonnegative entries. *J. ACM*, 51(4):671–697, 2004.
- [JVW86] Mark R Jerrum, Leslie G Valiant, and Vijay V Vazirani. Random generation of combinatorial structures from a uniform distribution. *Theoretical Computer Science*, 43:169–188, 1986.
- [KKL16] Tali Kaufman, David Kazhdan, and Alexander Lubotzky. Isoperimetric inequalities for ramanujan complexes and topological expanders. *Geometric and Functional Analysis*, 26(1):250–287, 2016.
- [KL14] Tali Kaufman and Alexander Lubotzky. High dimensional expanders and property testing. In *ITCS*, pages 501–506, 2014.
- [KM16] Tali Kaufman and David Mass. Walking on the edge and cosystolic expansion. *CoRR*, abs/1606.01844, 2016.
- [KM17] Tali Kaufman and David Mass. High dimensional random walks and colorful expansion. In *ITCS*, pages 4:1–4:27, 2017.
- [KO18] Tali Kaufman and Izhar Oppenheim. High order random walks: Beyond spectral gap. In *APPROX/RANDOM*, pages 47:1–47:17, 2018.
- [LM06] Nathan Linial and Roy Meshulam. Homological connectivity of random 2-complexes. *Combinatorica*, 26(4):475–487, 2006.
- [LMY19] Siqi Liu, Sidhanth Mohanty, and Elizabeth Yang. High-dimensional expanders from expanders. *CoRR*, abs/1907.10771, 2019.
- [LSV05] Alexander Lubotzky, Beth Samuels, and Uzi Vishne. Explicit constructions of ramanujan complexes of type. *Eur. J. Comb.*, 26(6):965–993, 2005.
- [LV06] László Lovász and Santosh Vempala. Simulated annealing in convex bodies and an  $o^*(n^4)$  volume algorithm. *Journal of Computer and System Sciences*, 72(2):392–417, 2006.
- [Mes01] Roy Meshulam. The clique complex and hypergraph matching. *Combinatorica*, 21(1):89–94, 2001.
- [MT05] Ravi Montenegro and Prasad Tetali. Mathematical aspects of mixing times in markov chains. *Foundations and Trends in Theoretical Computer Science*, 1(3), 2005.
- [MU05] Michael Mitzenmacher and Eli Upfal. *Probability and computing - randomized algorithms and probabilistic analysis*. Cambridge University Press, 2005.
- [MV87] Milena Mihail and Umesh Vazirani. On the expansion of 0/1 polytopes. *Journal of Combinatorial Theory*, 1987.
- [MW09] R. Meshulam and N. Wallach. Homological connectivity of random  $k$ -dimensional complexes. *Random Struct. Algorithms*, 34(3):408–417, 2009.
- [Opp18] Izhar Oppenheim. Local spectral expansion approach to high dimensional expanders part I: descent of spectral gaps. *Discrete & Computational Geometry*, 59(2):293–330, 2018.
- [PRT16] Ori Parzanchevski, Ron Rosenthal, and Ran J. Tessler. Isoperimetric inequalities in simplicial complexes. *Combinatorica*, 36(2):195–227, 2016.
- [Sch03] Alexander Schrijver. *Combinatorial optimization: polyhedra and efficiency*, volume 24. Springer Science & Business Media, 2003.
- [Sin92] Alistair Sinclair. Improved bounds for mixing rates of markov chains and multicommodity flow. *Combinatorics, probability and Computing*, 1(4):351–370, 1992.
- [Sin93] Alistair Sinclair. Algorithms for random generation and counting. progress in theoretical computer science, 1993.
- [Vig00] Eric Vigoda. Improved bounds for sampling colorings. *Journal of Mathematical Physics*, 41(3):1555–1569, 2000.
- [Wei06] Dror Weitz. Counting independent sets up to the tree threshold. In *STOC*, pages 140–149. ACM, 2006.
- [WLP09] EL Wilmer, David A Levin, and Yuval Peres. Markov chains and mixing times. *American Mathematical Soc., Providence*, 2009.