

CS475/CS675

Lecture 2: May 3, 2016

Cholesky factorization, tridiagonal,
band matrices

Reading: [TB] Chapt. 23 p. 172-176

Special Linear Systems

- Exploit special structures of linear systems
- More efficient LU factorization

- Symmetric systems
 - LDM^T factorization (variant of LU)
- Symmetric positive definite systems
 - GG^T factorization (a.k.a. Cholesky factorization)

LDM^T factorization

- Theorem: If all the leading principal submatrices of A are nonsingular, then there exist unique unit lower Δ matrices L and M , and a unique diagonal matrix D such that $A = LDM^T$.
- Partial Proof:
 - Factor $A = LU$
 - Define $D = \text{diag}(d_1, \dots, d_n)$, $d_i = u_{ii}$ $i = 1, \dots, n$
 - Let $M^T = D^{-1}U = \text{unit upper } \Delta$ ($M = \text{unit lower } \Delta$)
 - Thus $A = LU = LD(D^{-1}U) = LDM^T$
- Note: $\text{flops}(LU) = \text{flops}(LDM^T)$

Symmetric systems

- Theorem: If A is symmetric, then $A = LDL^T$
- Proof:
 - By previous result, $A = LDM^T$
 $\implies M^{-1}AM^{-T} = M^{-1}LDM^T M^{-T} = M^{-1}LD$
 - Since $M^{-1}AM^{-T}$ is symmetric, so is $M^{-1}LD$
 - Also, $M^{-1}L$ is lower $\Delta \implies M^{-1}LD$ is lower Δ
 - So $M^{-1}LD$ is both lower Δ and symmetric
 $\implies M^{-1}LD$ is diag $\implies M^{-1}L$ is diag
 - Since $M^{-1}L$ is also unit lower Δ ,
then $M^{-1}L = I \implies M = L$

Symmetric systems

- Notes

1. We can save about half the work by computing L and D only.
2. One way is to compute the U factor only during the LU factorization.

Positive definite systems

- Definition: A is positive definite iff $x^T A x > 0$ for all $x \neq 0$.
- Properties of positive definite matrices:

Positive definite systems

- Theorem: If $A \in \mathfrak{R}^{n \times n}$ is PD and $X \in \mathfrak{R}^{n \times k}$ has rank k , then $B = X^T A X \in \mathfrak{R}^{k \times k}$ is also PD
- Proof:
 - Let $z \in \mathfrak{R}^{k \times 1}$. Then $z^T B z = z^T X^T A X z = (X z)^T A (X z)$
 - If $X z = 0$, then X is not rank k .
 - Hence $z^T B z > 0$.
- Corollary: If A is PD, then all its principal submatrices are *PD*. In particular, all diag entries are positive.

Positive definite systems

- Corollary: If A is PD, then $A = LDM^T$ and D has positive diag entries.
- Proof:
 - Let $X = L^{-T}$.
Then $X^T AX = L^{-1}(LDM^T)L^{-T} = DM^T L^{-T}$ is PD.
 - By previous corollary, $\text{diag}(DM^T L^{-T})$ has positive entries.
 - Note that M^T and L^{-T} are unit upper Δ .
 $\Rightarrow M^T L^{-T}$ is also unit upper Δ
 $\Rightarrow \text{diag}(DM^T L^{-T}) = D$

Symmetric positive definite systems

- Theorem: If A is SPD, then there exists unique lower Δ G such that

$$A = GG^T$$

- Proof:

– $A = LDL^T$ and $D = \text{diag}(d_1, \dots, d_n)$, $d_i > 0$.

– Define $D^{\frac{1}{2}} \equiv \text{diag}(\sqrt{d_1}, \dots, \sqrt{d_n})$

– Let $G = LD^{\frac{1}{2}}$. Then G is lower Δ

$$\Rightarrow GG^T = LD^{\frac{1}{2}}(LD^{\frac{1}{2}})^T = LD^{\frac{1}{2}}D^{\frac{1}{2}}L^T = LDL^T = A$$

Symmetric positive definite systems

- Examples

Cholesky factorization

- $A = GG^T$ is called the Cholesky factorization of A and the lower ΔG is called the Cholesky factor.

Cholesky factorization

- Algorithm big picture

Cholesky factorization

For $k = 1, 2, \dots, n$

$$a_{kk} = \sqrt{a_{kk}}$$

For $j = k + 1, \dots, n$

$$a_{jk} = a_{jk} / a_{kk}$$

End

For $j = k + 1, \dots, n$

for $i = j, \dots, n$

$$a_{ij} = a_{ij} - a_{ik}a_{jk}$$

end

End

End

$$\text{flops}(\text{Cholesky}) \approx \frac{n^3}{3}$$

Banded systems

- Definition: A has upper bandwidth q if $a_{ij} = 0 \forall j > i + q$ and lower bandwidth p if $a_{ij} = 0 \forall i > j + p$.
- Picture

Banded systems

- If A is banded, so are LU, GG^T, LDM^T
- Theorem: Let $A = LU$. If A has upper bandwidth q and lower bandwidth p , then U has upper bandwidth q and L has lower bandwidth p .
- Picture

Band Gaussian Elimination

For $k = 1, 2, \dots, n - 1$

For $i = k + 1, \dots, \min(k + p, n)$

$$a_{ik} = a_{ik} / a_{kk}$$

end

for $i = k + 1, \dots, \min(k + p, n)$

for $j = k + 1, \dots, \min(k + q, n)$

$$a_{ij} = a_{ij} - a_{ik} a_{kj}$$

end

end

End

If $n \gg p$ and $n \gg q$, then $flops(\text{band GE}) \approx 2npq$

Tridiagonal systems

- Assume A is tridiagonal and symmetric
- Then

$$L =$$

$$D =$$

Tridiagonal System

- $A = LDL^T$ implies

$$\begin{aligned} a_{kk} &= (LDL^T)_{kk} \\ &= \sum_i \sum_j l_{ki} d_{ij} l_{jk}^T \\ &= \sum_i l_{ki} d_{ii} l_{ik}^T \\ &= \sum_i l_{ki}^2 d_{ii} \\ &= l_{k,k-1}^2 d_{k-1,k-1} + l_{kk}^2 d_{kk} \\ &\quad (i = k - 1) \\ &= l_{k-1}^2 d_{k-1} + d_k \end{aligned}$$

$$\begin{aligned} a_{k,k-1} &= \sum_i \sum_j l_{ki} d_{ij} l_{j,k-1}^T \\ &= \sum_i l_{ki} d_{ii} l_{i,k-1}^T \\ &= \sum_i l_{ki} d_{ii} l_{k-1,i} \\ &= l_{k,k-1} d_{k-1,k-1} l_{k-1,k-1} \\ &\quad (i = k - 1) \\ &= l_{k-1} d_{k-1} \end{aligned}$$

Tridiagonal Factorization

- Algorithm

$$d_1 = a_{11}$$

for $k = 2, \dots, n$

$$l_{k-1} = a_{k,k-1}/d_{k-1}$$

$$d_k = a_{kk} - l_{k-1}a_{k,k-1} \quad (l_{k-1}d_{k-1} = a_{k,k-1})$$

end

- $flops(tridiag) = O(n)$