

Bayes Nets (continued)

[RN2] Section 14.4

[RN3] Section 14.4

CS 486/686

University of Waterloo

Lecture 8: May 24, 2017

Outline


- Inference in Bayes Nets
- Variable Elimination

Inference in Bayes Nets

- The independence sanctioned by D-separation (and other methods) allows us to compute prior and posterior probabilities quite effectively.
- We'll look at a few simple examples to illustrate. We'll focus on networks without *loops*. (A loop is a cycle in the underlying *undirected* graph. Recall the directed graph has no cycles.)

Simple Forward Inference (Chain)

- Computing marginal requires simple forward “propagation” of probabilities



$$P(J) = \sum_{M,ET} P(J, M, ET)$$

(marginalization)

$$P(J) = \sum_{M,ET} P(J|M,ET)P(M|ET)P(ET)$$

(chain rule)

$$P(J) = \sum_{M,ET} P(J|M)P(M|ET)P(ET)$$

(conditional independence)


$$P(J) = \sum_M P(J|M) \sum_{ET} P(M|ET)P(ET)$$

(distribution of sum)

Note: all (final) terms are CPTs in the BN
 Note: only ancestors of J considered

Simple Forward Inference (Chain)

- Same idea applies when we have "upstream" evidence



$$P(J|ET) = \sum_M P(J, M | ET)$$

(marginalisation)

$$P(J|ET) = \sum_M P(J | M, ET) P(M | ET)$$

(chain rule)

$$P(J|ET) = \sum_M P(J | M) P(M | ET)$$

(conditional independence)

Simple Forward Inference (Pooling)

- Same idea applies with multiple parents


$$\begin{aligned} P(F_{ev}) &= \sum_{Flu, M, TS, ET} P(F_{ev}, Flu, M, TS, ET) \\ &= \sum_{Flu, M, TS, ET} P(F_{ev} | Flu, M, TS, ET) P(Flu | M, TS, ET) \\ &\quad P(M | TS, ET) P(TS | ET) P(ET) \\ &= \sum_{Flu, M, TS, ET} P(F_{ev} | Flu, M) P(Flu | TS) P(M | ET) P(TS) P(ET) \\ &= \sum_{Flu, M} P(F_{ev} | Flu, M) \left[\sum_{TS} P(Flu | TS) P(TS) \right] \\ &\quad \left[\sum_{ET} P(M | ET) P(ET) \right] \end{aligned}$$

- (1) by marginalisation; (2) by the chain rule;
(3) by conditional independence; (4) by distribution
 - note: all terms are CPTs in the Bayes net

Simple Forward Inference (Pooling)

- Same idea applies with evidence

$$\begin{aligned} P(\text{Fev} | \text{ts}, \sim m) &= \sum_{\text{Flu}} P(\text{Fev}, \text{Flu} | \text{ts}, \sim m) \\ &= \sum_{\text{Flu}} P(\text{Fev} | \text{Flu}, \text{ts}, \sim m) P(\text{Flu} | \text{ts}, \sim m) \\ &= \sum_{\text{Flu}} P(\text{Fev} | \text{Flu}, \sim m) P(\text{Flu} | \text{ts}) \end{aligned}$$



Simple Backward Inference

- When evidence is downstream of query variable, we must reason "backwards." This requires the use of Bayes rule:

$$\begin{aligned}P(ET \mid j) &= \alpha P(j \mid ET) P(ET) \\ &= \alpha \sum_M P(j, M \mid ET) P(ET) \\ &= \alpha \sum_M P(j \mid M, ET) P(M \mid ET) P(ET) \\ &= \alpha \sum_M P(j \mid M) P(M \mid ET) P(ET)\end{aligned}$$

- First step is just Bayes rule
 - normalizing constant α is $1/P(j)$; but we needn't compute it explicitly if we compute $P(ET \mid j)$ for each value of ET : we just add up terms $P(j \mid ET) P(ET)$ for all values of ET (they sum to $P(j)$)

Backward Inference (Pooling)

- Same ideas when several pieces of evidence lie "downstream"

$$P(ET|j,fev) = \alpha P(j,fev|ET) P(ET)$$

$$= \alpha \sum_{M,FI,TS} P(j,fev,M,FI,TS|ET) P(ET)$$

$$= \alpha \sum_{M,FI,TS} P(j|fev,M,FI,TS,ET) P(fev|M,FI,TS,ET) \\ P(M|FI,TS,ET) P(FI|TS,ET) P(TS|ET) P(ET)$$

$$= \alpha P(ET) \sum_M P(j|M) P(M|ET) \sum_{FI} P(fev|M,FI) \sum_{TS} \\ P(FI|TS) P(TS)$$

- Same steps as before; but now we compute prob of both pieces of evidence given hypothesis ET and combine them. Note: they are independent given M; but not given ET.

Variable Elimination

- The intuitions in the above examples give us a simple inference algorithm for networks without loops: the *polytree* algorithm.
- Instead we'll look at a more general algorithm that works for general BNs; but the polytree algorithm will be a special case.
- The algorithm, *variable elimination*, simply applies the summing out rule repeatedly.
 - To keep computation simple, it exploits the independence in the network and the ability to distribute sums inward

Factors

- A function $f(X_1, X_2, \dots, X_k)$ is also called a *factor*. We can view this as a table of numbers, one for each instantiation of the variables X_1, X_2, \dots, X_k .
 - A tabular rep'n of a factor is exponential in k
- Each CPT in a Bayes net is a factor:
 - e.g., $\Pr(C|A,B)$ is a function of three variables, A, B, C
- Notation: $f(\mathbf{X}, \mathbf{Y})$ denotes a factor over the variables $\mathbf{X} \cup \mathbf{Y}$. (Here \mathbf{X}, \mathbf{Y} are sets of variables.)

The Product of Two Factors

- Let $f(X,Y)$ & $g(Y,Z)$ be two factors with variables Y in common
- The *product* of f and g , denoted $h = f \times g$ (or sometimes just $h = fg$), is defined:

$$h(X,Y,Z) = f(X,Y) \times g(Y,Z)$$

f(A,B)		g(B,C)		h(A,B,C)			
ab	0.9	bc	0.7	abc	0.63	ab~c	0.27
a~b	0.1	b~c	0.3	a~bc	0.08	a~b~c	0.02
~ab	0.4	~bc	0.8	~abc	0.28	~ab~c	0.12
~a~b	0.6	~b~c	0.2	~a~bc	0.48	~a~b~c	0.12

Summing a Variable Out of a Factor

- Let $f(X,Y)$ be a factor with variable X (Y is a set)
- We *sum out* variable X from f to produce a new factor $h = \sum_X f$, which is defined:

$$h(Y) = \sum_{x \in \text{Dom}(X)} f(x,Y)$$

$f(A,B)$		$h(B)$	
ab	0.9	b	1.3
a~b	0.1	~b	0.7
~ab	0.4		
~a~b	0.6		


Restricting a Factor

- Let $f(X,Y)$ be a factor with variable X (Y is a set)
- We *restrict* factor f to $X=x$ by setting X to the value x and "deleting". Define $h = f_{X=x}$ as: $h(Y) = f(x,Y)$

$f(A,B)$		$h(B) = f_{A=a}$	
ab	0.9	b	0.9
a~b	0.1	~b	0.1
~ab	0.4		
~a~b	0.6		

Variable Elimination: No Evidence

- Computing prior probability of query var X can be seen as applying these operations on factors




- $$\begin{aligned} P(C) &= \sum_{A,B} P(C|B) P(B|A) P(A) \\ &= \sum_B P(C|B) \sum_A P(B|A) P(A) \\ &= \sum_B f_3(B,C) \sum_A f_2(A,B) f_1(A) \\ &= \sum_B f_3(B,C) f_4(B) = f_5(C) \end{aligned}$$

Define new factors: $f_4(B) = \sum_A f_2(A,B) f_1(A)$ and $f_5(C) = \sum_B f_3(B,C) f_4(B)$


Variable Elimination: No Evidence

- Here's the example with some numbers



$f_1(A)$		$f_2(A,B)$		$f_3(B,C)$		$f_4(B)$		$f_5(C)$	
a	0.9	ab	0.9	bc	0.7	b	0.85	c	0.625
$\sim a$	0.1	$a\sim b$	0.1	$b\sim c$	0.3	$\sim b$	0.15	$\sim c$	0.375
		$\sim ab$	0.4	$\sim bc$	0.2				
		$\sim a\sim b$	0.6	$\sim b\sim c$	0.8				

VE: No Evidence (Example 2)



$$\begin{aligned} P(D) &= \sum_{A,B,C} P(D|C) P(C|B,A) P(B) P(A) \\ &= \sum_C P(D|C) \sum_B P(B) \sum_A P(C|B,A) P(A) \\ &= \sum_C f_4(C,D) \sum_B f_2(B) \sum_A f_3(A,B,C) f_1(A) \\ &= \sum_C f_4(C,D) \sum_B f_2(B) f_5(B,C) \\ &= \sum_C f_4(C,D) f_6(C) \\ &= f_7(D) \end{aligned}$$

Define new factors: $f_5(B,C)$, $f_6(C)$, $f_7(D)$, in the obvious way

Variable Elimination: One View

- One way to think of variable elimination:
 - write out desired computation using the chain rule, exploiting the independence relations in the network
 - arrange the terms in a convenient fashion
 - distribute each sum (over each variable) in as far as it will go
 - i.e., the sum over variable X can be "pushed in" as far as the "first" factor mentioning X
 - apply operations "inside out", repeatedly eliminating and creating new factors (note that each step/removal of a sum eliminates one variable)

Variable Elimination Algorithm

- Given query var Q , remaining vars Z . Let F be the set of factors corresponding to CPTs for $\{Q\} \cup Z$.

1. Choose an elimination ordering Z_1, \dots, Z_n of variables in Z .
2. For each Z_j -- in the order given -- eliminate $Z_j \in Z$ as follows:
 - (a) Compute new factor $g_j = \sum_{Z_j} f_1 \times f_2 \times \dots \times f_k$,
where the f_i are the factors in F that include Z_j
 - (b) Remove the factors f_i (that mention Z_j) from F
and add new factor g_j to F
3. The remaining factors refer only to the query variable Q .
Take their product and normalize to produce $P(Q)$


VE: Example 2 again

Factors: $f_1(A)$ $f_2(B)$

$f_3(A,B,C)$ $f_4(C,D)$

Query: $P(D)$?

Elim. Order: A, B, C



Step 1: Add $f_5(B,C) = \sum_A f_3(A,B,C) f_1(A)$

Remove: $f_1(A), f_3(A,B,C)$

Step 2: Add $f_6(C) = \sum_B f_2(B) f_5(B,C)$

Remove: $f_2(B), f_5(B,C)$


Step 3: Add $f_7(D) = \sum_C f_4(C,D) f_6(C)$

Remove: $f_4(C,D), f_6(C)$

Last factor $f_7(D)$ is (possibly unnormalized) probability $P(D)$

Variable Elimination: Evidence

- Computing posterior of query variable given evidence is similar; suppose we observe $C=c$:



$$\begin{aligned} P(A|c) &= \alpha P(A) P(c|A) \\ &= \alpha P(A) \sum_B P(c|B) P(B|A) \\ &= \alpha f_1(A) \sum_B f_3(B,c) f_2(A,B) \\ &= \alpha f_1(A) \sum_B f_4(B) f_2(A,B) \\ &= \alpha f_1(A) f_5(A) \\ &= \alpha f_6(A) \end{aligned}$$

New factors: $f_4(B) = f_3(B,c)$; $f_5(A) = \sum_B f_2(A,B) f_4(B)$;
 $f_6(A) = f_1(A) f_5(A)$


Variable Elimination with Evidence

Given query var Q , evidence vars E
(observed to be e), remaining vars Z .
Let F be set of factors involving CPTs
for $\{Q\} \cup Z$.

1. Replace each factor $f \in F$ that mentions a variable(s) in E with its restriction $f_{E=e}$ (somewhat abusing notation)
2. Choose an elimination ordering Z_1, \dots, Z_n of variables in Z .
3. Run variable elimination as above.
4. The remaining factors refer only to the query variable Q .
Take their product and normalize to produce $P(Q)$

VE: Example 2 again with Evidence

Factors: $f_1(A)$ $f_2(B)$
 $f_3(A,B,C)$ $f_4(C,D)$
Query: $P(A)?$
Evidence: $D = d$
Elim. Order: C, B



Restriction: replace $f_4(C,D)$ with $f_5(C) = f_4(C,d)$

Step 1: Add $f_6(A,B) = \sum_C f_5(C) f_3(A,B,C)$

Remove: $f_3(A,B,C), f_5(C)$

Step 2: Add $f_7(A) = \sum_B f_6(A,B) f_2(B)$

Remove: $f_6(A,B), f_2(B)$

Last factors: $f_7(A), f_1(A)$. The product $f_1(A) \times f_7(A)$ is (possibly unnormalized) posterior. So... $P(A|d) = \alpha f_1(A) \times f_7(A)$.

Some Notes on the VE Algorithm


- After iteration j (elimination of Z_j), factors remaining in set F refer only to variables X_{j+1}, \dots, Z_n and Q . No factor mentions an evidence variable E after the initial restriction.
- Number of iterations: linear in number of variables
- Complexity is exponential in the number of variables.
 - Recall each factor has exponential size in its number of variables
 - Can't do any better than size of BN (since its original factors are part of the factor set)
 - When we create new factors, we might make a set of variables larger.

Some Notes on the VE Algorithm

- The size of the resulting factors is determined by elimination ordering! (We'll see this in detail)
- For *polytrees*, easy to find good ordering (e.g., work outside in).
- For general BNs, sometimes good orderings exist, sometimes they don't (then inference is exponential in number of vars).
 - Simply *finding* the optimal elimination ordering for general BNs is NP-hard.
 - Inference in general is NP-hard in general BNs


Elimination Ordering: Polytrees

- Inference is linear in size of network
 - ordering: eliminate only "singly-connected" nodes
 - e.g., in this network, eliminate D, A, C, X_1, \dots ; or eliminate X_1, \dots, X_k, D, A, C ; or mix up...
 - result: no factor ever larger than original CPTs
 - eliminating B before these gives factors that include all of A, C, X_1, \dots, X_k !!!




Effect of Different Orderings

- Suppose query variable is D. Consider different orderings for this network
 - A,F,H,G,B,C,E:
 - good: why?
 - E,C,A,B,G,H,F:
 - bad: why?
- Which ordering creates smallest factors?
 - either max size or total
- which creates largest factors?



Relevance



- Certain variables have no impact on the query.
 - In ABC network, computing $\Pr(A)$ with no evidence requires elimination of B and C.
 - But when you sum out these vars, you compute a trivial factor (whose value are all ones); for example:
 - eliminating C: $f_4(B) = \sum_C f_3(B,C) = \sum_C \Pr(C|B)$
 - 1 for any value of B (e.g., $\Pr(c|b) + \Pr(\sim c|b) = 1$)
- No need to think about B or C for this query

Relevance: A Sound Approximation

- Can restrict attention to *relevant* variables. Given query Q , evidence E :
 - Q is relevant
 - if any node Z is relevant, its parents are relevant
 - if $E \in E$ is a descendent of a relevant node, then E is relevant
- We can restrict our attention to the *subnetwork comprising only relevant variables* when evaluating a query Q

Relevance: Examples

- Query: $P(F)$
 - Relevant: F, C, B, A
- Query: $P(F|E)$
 - Relevant: F, C, B, A
 - Also: E , hence D, G
 - Intuitively, we need to compute $P(C|E) = \alpha P(C)P(E|C)$ to accurately compute $P(F|E)$
- Query: $P(F|E, C)$
 - Algorithm says all vars relevant; but really none except C, F since C cuts off all influence of others)
 - Algorithm is overestimating relevant set

