

# Lecture 5: Uncertainty

# CS486/686 Intro to Artificial Intelligence

2026-1-20

Pascal Poupart  
David R. Cheriton School of Computer Science

# Outline

- Probability theory
- Uncertainty via probabilities
- Probabilistic inference

# Terminology

- **Probability distribution:**
  - A specification of a probability for each event in our sample space
  - Probabilities must sum to 1
- Assume the world is described by two (or more) random variables
  - **Joint probability distribution**
    - Specification of probabilities for all combinations of events

# Joint distribution

- Given two random variables  $A$  and  $B$ :
- Joint distribution:

$\Pr(A = a \wedge B = b)$  for all  $a, b$

- **Marginalisation (sumout rule):**

$$\Pr(A = a) = \sum_b \Pr(A = a \wedge B = b)$$

$$\Pr(B = b) = \sum_a \Pr(A = a \wedge B = b)$$

# Example: Joint Distribution

sunny		~sunny
	cold	~cold
headache	0.108	0.012
~headache	0.016	0.064

$$P(\text{headache} \wedge \text{sunny} \wedge \text{cold}) = 0.108$$

$$P(\text{~headache} \wedge \text{sunny} \wedge \text{~cold}) = 0.064$$


$$P(\text{headache}) = 0.108 + 0.012 + 0.072 + 0.008 = 0.2$$



**marginalization**

# Conditional Probability

- $\Pr(A|B)$ : fraction of worlds in which  $B$  is true that also have  $A$  true




$H$  = “Have headache”  
 $F$  = “Have Flu”

$$\begin{aligned}\Pr(H) &= 1/10 \\ \Pr(F) &= 1/40 \\ \Pr(H|F) &= 1/2\end{aligned}$$

Headaches are rare and flu is rarer, but if you have the flu, then there is a 50-50 chance you will have a headache

# Conditional Probability



$H$  = “Have headache”  
 $F$  = “Have Flu”

$$\begin{aligned}\Pr(H) &= 1/10 \\ \Pr(F) &= 1/40 \\ \Pr(H|F) &= 1/2\end{aligned}$$

$$\begin{aligned}\Pr(H|F) &= \text{Fraction of flu inflicted worlds in which you have a headache} \\ &= (\# \text{ worlds with flu and headache}) / (\# \text{ worlds with flu}) \\ &= (\text{Area of “H and F” region}) / (\text{Area of “F” region}) \\ &= \Pr(H \wedge F) / \Pr(F)\end{aligned}$$

# Conditional Probability

- Definition:  $\Pr(A|B) = \Pr(A \wedge B) / \Pr(B)$


- Chain rule:  $\Pr(A \wedge B) = \Pr(A|B) \Pr(B)$

$$\Pr(A, B, C) = \Pr(A|B, C) \Pr(B, C)$$

$\Pr(B|C) \Pr(C)$

Memorize these rules!

# Inference



One day you wake up with a headache. You think “Drat! 50% of flues are associated with headaches so I must have a 50-50 chance of coming down with the flu”

$H$  = “Have headache”  
 $F$  = “Have Flu”

$$\begin{aligned}\Pr(H) &= 1/10 \\ \Pr(F) &= 1/40 \\ \Pr(H|F) &= 1/2\end{aligned}$$

Is your reasoning correct?

$$\begin{aligned}\Pr(F \wedge H) &= \Pr(F) \Pr(H|F) = (1/40)(1/2) = 1/80 \\ \Pr(F|H) &= \frac{\Pr(F, H)}{\Pr(H)} = \frac{1/80}{1/10} = 1/8\end{aligned}$$

# Example: Conditional Distribution

sunny		~sunny			
	cold	~cold			
headache	0.108	0.012	headache	0.072	0.008
~headache	0.016	0.064	~headache	0.144	0.576

$$\Pr(\text{headache} \wedge \text{cold} \mid \text{sunny}) = \frac{P(h, c, s)}{P(s)} = \frac{0.108}{0.108 + 0.012 + 0.016 + 0.064} = 0.54$$

$$\Pr(\text{headache} \wedge \text{cold} \mid \text{~sunny}) = \frac{P(h, c, \text{~s})}{P(\text{~s})} = \frac{0.072}{0.072 + 0.008 + 0.144 + 0.576} = 0.09$$

# Bayes Rule

- Note:  $\Pr(A|B)\Pr(B) = \Pr(A \wedge B) = \Pr(B \wedge A) = \Pr(B|A)\Pr(A)$
- Bayes Rule:  $\Pr(B|A) = \frac{\Pr(A|B)\Pr(B)}{\Pr(A)}$

**Memorize this!**

# Using Bayes' Rule for inference

- Often, we want to form a hypothesis about the world based on what we have observed
- Bayes' rule allows us to compute a belief about hypothesis  $H$ , given evidence  $e$

$$P(H|e) = \frac{P(e|H)P(H)}{P(e)}$$

Posterior probability

Likelihood

Prior probability

Normalizing constant

The diagram illustrates the components of Bayes' rule. The formula  $P(H|e) = \frac{P(e|H)P(H)}{P(e)}$  is shown. Red arrows point from the labels to the corresponding terms: 'Posterior probability' points to  $P(H|e)$ , 'Likelihood' points to  $P(e|H)$ , 'Prior probability' points to  $P(H)$ , and 'Normalizing constant' points to  $P(e)$ .

# More General Forms of Bayes Rule

$$P(A|B) = \frac{P(B|A)P(A)}{P(B|A)P(A) + P(B|\sim A)P(\sim A)}$$

$$P(A|B \wedge X) = \frac{P(B|A \wedge X)P(A|X)}{P(B|X)}$$

$$P(A = v_i|B) = \frac{P(B|A = v_i)P(A = v_i)}{\sum_{k=1}^n P(B|A = v_k)P(A = v_k)}$$

# Probabilistic Inference

- By probabilistic inference, we mean
  - given a *prior* distribution  $\Pr(X)$  over variables  $X$  of interest, representing degrees of belief
  - and given new evidence  $E = e$  for some variable  $E$
  - Revise your degrees of belief: *posterior*  $\Pr(X|E = e)$
- Applications:
  - Medicine:  $\Pr(disease|symptom1, symptom2, \dots, symptomN)$
  - Troubleshooting:  $\Pr(cause|test1, test2, \dots, testN)$

# Issues

- How do we specify the full joint distribution over a set of random variables  $X_1, X_2, \dots, X_n$ ?
  - Exponential number of possible worlds
  - e.g., if  $X_i$  is Boolean, then  $2^n$  numbers (or  $2^n - 1$  parameters, since they sum to 1)
  - These numbers are not robust/stable
- Inference is frightfully slow
  - Must sum over exponential number of worlds to answer queries
    - $\Pr(X_i) = \sum_{X_1} \dots \sum_{X_{i-1}} \sum_{X_{i+1}} \dots \sum_{X_n} \Pr(X_1, X_2, \dots, X_n)$
    - $\Pr(X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n | X_i) = \frac{\Pr(X_1, \dots, X_n)}{\Pr(X_i)} = \frac{\Pr(x_1, \dots, x_n)}{\sum_{X_1} \dots \sum_{X_{i-1}} \sum_{X_{i+1}} \dots \sum_{X_n} \Pr(X_1, \dots, X_n)}$

# Small Example: 3 Variables

sunny		$\sim$ sunny			
	cold	$\sim$ cold			
headache	0.108	0.012	headache	0.072	0.008
$\sim$ headache	0.016	0.064	$\sim$ headache	0.144	0.576

$$\Pr(\text{headache}) = 0.108 + 0.012 + 0.072 + 0.008 = 0.2$$

$$\begin{aligned}\Pr(\text{headache} \wedge \text{cold} | \text{sunny}) &= \Pr(\text{headache} \wedge \text{cold} \wedge \text{sunny}) / \Pr(\text{sunny}) \\ &= 0.108 / (0.108 + 0.012 + 0.016 + 0.064) = 0.54\end{aligned}$$

$$\begin{aligned}\Pr(\text{headache} \wedge \text{cold} | \sim \text{sunny}) &= \Pr(\text{headache} \wedge \text{cold} \wedge \sim \text{sunny}) / \Pr(\sim \text{sunny}) \\ &= 0.072 / (0.072 + 0.008 + 0.144 + 0.576) = 0.09\end{aligned}$$

# Intractable Inference

- How do we avoid the exponential blow up of joint distribution and probabilistic inference?
  - no solution in general
  - but in practice there is structure we can exploit
- We'll use **conditional independence**

# Independence

- Recall that  $X$  and  $Y$  are *independent* iff:

$$\begin{aligned}\Pr(X = x) &= \Pr(X = x | Y = y) \\ \Leftrightarrow \Pr(Y = y) &= \Pr(Y = y | X = x) \\ \Leftrightarrow \Pr(X = x, Y = y) &= \Pr(X = x) \Pr(Y = y) \\ \forall x \in \text{dom}(X), y \in \text{dom}(Y) &\end{aligned}$$

- Intuitively, learning the value of  $Y$  doesn't influence our beliefs about  $X$  and vice versa.
- Example:  $\Pr(\text{Sunny} | \text{ToothCavity}) = \Pr(\text{Sunny})$   
 $\Pr(\text{ToothCavity} | \text{Sunny}) = \Pr(\text{ToothCavity})$

# Conditional Independence

- Two *variables*  $X$  and  $Y$  are conditionally independent given variable  $Z$ 
$$\begin{aligned} \Pr(X = x|Z = z) &= \Pr(X = x|Y = y, Z = z) \\ \Leftrightarrow \Pr(Y = y|Z = z) &= \Pr(Y = y|X = x, Z = z) \\ \Leftrightarrow \Pr(X = x, Y = y|Z = z) &= \Pr(X = x|Z = z) \Pr(Y = y|Z = z) \\ \forall x \in \text{dom}(X), y \in \text{dom}(Y), z \in \text{dom}(Z) \end{aligned}$$
- If you know the value of  $Z$  (*whatever* it is), nothing you learn about  $Y$  will influence your beliefs about  $X$
- Example:  $\Pr(\text{ToothAche}|\text{ToothCavity}, \text{ToothCatch}) = \Pr(\text{ToothAche}|\text{ToothCavity})$   
 $\Pr(\text{ToothCatch}|\text{ToothCavity}, \text{ToothAche}) = \Pr(\text{ToothCatch}|\text{ToothCavity})$

# What good is independence?

- Suppose (say, Boolean) variables  $X_1, X_2, \dots, X_n$  are mutually independent
  - We can specify full joint distribution using only  $n$  parameters (linear) instead of  $2^n - 1$  (exponential)
- How? Simply specify  $\Pr(x_1), \dots, \Pr(x_n)$ 
  - From this we can recover the probability of any world or any (conjunctive) query easily
    - Recall  $\Pr(x_1, \dots, x_n) = \Pr(x_1) \dots \Pr(x_n)$

# Example

- 4 independent Boolean random vars  $X_1, X_2, X_3, X_4$

$$\Pr(x_1) = 0.4, \Pr(x_2) = 0.2, \Pr(x_3) = 0.5, \Pr(x_4) = 0.8$$

$$\begin{aligned}\Pr(x_1, \sim x_2, x_3, x_4) &= \Pr(x_1) (1 - \Pr(x_2)) \Pr(x_3) \Pr(x_4) \\ &= (0.4)(0.8)(0.5)(0.8) \\ &= 0.128\end{aligned}$$

$$\begin{aligned}\Pr(x_1, x_2, x_3 | x_4) &= \Pr(x_1) \Pr(x_2) \Pr(x_3) \mathbf{1} \\ &= (0.4)(0.2)(0.5)(1) \\ &= 0.04\end{aligned}$$

# The Value of Independence

- Complete independence reduces both *representation of joint distribution* and *inference* from  $O(2^n)$  to  $O(n)!!$
- **Unfortunately**, such complete mutual independence is very rare. Most realistic domains do not exhibit this property.
- **Fortunately**, most domains do exhibit a fair amount of conditional independence. We can exploit conditional independence for representation and inference as well.
- **Bayesian networks** do just this

# An Aside on Notation

- $\Pr(X)$  for variable  $X$  (or set of variables) refers to the *(marginal) distribution* over  $X$ .  $\Pr(X|Y)$  refers to the family of conditional distributions over  $X$ , one for each  $y \in \text{Dom}(Y)$ .
- Distinguish between  $\Pr(X)$  -- which is a distribution – and  $\Pr(x)$  or  $\Pr(\sim x)$  (or  $\Pr(x_i)$  for non-Boolean vars) -- which are numbers. Think of  $\Pr(X)$  as a function that accepts any  $x_i \in \text{Dom}(X)$  as an argument and returns  $\Pr(x_i)$ .
- Think of  $\Pr(X|Y)$  as a function that accepts any  $x_i$  and  $y_k$  and returns  $\Pr(x_i|y_k)$ . Note that  $\Pr(X|Y)$  is not a single distribution; rather it denotes the family of distributions (over  $X$ ) induced by the different  $y_k \in \text{Dom}(Y)$

# Exploiting Conditional Independence

- Consider a story:
  - If Pascal woke up too early  $E$ , Pascal probably needs coffee  $C$ ; if Pascal needs coffee, he's likely grumpy  $G$ . If he is grumpy then it's possible that the lecture won't go smoothly  $L$ . If the lecture does not go smoothly then the students will likely be sad  $S$ .



$E$  - Pascal woke up too early    $G$  - Pascal is grumpy    $S$  - Students are sad  
 $C$  - Pascal needs coffee    $L$  - The lecture did not go smoothly

# Conditional Independence



- If you learned any of  $E$ ,  $C$ ,  $G$ , or  $L$ , would your assessment of  $\Pr(S)$  change?
  - If any of these are seen to be true, you would increase  $\Pr(s)$  and decrease  $\Pr(\sim s)$ .
  - So  $S$  is *not independent* of  $E$ , or  $C$ , or  $G$ , or  $L$ .
- If you knew the value of  $L$  (true or false), would learning the value of  $E$ ,  $C$ , or  $G$  influence  $\Pr(S)$ ?
  - Influence that these factors have on  $S$  is mediated by their influence on  $L$ .
  - Students aren't sad because Pascal was grumpy, they are sad because of the lecture.
  - So  $S$  is *independent* of  $E$ ,  $C$ , and  $G$ , *given*  $L$

# Conditional Independence



- So  $S$  is *independent* of  $E$ , and  $C$ , and  $G$ , *given*  $L$
- Similarly:
  - $S$  is *independent* of  $E$ , and  $C$ , *given*  $G$
  - $G$  is *independent* of  $E$ , *given*  $C$
- This means that:

$$\Pr(S|L, \{G, C, E\}) = \Pr(S|L)$$

$$\Pr(L|G, \{C, E\}) = \Pr(L|G)$$

$$\Pr(G|C, \{E\}) = \Pr(G|C)$$

$\Pr(C|E)$  and  $\Pr(E)$  don't "simplify"

# Conditional Independence



- By the chain rule (for any instantiation of  $S \dots E$ ):

$$\Pr(S, L, G, C, E) = \Pr(S|L, G, C, E) \Pr(L|G, C, E) \Pr(G|C, E) \Pr(C|E) \Pr(E)$$


- By our independence assumptions:

$$\Pr(S, L, G, C, E) = \Pr(S|L) \Pr(L|G) \Pr(G|C) \Pr(C|E) \Pr(E)$$

- We can specify the full joint by specifying five *local conditional distributions*:

$$\Pr(S|L); \Pr(L|G); \Pr(G|C); \Pr(C|E); \text{ and } \Pr(E)$$

# Example Quantification




- Specifying the joint requires only 9 parameters (if we note that half of these are “1 minus” the others), instead of 31 for the explicit representation
  - linear in number of variables instead of exponential!
  - linear generally if dependence has a chain structure

# Inference is Easy



- Want to know  $\Pr(g)$ ? Use sum out rule:

$$\begin{aligned} P(g) &= \sum_{c_i \in \text{Dom}(C)} \Pr(g | c_i) \Pr(c_i) \\ &= \sum_{c_i \in \text{Dom}(C)} \Pr(g | c_i) \sum_{e_i \in \text{Dom}(E)} \Pr(c_i | e_i) \Pr(e_i) \end{aligned}$$



These are all terms specified in our local distributions!

# Inference is Easy



- Computing  $\Pr(g)$  in more concrete terms:

$$\Pr(c) = \Pr(c|e)\Pr(e) + \Pr(c|\sim e)\Pr(\sim e) = \cancel{0.2} \overset{0.9}{*} 0.7 + 0.5 * 0.3 = 0.78$$

$$\Pr(\sim c) = \Pr(\sim c|e)\Pr(e) + \Pr(\sim c|\sim e)\Pr(\sim e) = 0.22$$

$$\Pr(\sim c) = 1 - \Pr(c), \text{ as well}$$

$$\Pr(g) = \Pr(g|c)\Pr(c) + \Pr(g|\sim c)\Pr(\sim c) = 0.3 * 0.78 + 1.0 * 0.22 = 0.454$$

$$\Pr(\sim g) = 1 - \Pr(g) = 0.546$$