

# Sylvester-Gallai Configurations and Algebraic Complexity

**Rafael Oliveira**  
University of Waterloo

**Akash Kumar Sengupta**  
Columbia University

IIT Bombay  
February 2023

# Overview

- Introduction
  - Sylvester-Gallai Configurations
  - Previous (and current) Work
- Our Results
  - Radical Sylvester-Gallai Theorem for Cubics
  - Our Tools
  - Complete proof overview
- Conclusion & Open Problems
- Extra: SG generalization for PIT and LCCs
- Proof of Structure Theorem

# (Very brief) History

- [Sylvester 1893]**: given a finite set of points  $\mathcal{F} \subset \mathbb{R}^N$  such that:
- ▶ any line containing two points of  $\mathcal{F}$  must pass through a third.

Must  $\mathcal{F}$  be contained in a line?

# (Very brief) History

- [Sylvester 1893]**: given a finite set of points  $\mathcal{F} \subset \mathbb{R}^N$  such that:
- ▶ any line containing two points of  $\mathcal{F}$  must pass through a third.

Must  $\mathcal{F}$  be contained in a line?

**[Melchior 1940, Gallai 1944]**: YES.

# (Very brief) History

**[Sylvester 1893]**: given a finite set of points  $\mathcal{F} \subset \mathbb{R}^N$  such that:

- ▶ any line containing two points of  $\mathcal{F}$  must pass through a third.

Must  $\mathcal{F}$  be contained in a line?

**[Melchior 1940, Gallai 1944]**: YES.

Problem depends on base field.

[Folklore]: over  $\mathbb{C}$ , torsion points of elliptic curves give  
2-dimensional configurations.

**[Serre 1966]**: given a finite set of points  $\mathcal{F} \subset \mathbb{C}^N$  such that:

- ▶ any line containing two points of  $\mathcal{F}$  must pass through a third.

Must  $\mathcal{F}$  be contained in a **complex plane**?

# (Very brief) History

**[Sylvester 1893]**: given a finite set of points  $\mathcal{F} \subset \mathbb{R}^N$  such that:

- ▶ any line containing two points of  $\mathcal{F}$  must pass through a third.

Must  $\mathcal{F}$  be contained in a line?

**[Melchior 1940, Gallai 1944]**: YES.

Problem depends on base field.

[Folklore]: over  $\mathbb{C}$ , torsion points of elliptic curves give  
2-dimensional configurations.

**[Serre 1966]**: given a finite set of points  $\mathcal{F} \subset \mathbb{C}^N$  such that:

- ▶ any line containing two points of  $\mathcal{F}$  must pass through a third.

Must  $\mathcal{F}$  be contained in a **complex plane**?

**[Hirzebruch 1983]** YES, using deep results from AG.

# TCS History

Several variations and generalizations.

# TCS History

Several variations and generalizations.

- ▶ Coloured versions
  - ▶ PIT [**Dvir Shpilka 2007**] (conjectured coloured version)



# TCS History

Several variations and generalizations.

- ▶ Coloured versions
  - ▶ PIT [**Dvir Shpilka 2007**] (conjectured coloured version)
- ▶ Higher-dimensional analogs:
  - ▶ PIT [**Kayal Saraf 2009, Saxena Seshadri 2013**]

# TCS History

Several variations and generalizations.

- ▶ Coloured versions
  - ▶ PIT **[Dvir Shpilka 2007]** (conjectured coloured version)
- ▶ Higher-dimensional analogs:
  - ▶ PIT **[Kayal Saraf 2009, Saxena Seshadri 2013]**
- ▶ Robust analogs:
  - ▶ Coding Theory (LCCs) **[BDWY 2011]**
  - ▶ Reconstruction (aka tensor decomposition) **[Sinha 2016]**

# TCS History

Several variations and generalizations.

- ▶ Coloured versions
  - ▶ PIT [**Dvir Shpilka 2007**] (conjectured coloured version)
- ▶ Higher-dimensional analogs:
  - ▶ PIT [**Kayal Saraf 2009, Saxena Seshadri 2013**]
- ▶ Robust analogs:
  - ▶ Coding Theory (LCCs) [**BDWY 2011**]
  - ▶ Reconstruction (aka tensor decomposition) [**Sinha 2016**]
- ▶ Higher-degree analogs:
  - ▶ PIT [**Gupta 2014**]

# TCS History

Several variations and generalizations.

- ▶ Coloured versions
  - ▶ PIT **[Dvir Shpilka 2007]** (conjectured coloured version)
- ▶ Higher-dimensional analogs:
  - ▶ PIT **[Kayal Saraf 2009, Saxena Seshadri 2013]**
- ▶ Robust analogs:
  - ▶ Coding Theory (LCCs) **[BDWY 2011]**
  - ▶ Reconstruction (aka tensor decomposition) **[Sinha 2016]**
- ▶ Higher-degree analogs:
  - ▶ PIT **[Gupta 2014]**

Underlying theme:

Are Sylvester-Gallai type configurations always low-dimensional?

# Robust Sylvester-Gallai

## Definition (Robust linear Sylvester Gallai)

$\mathcal{F} := \{v_1, \dots, v_m\} \subset \mathbb{C}^N$  is a  $\delta$ -linear-SG configuration if for all  $i \in [m]$ , there are  $\delta(m-1)$  indices  $j$  such that:

there is  $k \neq i, j$  such that  $v_k \in \text{span}_{\mathbb{C}} \{v_i, v_j\}$ .

# Robust Sylvester-Gallai

## Definition (Robust linear Sylvester Gallai)

$\mathcal{F} := \{v_1, \dots, v_m\} \subset \mathbb{C}^N$  is a  $\delta$ -linear-SG configuration if for all  $i \in [m]$ , there are  $\delta(m-1)$  indices  $j$  such that:

there is  $k \neq i, j$  such that  $v_k \in \text{span}_{\mathbb{C}} \{v_i, v_j\}$ .

## Theorem (robust linear SG theorem – [BDWY 2011])

*If  $\mathcal{F}$  is  $\delta$ -linear-SG configuration, then  $\dim \text{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1/\delta^2)$ .*

# Robust Sylvester-Gallai

Definition (Robust linear Sylvester Gallai)

$\mathcal{F} := \{v_1, \dots, v_m\} \subset \mathbb{C}^N$  is a  $\delta$ -linear-SG configuration if for all  $i \in [m]$ , there are  $\delta(m-1)$  indices  $j$  such that:

there is  $k \neq i, j$  such that  $v_k \in \text{span}_{\mathbb{C}} \{v_i, v_j\}$ .

Theorem (robust linear SG theorem – **[BDWY 2011]**)

If  $\mathcal{F}$  is  $\delta$ -linear-SG configuration, then  $\dim \text{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1/\delta^2)$ .

► Improved to  $O(1/\delta)$  by **[Dvir Saraf Wigderson 2014]**.

# Why Should I Care?

- ▶ Mathematicians & complexity theorists:

It's a **structural** study of **cancellations/relations** (syzygies).



# Why Should I Care?

- ▶ Mathematicians & complexity theorists:

It's a **structural** study of **cancellations/relations** (**syzygies**).

Cancellations in SG configurations make them quite complex!

# Mayr-Meyer



[Mayr Meyer 1982]: “cancellations in algebraic geometry are EXPSPACE hard”

[Brownawell 1987, Kollar 1988]: “radical cancellations” are in PSPACE.

# Where are the cancellations?

- ▶  $\mathcal{F} = \{v_1, \dots, v_m\} \subset \mathbb{R}^2$  is a SG configuration if for all  $i, j \in [m]$ , there is  $k \neq i, j$  such that  $v_i, v_j, v_k$  colinear.

# Where are the cancellations?

- ▶  $\mathcal{F} = \{v_1, \dots, v_m\} \subset \mathbb{R}^2$  is a SG configuration if for all  $i, j \in [m]$ , there is  $k \neq i, j$  such that  $v_i, v_j, v_k$  colinear.
- ▶ **Duality:**  $\mathcal{F} = \{\ell_1, \dots, \ell_m\} \subset \mathbb{R}[x, y]_1$  is a SG configuration if for all  $i, j \in [m]$ , there is  $k \neq i, j$  such that  $\ell_k \in (\ell_i, \ell_j)$ .

# Where are the cancellations?

- ▶  $\mathcal{F} = \{v_1, \dots, v_m\} \subset \mathbb{R}^2$  is a SG configuration if for all  $i, j \in [m]$ , there is  $k \neq i, j$  such that  $v_i, v_j, v_k$  colinear.
- ▶ **Duality:**  $\mathcal{F} = \{\ell_1, \dots, \ell_m\} \subset \mathbb{R}[x, y]_1$  is a SG configuration if for all  $i, j \in [m]$ , there is  $k \neq i, j$  such that  $\ell_k \in (\ell_i, \ell_j)$ .

1.  $\ell_k \in (\ell_i, \ell_j) \Leftrightarrow \exists \alpha_i, \alpha_j, \alpha_k \in \mathbb{R}$  such that

$$\alpha_i \ell_i + \alpha_j \ell_j + \alpha_k \ell_k = 0$$

# Where are the cancellations?

- ▶  $\mathcal{F} = \{v_1, \dots, v_m\} \subset \mathbb{R}^2$  is a SG configuration if for all  $i, j \in [m]$ , there is  $k \neq i, j$  such that  $v_i, v_j, v_k$  colinear.
- ▶ **Duality**:  $\mathcal{F} = \{\ell_1, \dots, \ell_m\} \subset \mathbb{R}[x, y]_1$  is a SG configuration if for all  $i, j \in [m]$ , there is  $k \neq i, j$  such that  $\ell_k \in (\ell_i, \ell_j)$ .

1.  $\ell_k \in (\ell_i, \ell_j) \Leftrightarrow \exists \alpha_i, \alpha_j, \alpha_k \in \mathbb{R}$  such that

$$\alpha_i \ell_i + \alpha_j \ell_j + \alpha_k \ell_k = 0$$

2. Are these relations enough to show that  $\dim\langle \mathcal{F} \rangle = 1$ ?

- ▶ **(Non-linear) Generalization [Gupta 2014]**:
  - ▶  $\mathcal{F} = \{F_1, \dots, F_m\} \subset \mathbb{C}[x_1, \dots, x_N]$  is a SG configuration if for all  $i, j \in [m]$ , there is  $k \neq i, j$  such that

$$F_k \in \text{rad}(F_i, F_j)$$

# Generalization – geometrically



# General conjecture

Definition (Radical Sylvester Gallai – [Gupta 2014])

$\mathcal{F} = \{F_1, \dots, F_m\} \subset \mathbb{C}[x_1, \dots, x_N]$  is a  $d$ -radical-SG config. if:

1.  $F_i$  irreducible for all  $i \in [m]$
2.  $\deg(F_i) \leq d$  for all  $i \in [m]$  (low degree)
3.  $F_i \notin (F_j)$  for  $i \neq j$  (“distinct”)
4. for all  $i, j$ , there is  $k \neq i, j$  such that (SG dependency)

$$F_k \in \text{rad}(F_i, F_j)$$



# General conjecture

Definition (Radical Sylvester Gallai – **[Gupta 2014]**)

$\mathcal{F} = \{F_1, \dots, F_m\} \subset \mathbb{C}[x_1, \dots, x_N]$  is a  $d$ -radical-SG config. if:

1.  $F_i$  irreducible for all  $i \in [m]$
2.  $\deg(F_i) \leq d$  for all  $i \in [m]$  (low degree)
3.  $F_i \notin (F_j)$  for  $i \neq j$  (“distinct”)
4. for all  $i, j$ , there is  $k \neq i, j$  such that (SG dependency)

$$F_k \in \text{rad}(F_i, F_j)$$

Conjecture (**[Gupta 2014]**)

There is  $\lambda : \mathbb{N} \rightarrow \mathbb{N}$  s.t. if  $\mathcal{F}$  is a  $d$ -radical-SG configuration, then

$$\text{tr-deg}(\mathcal{F}) \leq \lambda(d).$$

Informally: must every SG configuration be in “few variables”?

# General conjecture

Definition (Radical Sylvester Gallai – [Gupta 2014])

$\mathcal{F} = \{F_1, \dots, F_m\} \subset \mathbb{C}[x_1, \dots, x_N]$  is a  $d$ -radical-SG config. if:

1.  $F_i$  irreducible for all  $i \in [m]$
2.  $\deg(F_i) \leq d$  for all  $i \in [m]$  (low degree)
3.  $F_i \notin (F_j)$  for  $i \neq j$  (“distinct”)
4. for all  $i, j$ , there is  $k \neq i, j$  such that (SG dependency)

$$F_k \in \text{rad}(F_i, F_j)$$

## Conjecture

There is  $\lambda : \mathbb{N} \rightarrow \mathbb{N}$  s.t. if  $\mathcal{F}$  is a  $d$ -radical-SG configuration, then

$$\dim \text{span}_{\mathbb{C}} \{\mathcal{F}\} \leq \lambda(d).$$

# Previous (and current) Works

Theorem (Linear SG – **[Hirzebruch 1983]**)

*If  $\mathcal{F}$  is 1-radical-SG configuration, then  $\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} \leq 2$ .*

Another proof given by **[Dvir Saraf Wigderson 2014]**

# Previous (and current) Works

Theorem (Linear SG – **[Hirzebruch 1983]**)

*If  $\mathcal{F}$  is 1-radical-SG configuration, then  $\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} \leq 2$ .*

Another proof given by **[Dvir Saraf Wigderson 2014]**

Theorem (Quadratic radical SG theorem – **[Shpilka 2020]**)

*If  $\mathcal{F}$  is 2-radical-SG configuration, then  $\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1)$ .*

# Previous (and current) Works

Theorem (Linear SG – **[Hirzebruch 1983]**)

*If  $\mathcal{F}$  is 1-radical-SG configuration, then  $\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} \leq 2$ .*

Another proof given by **[Dvir Saraf Wigderson 2014]**

Theorem (Quadratic radical SG theorem – **[Shpilka 2020]**)

*If  $\mathcal{F}$  is 2-radical-SG configuration, then  $\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1)$ .*

Theorem (Cubic radical SG theorem – **[O. Sengupta 2022]**)

*If  $\mathcal{F}$  is 3-radical-SG configuration, then  $\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1)$ .*

- Introduction
    - Sylvester-Gallai Configurations
    - Previous (and current) Work
  - Our Results
    - Radical Sylvester-Gallai Theorem for Cubics
    - Our Tools
    - Complete proof overview
  - Conclusion & Open Problems
  - Extra: SG generalization for PIT and LCCs
  - Proof of Structure Theorem
-

# Quadratic Case (1-page Amir)

Theorem (Quadratic radical SG theorem – [Shpilka 2020])

*If  $\mathcal{F}$  is 2-radical-SG configuration, then  $\dim \text{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1)$ .*

# Quadratic Case (1-page Amir)

Theorem (Quadratic radical SG theorem – [Shpilka 2020])

*If  $\mathcal{F}$  is 2-radical-SG configuration, then  $\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1)$ .*

Proof outline:

- ▶ **Structure theorem:** how can  $F_k \in \operatorname{rad}(F_i, F_j)$ ?
  1.  $F_k \in \operatorname{span}_{\mathbb{C}} \{F_i, F_j\}$



# Quadratic Case (1-page Amir)

Theorem (Quadratic radical SG theorem – [Shpilka 2020])

*If  $\mathcal{F}$  is 2-radical-SG configuration, then  $\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1)$ .*

Proof outline:

► **Structure theorem:** how can  $F_k \in \operatorname{rad}(F_i, F_j)$ ?

1.  $F_k \in \operatorname{span}_{\mathbb{C}} \{F_i, F_j\}$
2.  $\ell^2 \in \operatorname{span}_{\mathbb{C}} \{F_i, F_j\}$

# Quadratic Case (1-page Amir)

Theorem (Quadratic radical SG theorem – [Shpilka 2020])

*If  $\mathcal{F}$  is 2-radical-SG configuration, then  $\dim \text{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1)$ .*

Proof outline:

► **Structure theorem:** how can  $F_k \in \text{rad}(F_i, F_j)$ ?

1.  $F_k \in \text{span}_{\mathbb{C}} \{F_i, F_j\}$
2.  $\ell^2 \in \text{span}_{\mathbb{C}} \{F_i, F_j\}$
3.  $F_i, F_j, F_k \in (x, y)$  for some linear  $x, y$

# Quadratic Case (1-page Amir)

Theorem (Quadratic radical SG theorem – [Shpilka 2020])

*If  $\mathcal{F}$  is 2-radical-SG configuration, then  $\dim \text{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1)$ .*

Proof outline:

- ▶ **Structure theorem:** how can  $F_k \in \text{rad}(F_i, F_j)$ ?
  1.  $F_k \in \text{span}_{\mathbb{C}} \{F_i, F_j\}$
  2.  $\ell^2 \in \text{span}_{\mathbb{C}} \{F_i, F_j\}$
  3.  $F_i, F_j, F_k \in (x, y)$  for some linear  $x, y$
- ▶ Main idea: "linearize" the configuration  
quadratics **not** robust **linear** configuration  $\Rightarrow$  must **look alike**.

# Quadratic Case (1-page Amir)

Theorem (Quadratic radical SG theorem – [Shpilka 2020])

*If  $\mathcal{F}$  is 2-radical-SG configuration, then  $\dim \text{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1)$ .*

Proof outline:

- ▶ **Structure theorem:** how can  $F_k \in \text{rad}(F_i, F_j)$ ?
  1.  $F_k \in \text{span}_{\mathbb{C}} \{F_i, F_j\}$
  2.  $\ell^2 \in \text{span}_{\mathbb{C}} \{F_i, F_j\}$
  3.  $F_i, F_j, F_k \in (x, y)$  for some linear  $x, y$
- ▶ Main idea: "linearize" the configuration  
quadratics **not** robust **linear** configuration  $\Rightarrow$  must **look alike**.

Upshot: **non-linear** SG dependencies involve special **linear** forms.

# Quadratic Case (1-page Amir)

Theorem (Quadratic radical SG theorem – [Shpilka 2020])

*If  $\mathcal{F}$  is 2-radical-SG configuration, then  $\dim \text{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1)$ .*

Proof outline:

- ▶ **Structure theorem:** how can  $F_k \in \text{rad}(F_i, F_j)$ ?
  1.  $F_k \in \text{span}_{\mathbb{C}} \{F_i, F_j\}$
  2.  $\ell^2 \in \text{span}_{\mathbb{C}} \{F_i, F_j\}$
  3.  $F_i, F_j, F_k \in (x, y)$  for some linear  $x, y$
- ▶ Main idea: "linearize" the configuration  
quadratics **not** robust **linear** configuration  $\Rightarrow$  must **look alike**.
- ▶ Extract **linear** Sylvester-Gallai configuration from remaining linear forms  
(combinatorially involved)

# Some Notation

- Graded rings:  $R = \bigoplus_{d \geq 0} R_d$  such that

$$R_i R_j \subset R_{i+j}$$

$R_d :=$  set of elements of degree  $d$ .

# Some Notation

- ▶ Graded rings:  $R = \bigoplus_{d \geq 0} R_d$  such that

$$R_i R_j \subset R_{i+j}$$

$R_d :=$  set of elements of degree  $d$ .

- ▶ Polynomial ring graded by degree

# Some Notation

- ▶ Graded rings:  $R = \bigoplus_{d \geq 0} R_d$  such that

$$R_i R_j \subset R_{i+j}$$

$R_d :=$  set of elements of degree  $d$ .

- ▶ Polynomial ring graded by degree
- ▶ Given graded vector space  $V = V_1 + \cdots + V_d$  can construct graded algebra  $\mathbb{C}[V]$



# How can we induct?

Approach to induct generalizes [Shpilka 2020] in several ways.

# How can we induct?

Approach to induct generalizes [Shpilka 2020] in several ways.

- **Observation:** if there is vector space  $V = V_1 + V_2$  such that  $\mathcal{F} \subset \mathbb{C}[V]$ , then

$$\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} \leq (\dim V)^3.$$

Enough to construct small algebra  $\mathbb{C}[V]$  with  $\dim V = O(1)$ .

# How can we induct?

Approach to induct generalizes [Shpilka 2020] in several ways.

- ▶ **Observation:** if there is vector space  $V = V_1 + V_2$  such that  $\mathcal{F} \subset \mathbb{C}[V]$ , then

$$\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} \leq (\dim V)^3.$$

Enough to construct small algebra  $\mathbb{C}[V]$  with  $\dim V = O(1)$ .

- ▶ How to reduce degree? (from 3  $\rightarrow$  2) Let  $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$ .
  - ▶ it may not be true that  $\mathcal{F}_1 \cup \mathcal{F}_2$  is a 2-radical-SG configuration

# How can we induct?

Approach to induct generalizes [Shpilka 2020] in several ways.

- ▶ **Observation:** if there is vector space  $V = V_1 + V_2$  such that  $\mathcal{F} \subset \mathbb{C}[V]$ , then

$$\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} \leq (\dim V)^3.$$

Enough to construct small algebra  $\mathbb{C}[V]$  with  $\dim V = O(1)$ .

- ▶ How to reduce degree? (from 3  $\rightarrow$  2) Let  $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$ .
  - ▶ it may not be true that  $\mathcal{F}_1 \cup \mathcal{F}_2$  is a 2-radical-SG configuration
  - ▶ if could prove
    1. there is small  $V = V_1 + V_2$  s.t.  $\mathcal{F}_3 \subset \mathbb{C}[V]$
    2. we could solve 2-radical-SG over the algebra  $\mathbb{C}[V]$

then done!

# How can we induct?

Approach to induct generalizes [Shpilka 2020] in several ways.

- ▶ **Observation:** if there is vector space  $V = V_1 + V_2$  such that  $\mathcal{F} \subset \mathbb{C}[V]$ , then

$$\dim \text{span}_{\mathbb{C}} \{\mathcal{F}\} \leq (\dim V)^3.$$

Enough to construct small algebra  $\mathbb{C}[V]$  with  $\dim V = O(1)$ .

- ▶ How to reduce degree? (from 3  $\rightarrow$  2) Let  $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$ .
  - ▶ it may not be true that  $\mathcal{F}_1 \cup \mathcal{F}_2$  is a 2-radical-SG configuration
  - ▶ if could prove
    1. there is small  $V = V_1 + V_2$  s.t.  $\mathcal{F}_3 \subset \mathbb{C}[V]$
    2. we could solve 2-radical-SG over the algebra  $\mathbb{C}[V]$then done!
- ▶ Can we do both? YES!

Need a lot of new tools!

# Inductive radical SG problem

Original radical SG configuration:

**Definition (Radical Sylvester Gallai )**

$\mathcal{F} = \{F_1, \dots, F_m\} \subset \mathbb{C}[x_1, \dots, x_N]$  is a  $d$ -radical-SG config. if:

1.  $F_i$  irreducible for all  $i \in [m]$
2.  $\deg(F_i) \leq d$  for all  $i \in [m]$  (low degree)
3.  $F_i \notin (F_j)$  for  $i \neq j$  (“distinct”)
4. for all  $i, j$ , there is  $k \neq i, j$  such that (SG dependency)

$$F_k \in \text{rad}(F_i, F_j) \Leftrightarrow |\mathcal{F} \cap \text{rad}(F_i, F_j)| \geq 3$$

# Inductive radical SG problem

Inductive radical SG configuration:

Definition (Radical Sylvester Gallai over algebra)

Let  $V = V_1 + \cdots + V_d$ .  $\mathcal{F} = \{F_1, \dots, F_m\} \subset \mathbb{C}[x_1, \dots, x_N]$  is a  $(d, V)$ -radical-SG configuration if:

1.  $F_i$  irreducible for all  $i \in [m]$
2.  $\deg(F_i) \leq d$  for all  $i \in [m]$  (low degree)
3.  $F_i \notin (F_j)$  for  $i \neq j$  (“distinct”)

# Inductive radical SG problem

Inductive radical SG configuration:

Definition (Radical Sylvester Gallai over algebra)

Let  $V = V_1 + \cdots + V_d$ .  $\mathcal{F} = \{F_1, \dots, F_m\} \subset \mathbb{C}[x_1, \dots, x_N]$  is a  $(d, V)$ -radical-SG configuration if:

1.  $F_i$  irreducible for all  $i \in [m]$
2.  $\deg(F_i) \leq d$  for all  $i \in [m]$  (low degree)
3.  $F_i \notin (F_j)$  for  $i \neq j$  (“distinct”)
4. for all  $i, j$  (SG dependency over algebra)

$$|\mathcal{F} \cap \text{rad}(F_i, F_j)| \geq 3 \quad \text{or} \quad \text{rad}(F_i, F_j) \cap \mathbb{C}[V] \not\subset (F_i) \cup (F_j)$$

Upshot: can have pairs  $i, j$  with no dependence in  $\mathcal{F}$ , but it has to have dependence in algebra  $\mathbb{C}[V]$ .



- Introduction
    - Sylvester-Gallai Configurations
    - Previous (and current) Work
  - Our Results
    - Radical Sylvester-Gallai Theorem for Cubics
    - Our Tools
    - Complete proof overview
  - Conclusion & Open Problems
  - Extra: SG generalization for PIT and LCCs
  - Proof of Structure Theorem
-

# Algebras

- ▶ Not all algebras are created equally...
  1. Best algebra: polynomial rings (commutative and free)

# Algebras

- ▶ Not all algebras are created equally...
  1. Best algebra: polynomial rings (commutative and free)
  2. Could  $\mathcal{F}_3$  be in a small sub-polynomial ring?

# Algebras

- ▶ Not all algebras are created equally...
  1. Best algebra: polynomial rings (commutative and free)
  2. Could  $\mathcal{F}_3$  be in a small sub-polynomial ring?
  3. May not be possible in our case:

$$F = x(y_1z_1 + y_2z_2 + \cdots + y_nz_n) + uw^2 \in \mathcal{F}_3$$

# Algebras

- ▶ Not all algebras are created equally...
  1. Best algebra: polynomial rings (commutative and free)
  2. Could  $\mathcal{F}_3$  be in a small sub-polynomial ring?
  3. May not be possible in our case:

$$F = x(y_1z_1 + y_2z_2 + \cdots + y_nz_n) + uw^2 \in \mathcal{F}_3$$

- ▶ What is next best thing?

# Algebras

- ▶ Not all algebras are created equally...
  1. Best algebra: polynomial rings (commutative and free)
  2. Could  $\mathcal{F}_3$  be in a small sub-polynomial ring?
  3. May not be possible in our case:

$$F = x(y_1z_1 + y_2z_2 + \cdots + y_nz_n) + uw^2 \in \mathcal{F}_3$$

- ▶ What is next best thing?
  - ▶ Subalgebras that are **isomorphic** to a polynomial ring AND **behave well** with  $\mathbb{C}[x_1, \dots, x_N]$

# Algebras

- ▶ Not all algebras are created equally...
  1. Best algebra: polynomial rings (commutative and free)
  2. Could  $\mathcal{F}_3$  be in a small sub-polynomial ring?
  3. May not be possible in our case:

$$F = x(y_1z_1 + y_2z_2 + \cdots + y_nz_n) + uw^2 \in \mathcal{F}_3$$

- ▶ What is next best thing?
  - ▶ Subalgebras that are **isomorphic** to a polynomial ring AND **behave well** with  $\mathbb{C}[x_1, \dots, x_N]$
  - ▶ Algebras generated by prime sequences!  
Key properties: **Regular Sequence** & **Intersection flatness**

# Algebras

- ▶ Not all algebras are created equally...
  1. Best algebra: polynomial rings (commutative and free)
  2. Could  $\mathcal{F}_3$  be in a small sub-polynomial ring?
  3. May not be possible in our case:

$$F = x(y_1z_1 + y_2z_2 + \cdots + y_nz_n) + uw^2 \in \mathcal{F}_3$$

- ▶ What is next best thing?
  - ▶ Subalgebras that are **isomorphic** to a polynomial ring AND **behave well** with  $\mathbb{C}[x_1, \dots, x_N]$
  - ▶ Algebras generated by prime sequences!  
Key properties: **Regular Sequence** & **Intersection flatness**
    1. Regular sequence  $\Rightarrow$  “free as polynomial ring”



# Algebras

- ▶ Not all algebras are created equally...
  1. Best algebra: polynomial rings (commutative and free)
  2. Could  $\mathcal{F}_3$  be in a small sub-polynomial ring?
  3. May not be possible in our case:

$$F = x(y_1z_1 + y_2z_2 + \cdots + y_nz_n) + uw^2 \in \mathcal{F}_3$$

- ▶ What is next best thing?
  - ▶ Subalgebras that are **isomorphic** to a polynomial ring AND **behave well** with  $\mathbb{C}[x_1, \dots, x_N]$
  - ▶ Algebras generated by prime sequences!  
Key properties: **Regular Sequence** & **Intersection flatness**
    1. Regular sequence  $\Rightarrow$  “free as polynomial ring”
    2. Intersection flatness  $\Rightarrow$  behaves nicely with  $\mathbb{C}[x_1, \dots, x_N]$   
Primes in the small subalgebra are also primes in  $\mathbb{C}[x_1, \dots, x_N]$

# Wide Ananyan-Hochster Algebras

- Suppose I have an algebra  $\mathbb{C}[F_1, \dots, F_k]$  of low degree polynomials which is not nice

Can I convert it into a nice algebra without blowing up the size too much?

# Wide Ananyan-Hochster Algebras

- ▶ Suppose I have an algebra  $\mathbb{C}[F_1, \dots, F_k]$  of low degree polynomials which is not nice

Can I convert it into a nice algebra without blowing up the size too much?

- ▶ **[Ananyan Hochster 2020]** construct such algebras (and much more!)

1. Basic idea: if  $V = V_1 + V_2$  is such that ANY  $Q \in V_2$  has

$$\text{rank}(Q) \geq \dim V + 3$$

then  $\mathbb{C}[V]$  is a nice algebra ( $V$  nice vector space).

# Wide Ananyan-Hochster Algebras

- ▶ Suppose I have an algebra  $\mathbb{C}[F_1, \dots, F_k]$  of low degree polynomials which is not nice

Can I convert it into a nice algebra without blowing up the size too much?

- ▶ **[Ananyan Hochster 2020]** construct such algebras (and much more!)

1. Basic idea: if  $V = V_1 + V_2$  is such that ANY  $Q \in V_2$  has

$$\text{rank}(Q) \geq \dim V + 3$$

then  $\mathbb{C}[V]$  is a nice algebra ( $V$  nice vector space).

- ▶ In **[O. Sengupta 2022]** we build upon this to construct **wide Ananyan-Hochster algebras**

1. generated by prime sequences (or better)
2. robust to “small increases”

# Our approach

1. Solve  $(2, V)$ -radical-SG problem

Proposition (**[O. Sengupta 2022]**)

*If  $V$  is wide AH vector space and  $\mathcal{F}$  is  $(2, V)$ -radical-SG configuration, then*

$$\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1 + (\dim V)^2)$$

# Our approach

1. Solve  $(2, V)$ -radical-SG problem

Proposition (**[O. Sengupta 2022]**)

*If  $V$  is wide AH vector space and  $\mathcal{F}$  is  $(2, V)$ -radical-SG configuration, then*

$$\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1 + (\dim V)^2)$$

Generalizes **[Shpilka 2020]** main result.

# Our approach

1. Solve  $(2, V)$ -radical-SG problem

**Proposition ([O. Sengupta 2022])**

*If  $V$  is wide AH vector space and  $\mathcal{F}$  is  $(2, V)$ -radical-SG configuration, then*

$$\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}\} = O(1 + (\dim V)^2)$$

2. Now we need to construct  $V$  wide such that  $\mathcal{F}_3 \subset \mathbb{C}[V]$ .

# Structure Theorems

Theorem (Structure theorem for cubics [O. Sengupta 2022])

Let  $F, G$  be irreducible homogeneous cubics. One of the following must hold:

1.  $(F, G)$  is radical
2.  $(F, G) \subset (x, y)$  for  $x, y$  linear forms
3.  $(F, G) \subset (Q, x)$  for  $Q$  irreducible quadratic and  $x$  linear
4.  $xy^2 \in \text{span}_{\mathbb{C}} \{F, G\}$  for  $x, y$  linear forms
5.  $(F, G) \subset I_{md}$  where  $I_{md}$  cuts out *variety of minimal degree*

Example of variety of minimal degree:

(twisted cubic)

$$\begin{pmatrix} x & y & z \\ y & z & w \end{pmatrix} \mapsto (y^2 - xz, z^2 - yw, xw - yz)$$



# More Structure Theorems

In addition to the above, several new structure theorems which hold for general degree  $d$

# More Structure Theorems

In addition to the above, several new structure theorems which hold for general degree  $d$

- ▶ Discriminant lemma (decide radical or not)
  - ▶ generalizes fact that discriminant of univariate polynomial  $p(x)$  is zero  $\Leftrightarrow p(x)$  has multiple roots
  - ▶ quantitative bounds when combined with wide AH algebras

Key property: **Cohen-Macaulayness**

# More Structure Theorems

In addition to the above, several new structure theorems which hold for general degree  $d$

- ▶ Discriminant lemma (decide radical or not)
  - ▶ generalizes fact that discriminant of univariate polynomial  $p(x)$  is zero  $\Leftrightarrow p(x)$  has multiple roots
  - ▶ quantitative bounds when combined with wide AH algebras

Key property: **Cohen-Macaulayness**

- ▶ **Transfer principle**: generalize several properties of polynomial rings to wide AH algebras
  - ▶ elimination theorems in wide AH algebras
  - ▶ primality and reducedness criteria in AH algebras
  - ▶ more...

Key property: **Intersection Flatness**

# Proof overview

- ▶  $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$  our 3-radical-SG configuration
- ▶ Solved  $(2, V)$ -radical-SG problem over  $V$  low dimensional wide AH vector space

# Proof overview

- ▶  $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$  our 3-radical-SG configuration
- ▶ Solved  $(2, V)$ -radical-SG problem over  $V$  low dimensional wide AH vector space
- ▶ Need to prove  $\mathcal{F}_3 \subset \mathbb{C}[V]$  for some small wide AH vector space  $V$ 
  1. If  $\mathcal{F}_3$  is a  $\delta$ -linear-SG configuration then  $\dim \operatorname{span}_{\mathbb{C}} \{\mathcal{F}_3\} = O(1)$ .  
Apply our wide AH process to  $\operatorname{span}_{\mathbb{C}} \{\mathcal{F}_3\}$  to get  $V$ .

# Proof overview

- ▶  $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$  our 3-radical-SG configuration
- ▶ Solved  $(2, V)$ -radical-SG problem over  $V$  low dimensional wide AH vector space
- ▶ Need to prove  $\mathcal{F}_3 \subset \mathbb{C}[V]$  for some small wide AH vector space  $V$

1. If  $\mathcal{F}_3$  is a  $\delta$ -linear-SG configuration then  $\dim \text{span}_{\mathbb{C}} \{\mathcal{F}_3\} = O(1)$ .

Apply our wide AH process to  $\text{span}_{\mathbb{C}} \{\mathcal{F}_3\}$  to get  $V$ .

2.  $\mathcal{F}_3$  not  $\delta$ -linear-SG configuration, then there are cubics  $C_1, C_2, C_3$  such that **most**  $F_i \in \mathcal{F}_3$  is such that  $(F_i, C_j)$  **not-radical** ( $j \in [3]$ ).

Most of the wide vector space  $V$  comes from  $C_1, C_2, C_3$ .

With a bit of work, transform  $\mathcal{F}$  into a  $(2, V)$ -radical-SG configuration.

- Introduction
    - Sylvester-Gallai Configurations
    - Previous (and current) Work
  - Our Results
    - Radical Sylvester-Gallai Theorem for Cubics
    - Our Tools
    - Complete proof overview
  - Conclusion & Open Problems
  - Extra: SG generalization for PIT and LCCs
  - Proof of Structure Theorem
-

# Why is 3 important?

Challenges in degree 3 similar to challenges in general case

- ▶ geometry is more complex
  - ▶ need more general structural lemmas
  - ▶ structure theorem for cubics is more involved than for quadratics



# Why is 3 important?

Challenges in degree 3 similar to challenges in general case

- ▶ geometry is more complex
  - ▶ need more general structural lemmas
  - ▶ structure theorem for cubics is more involved than for quadratics
- ▶ it may not be possible to “linearize” the configuration

# Why is 3 important?

Challenges in degree 3 similar to challenges in general case

- ▶ geometry is more complex
  - ▶ need more general structural lemmas
  - ▶ structure theorem for cubics is more involved than for quadratics
- ▶ it may not be possible to “linearize” the configuration
- ▶ if we want **principled** approach, need to devise an **inductive** version of SG
  - ▶ reducing from cubic to quadratic is harder than from quadratic to linear

# Why is 3 important?

Challenges in degree 3 similar to challenges in general case

- ▶ geometry is more complex
  - ▶ need more general structural lemmas
  - ▶ structure theorem for cubics is more involved than for quadratics
- ▶ it may not be possible to “linearize” the configuration
- ▶ if we want **principled** approach, need to devise an **inductive** version of SG
  - ▶ reducing from cubic to quadratic is harder than from quadratic to linear

All of the above (and a little bit more) in **[O. Sengupta 2022]**!

# Conclusion

- ▶ Proved 3-radical-SG conjecture is true.

# Conclusion

- ▶ Proved 3-radical-SG conjecture is true.
- ▶ Inductive, generalizable SG problem (SG over algebra)  
In previous versions, unclear how to solve SG inductively.

# Conclusion

- ▶ Proved 3-radical-SG conjecture is true.
- ▶ Inductive, generalizable SG problem (SG over algebra)  
In previous versions, unclear how to solve SG inductively.
- ▶ Introduced several new algebro-geometric techniques:
  1. **wide AH algebras**
    - ▶ subalgebras “like subpolynomial rings”
    - ▶ robust to small augmentations
  2. **discriminant-based** reducedness testing & quantitative bounds
  3. **transfer principle**:  
polynomial rings  $\rightarrow$  algebras generated by prime sequences
  4. Exploration of **Cohen-Macaulayness** in SG configurations
  5. Structure theorem for intersection of cubics

# Open Questions

Open Question (Radical Sylvester-Gallai over an algebra)

*There is  $\lambda : \mathbb{N}^2 \rightarrow \mathbb{N}$  such that if  $\mathcal{F}$  is a  $(d, V)$ -radical-SG configuration, then*

$$\dim \operatorname{span}_{\mathbb{C}} \{ \mathcal{F} \} \leq \lambda(d, \dim V).$$

Several variants – robust, coloured, higher-codimensional... this is just the beginning of the rabbit hole.

# Open Questions

## Open Question (Radical Sylvester-Gallai over an algebra)

*There is  $\lambda : \mathbb{N}^2 \rightarrow \mathbb{N}$  such that if  $\mathcal{F}$  is a  $(d, V)$ -radical-SG configuration, then*

$$\dim \operatorname{span}_{\mathbb{C}} \{ \mathcal{F} \} \leq \lambda(d, \dim V).$$

Several variants – robust, coloured, higher-codimensional... this is just the beginning of the rabbit hole.

Radical Sylvester-Gallai seems instrumental first step towards main conjecture of **[Gupta 2014]**, as in **[Shpilka 2020, Peleg Shpilka 2020]**.



# Open Questions

## Open Question (Radical Sylvester-Gallai over an algebra)

*There is  $\lambda : \mathbb{N}^2 \rightarrow \mathbb{N}$  such that if  $\mathcal{F}$  is a  $(d, V)$ -radical-SG configuration, then*

$$\dim \operatorname{span}_{\mathbb{C}} \{ \mathcal{F} \} \leq \lambda(d, \dim V).$$

Several variants – robust, coloured, higher-codimensional... this is just the beginning of the rabbit hole.

Radical Sylvester-Gallai seems instrumental first step towards main conjecture of **[Gupta 2014]**, as in **[Shpilka 2020, Peleg Shpilka 2020]**.

More generally: can we parametrize cancellations in algebra?

# Future Directions

A sneak peek into the rabbit hole:

## Open Question (Complexity theory for Algebraic Geometry)

*Can we pin down the complexity of basic algebro-geometric questions?*

- ▶ *primary decomposition*
- ▶ *radical ideal membership*
- ▶ *projective dimension*
- ▶ *free resolutions*

# Future Directions

A sneak peek into the rabbit hole:

## Open Question (Complexity theory for Algebraic Geometry)

*Can we pin down the complexity of basic algebro-geometric questions?*

- ▶ *primary decomposition*
- ▶ *radical ideal membership*
- ▶ *projective dimension*
- ▶ *free resolutions*

[Ananyan Hochster 2020] gives us **upper bound** (non-explicit) on parametrization of cancellations/relations (and in the above problems).

- ▶ can we get explicit (and eventually tight) parametrizations?
- ▶ important special cases as complexity classes?

# References I



Ananyan, Tigran and Hochster, Melvin (2020)

Small subalgebras of polynomial rings and Stillman's conjecture  
*Journal of the American Mathematical Society*



Barak, Boaz and Dvir, Zeev and Yehudayoff, Amir and Wigderson, Avi (2011)

Rank Bounds for Design Matrices with Applications to Combinatorial Geometry and Locally Correctable Codes  
*Forty-Third Annual ACM Symposium on Theory of Computing*



Brownawell, W. Dale. (1987)

Bounds for the degrees in the Nullstellensatz.  
*Annals of Mathematics* 126.3



Colliot-Thélène, J-L and Sansuc, J-J and Swinnerton-Dyer, P (1987)

Intersections of two quadrics and Châtelet surfaces. I.  
*Journal für die reine und angewandte Mathematik*

# References II



Dvir, Zeev and Shpilka, Amir (2007)

Locally decodable codes with two queries and polynomial identity testing for depth 3 circuits

[SIAM Journal on Computing](#)



Dvir, Zeev and Saraf, Shubhangi and Wigderson, Avi (2014)

Improved rank bounds for design matrices and a new proof of Kelly's theorem

[Forum of Mathematics, Sigma](#)



Garg, Abhibhav and Oliveira, Rafael and Sengupta, Akash K (2022)

Robust Radical Sylvester-Gallai Theorem for Quadratics

[SoCG 2022](#)



Gallai, Tibor (1944)

Solution of problem 4065

[American Mathematical Monthly](#)

# References III



Gupta, Ankit (2014)

Algebraic Geometric Techniques for Depth-4 PIT & Sylvester-Gallai Conjectures for Varieties.

[Electron. Colloquium Comput. Complex.](#)



Hansen, Sten (1966)

A generalization of a theorem of Sylvester on the lines determined by a finite point set

[Mathematica Scandinavica](#)



Hodge, William Vallance Douglas and Pedoe, Daniel (1994)

Methods of Algebraic Geometry: Volume 2

[Cambridge University Press](#)



Hirzebruch, (1983)

Arrangements of lines and algebraic surfaces

[Arithmetic and Geometry](#), pages 113–140.

# References IV



Kayal, Neeraj and Saraf, Shubhangi (2009)

Blackbox polynomial identity testing for depth 3 circuits

[50th Annual IEEE Symposium on Foundations of Computer Science](#)



Kollár, János. (1988)

Sharp effective nullstellensatz.

[Journal of the American Mathematical Society](#)



Mayr, Ernst and Meyer, Albert (1982)

The complexity of the word problems for commutative semigroups and polynomial ideals

[Advances in Mathematics](#), 305-329



Melchior, Eberhard (1940)

Über vielseitigkeit der projektiven ebene

[Deutsche Math](#), volume 5, pages 461–275

# References V



Oliveira, Rafael and Sengupta, Akash K (2022)

Radical Sylvester-Gallai theorem for cubics

[Manuscript](#)



Peleg, Shir and Shpilka, Amir (2020)

Polynomial time deterministic identity testing algorithm for  $\Sigma^{[3]}\Pi\Sigma\Pi^{[2]}$  circuits via Edelstein-Kelly type theorem for quadratic polynomials

[53rd Annual ACM SIGACT Symposium on Theory of Computing](#)



Saxena, Nitin and Seshadhri, Comandur

From Sylvester-Gallai configurations to rank bounds: Improved blackbox identity test for depth-3 circuits

[Journal of the ACM \(JACM\)](#)



Sinha, Gaurav (2016)

Reconstruction of Real Depth-3 Circuits with Top Fan-In 2

[31st Conference on Computational Complexity](#)



# References VI



Serre, Jean-Pierre (1966)

Advanced problem 5359

[Amer. Math. Monthly](#), volume 73, number 1, page 89



Shpilka, Amir (2020)

Sylvester-Gallai type theorems for quadratic polynomials

[Discrete Analysis](#)



Sylvester, James Joseph (1893)

Mathematical question 11851

[Educational Times](#), volume 59, number 98, page 256

- Introduction
  - Sylvester-Gallai Configurations
  - Previous (and current) Work
- Our Results
  - Radical Sylvester-Gallai Theorem for Cubics
  - Our Tools
  - Complete proof overview
- Conclusion & Open Problems
- Extra: SG generalization for PIT and LCCs
- Proof of Structure Theorem

# SG configurations in PIT and Reconstruction

- ▶ PIT/Reconstruction break down into two cases:
  1. SG circuits: where a lot of cancellations/relations can happen. In this case the circuit may not be unique/have less structure (hard case)
  2. non-SG circuits: few relations can happen. This case is easier, since we can “isolate” the gates.

# Proving structure theorem

- ▶ Look at primary decomposition (minimal primes + multiplicity)
  1.  $(F, G)$  is Cohen-Macaulay  $\Rightarrow$  unmixed (and much more)

# Proving structure theorem

- ▶ Look at primary decomposition (minimal primes + multiplicity)
  1.  $(F, G)$  is **Cohen-Macaulay**  $\Rightarrow$  **unmixed** (and much more)
  2. From primary decomposition:

$$\deg(F, G) = \sum_{\mathfrak{p}} m(\mathfrak{p}) \cdot \deg(\mathfrak{p})$$

# Proving structure theorem

- ▶ Look at primary decomposition (minimal primes + multiplicity)
  1.  $(F, G)$  is **Cohen-Macaulay**  $\Rightarrow$  **unmixed** (and much more)
  2. From primary decomposition:

$$\deg(F, G) = \sum_{\mathfrak{p}} m(\mathfrak{p}) \cdot \deg(\mathfrak{p})$$

3.  $\deg(F, G) = 9$ , since  $F, G$  cubics with  $\gcd(F, G) = 1$

# Proving structure theorem

- ▶ Look at primary decomposition (minimal primes + multiplicity)
  1.  $(F, G)$  is **Cohen-Macaulay**  $\Rightarrow$  **unmixed** (and much more)
  2. From primary decomposition:

$$\deg(F, G) = \sum_{\mathfrak{p}} m(\mathfrak{p}) \cdot \deg(\mathfrak{p})$$

3.  $\deg(F, G) = 9$ , since  $F, G$  cubics with  $\gcd(F, G) = 1$
4. if  $m(\mathfrak{p}) = 1$  for all  $\mathfrak{p}$  then  $(F, G)$  is radical

# Proving structure theorem

- ▶ Look at primary decomposition (minimal primes + multiplicity)
  1.  $(F, G)$  is **Cohen-Macaulay**  $\Rightarrow$  **unmixed** (and much more)
  2. From primary decomposition:

$$\deg(F, G) = \sum_{\mathfrak{p}} m(\mathfrak{p}) \cdot \deg(\mathfrak{p})$$

3.  $\deg(F, G) = 9$ , since  $F, G$  cubics with  $\gcd(F, G) = 1$
4. if  $m(\mathfrak{p}) = 1$  for all  $\mathfrak{p}$  then  $(F, G)$  is radical
5. if  $(F, G) \subset (x, y)$  we are done, so assume this is not the case.  
Then  $\deg(\mathfrak{p}) \geq 2$  for all  $\mathfrak{p}$ .



# Proving structure theorem

- ▶ Look at primary decomposition (minimal primes + multiplicity)
  1.  $(F, G)$  is **Cohen-Macaulay**  $\Rightarrow$  **unmixed** (and much more)
  2. From primary decomposition:

$$\deg(F, G) = \sum_{\mathfrak{p}} m(\mathfrak{p}) \cdot \deg(\mathfrak{p})$$

3.  $\deg(F, G) = 9$ , since  $F, G$  cubics with  $\gcd(F, G) = 1$
4. if  $m(\mathfrak{p}) = 1$  for all  $\mathfrak{p}$  then  $(F, G)$  is radical
5. if  $(F, G) \subset (x, y)$  we are done, so assume this is not the case.  
Then  $\deg(\mathfrak{p}) \geq 2$  for all  $\mathfrak{p}$ .
6.  $9 = \{2, 3\} \cdot d + \text{"stuff of degree } \geq 2\text{"}$  so  $d \in \{2, 3\}$

# Proving structure theorem

- ▶ Look at primary decomposition (minimal primes + multiplicity)
  1.  $(F, G)$  is **Cohen-Macaulay**  $\Rightarrow$  **unmixed** (and much more)
  2. From primary decomposition:

$$\deg(F, G) = \sum_{\mathfrak{p}} m(\mathfrak{p}) \cdot \deg(\mathfrak{p})$$

3.  $\deg(F, G) = 9$ , since  $F, G$  cubics with  $\gcd(F, G) = 1$
4. if  $m(\mathfrak{p}) = 1$  for all  $\mathfrak{p}$  then  $(F, G)$  is radical
5. if  $(F, G) \subset (x, y)$  we are done, so assume this is not the case.  
Then  $\deg(\mathfrak{p}) \geq 2$  for all  $\mathfrak{p}$ .
6.  $9 = \{2, 3\} \cdot d + \text{"stuff of degree } \geq 2\text{"}$  so  $d \in \{2, 3\}$
7.  $d = 2 \Rightarrow \mathfrak{p} = (Q, x)$  for  $Q$  quadratic and  $x$  linear

# Proving structure theorem

- ▶ Look at primary decomposition (minimal primes + multiplicity)
  1.  $(F, G)$  is **Cohen-Macaulay**  $\Rightarrow$  **unmixed** (and much more)
  2. From primary decomposition:

$$\deg(F, G) = \sum_{\mathfrak{p}} m(\mathfrak{p}) \cdot \deg(\mathfrak{p})$$

3.  $\deg(F, G) = 9$ , since  $F, G$  cubics with  $\gcd(F, G) = 1$
4. if  $m(\mathfrak{p}) = 1$  for all  $\mathfrak{p}$  then  $(F, G)$  is radical
5. if  $(F, G) \subset (x, y)$  we are done, so assume this is not the case.  
Then  $\deg(\mathfrak{p}) \geq 2$  for all  $\mathfrak{p}$ .
6.  $9 = \{2, 3\} \cdot d + \text{"stuff of degree } \geq 2\text{"}$  so  $d \in \{2, 3\}$
7.  $d = 2 \Rightarrow \mathfrak{p} = (Q, x)$  for  $Q$  quadratic and  $x$  linear
8.  $d = 3$  and  $(F, G)$  **degenerate**  $\Rightarrow \mathfrak{p} = (F, x)$  for  $x$  linear

# Proving structure theorem

- ▶ Look at primary decomposition (minimal primes + multiplicity)
  1.  $(F, G)$  is **Cohen-Macaulay**  $\Rightarrow$  **unmixed** (and much more)
  2. From primary decomposition:

$$\deg(F, G) = \sum_{\mathfrak{p}} m(\mathfrak{p}) \cdot \deg(\mathfrak{p})$$

3.  $\deg(F, G) = 9$ , since  $F, G$  cubics with  $\gcd(F, G) = 1$
4. if  $m(\mathfrak{p}) = 1$  for all  $\mathfrak{p}$  then  $(F, G)$  is radical
5. if  $(F, G) \subset (x, y)$  we are done, so assume this is not the case.  
Then  $\deg(\mathfrak{p}) \geq 2$  for all  $\mathfrak{p}$ .
6.  $9 = \{2, 3\} \cdot d + \text{"stuff of degree } \geq 2\text{"}$  so  $d \in \{2, 3\}$
7.  $d = 2 \Rightarrow \mathfrak{p} = (Q, x)$  for  $Q$  quadratic and  $x$  linear
8.  $d = 3$  and  $(F, G)$  **degenerate**  $\Rightarrow \mathfrak{p} = (F, x)$  for  $x$  linear
9.  $d = 3$  and  $(F, G)$  **non-degenerate**  $\Rightarrow \mathfrak{p}$  defines variety of minimal degree